

ABSTRACT

Precision Theory for LHC/FCC: New Results for the Five Point Function and Interface between KKMC-hh and MG5_aMC@NLO

Yang Liu, Ph.D.

Advisor: B. F. L. Ward, Ph.D.

The development of large colliders provides us with the opportunity to discover the fundamental particles in nature and explore the interactions among them. The Standard Model (SM) of particle physics reflects our best knowledge of elementary particles and their interactions at present, which is formulated by a gauge quantum field theory with gauge symmetry $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$. With the discovery of the Higgs boson, the era of the sub-1% precision on processes such as Z and W production is approaching us. In order to achieve the 1% theoretical precision tag, we have to take radiative corrections into account and develop more precise Monte Carlo generators.

In this dissertation, we first developed the computer realization of the magic spinor product method in loop integrals proposed by B. F. L. Ward to evaluate the general five-point function numerically. The result from magic spinor product method agrees with that from LoopTools overall. Additionally, we also developed an approach to achieve the next-to-the-leading order QCD and the electroweak (EW) exact $O(\alpha_s \otimes \alpha^2 L)$ corrections, interfacing MG5_aMC@NLO with KKMC-hh by merging their LHE

files. By comparing the results of the Drell-Yan process obtained by KKMC-hh, MG5_aMC@NLO and KKMC-hh interfaced with MG5_aMC@NLO , at $\sqrt{s} = 13$ TeV with the ATLAS cuts on the Z/γ^* production and decay to lepton pairs, respectively, we find that the results derived from KKMC-hh interfaced with MG5_aMC@NLO would generate enhancements from those derived from MG5_aMC@NLO, which is due to the EW corrections provided by KKMC-hh.

Precision Theory for LHC/FCC: New Results for the Five Point Function and
Interface between KKMC-hh and MG5_aMC@NLO

by

Yang Liu, B.S., M.A., M.S.,

A Dissertation

Approved by the Department of Physics

Dwight P. Russell, Ph.D., Interim Chairperson

Submitted to the Graduate Faculty of
Baylor University in Partial Fulfillment of the
Requirements for the Degree
of
Doctor of Philosophy

Approved by the Dissertation Committee

Bennie F. L. Ward, Ph.D., Chairperson

Gerald B. Cleaver, Ph.D.

Kenichi Hatakeyama, Ph.D.

Mark Sepanski, Ph.D.

Anzhong Wang, Ph.D.

Accepted by the Graduate School
May 2021

J. Larry Lyon, Ph.D., Dean

Copyright © 2021 by Yang Liu

All rights reserved

TABLE OF CONTENTS

LIST OF FIGURES	viii
LIST OF TABLES.....	x
ACKNOWLEDGMENTS	xi
DEDICATION.....	xii
 CHAPTER ONE	
Standard Model of Electroweak Interaction.....	1
1.1 Gauge Invariance	3
1.1.1 Abelian Gauge Invariance: Quantum Electrodynamics.....	3
1.1.2 Non-Abelian Gauge Invariance	5
1.2 Spontaneous Symmetry Breaking.....	7
1.3 The Higgs Mechanism	11
1.3.1 The Higgs Mechanism in the Abelian Theory	11
1.3.2 The Higgs Mechanism in the Non-Abelian Theory	15
1.4 Standard Model of the Electroweak Interactions	18
1.4.1 The General Ideas for Building Spontaneous Broken Gauge Theories	18
1.4.2 The Glashow-Weinberg-Salam Theory.....	21
 CHAPTER TWO	
Techniques for the Calculation of Electroweak Radiative Corrections at the One-Loop Level	29
2.1 The Model.....	30
2.2 Renormalization in the Electroweak SM	35
2.2.1 Renormalization Constants and Counterterms	37
2.2.2 Renormalization Conditions	38
2.2.3 Explicit Form of Renormalization Constants.....	43
2.3 One-Loop Integrals.....	46
2.3.1 Scalar One-loop Integrals for $N \leq 4$	46
2.3.2 Tensor Integral Reduction	52
2.4 Example for One-loop Radiative Correction Calculations	62
 CHAPTER THREE	
Spinor Techniques	67
3.1 Introduction	67
3.2 The CALKUL Approach.....	69
3.3 The "Chinese Magic" Polarization Scheme.....	77
3.4 Kleiss and Stirling Spinor Technique	85

CHAPTER FOUR	
Numerical Computations for the Five Point Function	89
4.1 Magic Spinor Product Approach in Loop Integrals	90
4.2 Numerical Results for the Five-Point Function E_0	100
CHAPTER FIVE	
Quantum Chromodynaics	107
5.1 Introduction to Quantum Chromodynamics	107
5.2 Renormalization Group Equation and Asymptotic Freedom	117
5.2.1 Renormalization Group Equation	117
5.2.2 Asymptotic Freedom in QCD	123
CHAPTER SIX	
Yennie-Frautschi-Suura Resummation	129
6.1 Resummation of Virtual Photon Radiative Corrections	130
6.2 Resummation of Real Photon Radiative Corrections	134
6.3 Details of Infrared Factors	137
6.4 Details of Noninfrared Virtual Photon Terms	139
6.5 Details of Noninfrared Real Photon Terms	141
CHAPTER SEVEN	
CEEX and KKMC-hh	144
7.1 Amplitudes for Exclusive Exponentiation	145
7.1.1 Master Formula	146
7.1.2 Pure Virtual Corrections	149
7.1.3 One Real Photon with Virtual Corrections	151
7.1.4 Two Real Photons with Virtul Corrections	153
7.1.5 Three Real Photons	158
7.2 Amplitudes for Coherent Exclusive Exponentiation	160
7.2.1 Master Formula	160
7.2.2 IR Structure in CEEX	166
7.2.3 Narrow Neutral Resonance in CEEX	176
7.2.4 Virtual Corrections, No Photons	186
7.2.5 One Real Photon	190
7.2.6 Two Real Photons	198
7.3 Relations between CEEX and EEX	210
7.4 Monte Carlo Algorithm	212
7.4.1 Weights and Distributions	213
7.4.2 Phase-space Reorganization	220
7.4.3 MC generation of the FSR photon momenta	222
7.4.4 Common IR Boundary For ISR and FSR	231
7.4.5 Entire MC Algorithm Top-to-Bottom	238
CHAPTER EIGHT	
Interface between KKMC-hh and MG5_aMC@NLO	240
8.1 Overview of Madgraph_aMC@NLO	240
8.2 Interfacing KKMC-hh with MG5_aMC@NLO	242

8.3 Results for Interfacing KKMC-hh with MG5_aMC@NLO.....	247
CHAPTER NINE	
Overall Summary	253
APPENDICES.....	255
APPENDIX A	
Feynman Rules of the Electroweak SM	256
APPENDIX B	
Feynman Rules of Quantum Chromodynamics	271
APPENDIX C	
The $SU(3)$ Group	277
APPENDIX D	
The Global Positioning of Spin GPS Scheme.....	280
APPENDIX E	
The Drell-Yan Process.....	284
BIBLIOGRAPHY.....	296

LIST OF FIGURES

Figure 1.1. The elementary particles of the SM	2
Figure 1.2. The potential $V(\phi)$ of the scalar field ϕ in the case $\mu^2 > 0$ (left) and $\mu^2 < 0$ (right)	8
Figure 4.1. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 0^\circ$	101
Figure 4.2. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 15^\circ$	101
Figure 4.3. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 30^\circ$	102
Figure 4.4. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 45^\circ$	102
Figure 4.5. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 60^\circ$	102
Figure 4.6. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 75^\circ$	103
Figure 4.7. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 90^\circ$	103
Figure 4.8. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 105^\circ$	103
Figure 4.9. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 120^\circ$	104
Figure 4.10. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 135^\circ$	104
Figure 4.11. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 150^\circ$	104
Figure 4.12. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} =$ 500 GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 165^\circ$	105

Figure 4.13. Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_2} = 91$ GeV and $\theta'_1 = 180^\circ$	105
Figure 8.1. Muon transverse momentum distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.	249
Figure 8.2. Muon pseudorapidity distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.	250
Figure 8.3. Dimuon transverse momentum distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.	251
Figure 8.4. Dimuon invariant mass distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.	252
Figure 8.5. Dimuon rapidity distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.	252

LIST OF TABLES

Table 8.1. Cross Sections obtained by MG5_aMC@NLO \otimes KKMC-hh, KKMC-hh, MG5_aMC@NLO, respectively.....	248
--	-----

ACKNOWLEDGMENTS

Firstly, I would like to thank my mentor Dr. Bennie F. L. Ward for his guidance and support during the years I have been studying and working under his supervision. I owe him my sincerest gratitude.

Special thanks to Dr. Mark Sepanski for the assistance he has given me. I would like to express my gratitude for his enlightening discussions regarding graduate math classes.

I would like to acknowledge my committee members, Dr. Gerald Cleaver, Dr. Kenichi Hatakeyama, Dr. Mark Sepanski and Dr. Anzhong Wang for their time and patience.

I would also like to thank the Physics Department of Baylor University and our staff members Mrs. Chava Baker and Mrs. Marian Nunn-Graves for their dedication to duty.

To my loving wife Natalie Rose Cowan-Liu

CHAPTER ONE

Standard Model of Electroweak Interaction

The standard model (SM) of particle physics represents our best understanding of elementary particles and their interactions. It is one of the most successful theories, because its predictions are confirmed with exceptional precision in many experiments. The SM is a gauge quantum field theory with gauge symmetry $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ [1–16]. $SU(3)_C$ is the gauge symmetry of quantum chromodynamics (QCD), the theory describing strong interactions, and $SU(2)_L \otimes U(1)_Y$ is the gauge symmetry of electroweak interactions. The SM is constructed by invariance under Poincaré group (translations, rotations and Lorentz boosts) and renormalizability. The constituents of matter are spin-1/2 particles (fermions), 6 leptons and 6 quarks that pair up to transform under $SU(2)_L$. The interactions of SM are mediated by spin-1 particles—gauge bosons. The elementary particles of the SM are listed in Figure 1.1.

If an elementary particle carries the charge of a certain force, it is involved with the corresponding interaction. Quarks carry color charge (red, green, blue) and interact through the strong force mediated by massless gluons. The gluon has eight different states, which carries a combination of color and anti-color charge in each state (color $SU(3)$ octet). The up, charm and top quarks carry a fractional electric charge of $2/3e$, while down, strange and bottom quarks carry a fractional electric charge of $-1/3e$. The charged leptons (electron, muon and tau) have an integer charge of $-e$. All electrically charged particles participate in the electromagnetic interaction mediated by the massless photon. Each charged lepton is paired with a neutral




Figure 1.1: The elementary particles of the SM

lepton (electron-, muon-, and tau-neutrino) with extremely low mass. All the particles participate in the weak interactions since they all carry an isospin, of which the z -component is either $+1/2$ (u, c, t-quark and neutrinos) or $-1/2$ (d, s, b-quark and charged leptons). The weak interactions are mediated by the neutral Z or the electrically charged W^\pm vector bosons. The masses of elementary particles in SM are acquired through the interactions with Higgs fields.

In this chapter, we aim to give a brief introduction of the standard model of electroweak interactions [17–19]. We will introduce gauge invariance first. The spontaneous symmetry breaking and Higgs mechanism will be discussed next. Last we will review the construction for the Lagrangian of the electroweak interactions.

1.1 Gauge Invariance

1.1.1 Abelian Gauge Invariance: Quantum Electrodynamics

Gauge theories are built with internal symmetries. For example, consider the $U(1)$ group of phase transformations of a free massive fermion field $\psi(x)$:

$$\psi(x) \rightarrow e^{-i\alpha}\psi(x), \quad (1.1)$$

where α is an arbitrary phase parameter. The corresponding Lagrangian density

$$L(x) = \bar{\psi}(x)(i\not{\partial} - m)\psi(x) \quad (1.2)$$

is invariant under these transformations. According to the Nöther theorem, this symmetry leads to a conserved current,

$$j_\mu(x) = \bar{\psi}(x)\gamma_\mu\psi(x), \quad \partial^\mu j_\mu = 0. \quad (1.3)$$

The conserved charge, namely, generator of the $U(1)$ symmetry group, can be written as an integral over the charge density:

$$Q = \int d^3x j_0(x). \quad (1.4)$$

The invariance of the Lagrangian (1.2) under phase rotation indicates that the phase parameter α has no physical significance so that it could be chosen arbitrarily. It is unnatural to select a uniquely fixed α over all of the space-time, and it would be more natural to choose α locally,

$$\psi(x) \rightarrow e^{-i\alpha(x)}\psi(x), \quad (1.5)$$

where α depends on space-time in an arbitrary way. However, this modification brings a new problem. The Lagrangian (1.2) is no more invariant under the local phase rotations (1.5), because the derivative $\partial_\mu\psi(x)$ is transformed under phase rotation

(1.5) by

$$\partial_\mu\psi(x) \rightarrow e^{-i\alpha(x)}\partial_\mu\psi(x) - ie^{i\alpha(x)}\partial_\mu\alpha(x)\psi(x). \quad (1.6)$$

To solve this problem, We need to introduce a covariant derivative D_μ , which has the property that $D\psi$ transforms under phase rotations like ψ :

$$D_\mu\psi(x) \rightarrow e^{-i\alpha(x)}D_\mu\psi(x). \quad (1.7)$$

Such a covariant derivative can only be introduced if there exists another field, a vector field A_μ which interacts with the spinor field ψ . The covariant derivative D_μ is chosen as

$$D_\mu = \partial_\mu + igA_\mu \quad (1.8)$$

where g is an arbitrary coupling constant, and A_μ transforms under a local phase transformation (gauge transformation) as follows:

$$A_\mu(x) \rightarrow A_\mu(x) + \frac{1}{g}\partial_\mu\alpha(x). \quad (1.9)$$

We could easily verify that covariant (1.8) satisfies the requirement (1.7). Thus the invariance of the Lagrangian (1.2) under gauge transformations is recovered after replacing ∂_μ with D_μ . However, we must add the kinetic term of the A_μ field for consistency, which must be gauge invariant in itself (only involving the gauge-invariant field strength).

$$F_{\mu\nu} = \partial_\nu A_\mu - \partial_\mu A_\nu. \quad (1.10)$$

Therefore we obtain the Lagrangian which is invariant under gauge transformation

$$L = \bar{\psi}(i\not{D} - m)\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (1.11)$$

If we identify the spinor field ψ with electron field, the vector field A_μ with photon field and replace g by e (electric charge), we obtain the Lagrangian for quantum electrodynamics (QED). Note that only the minimal coupling of the photon field to the electron field of the type $e\bar{\psi}\gamma_\mu\psi A^\mu$ is allowed due to the requirement of local gauge invariance. Furthermore a mass term for the photon field of the type $m^2 A_\mu A^\mu$ is forbidden to arise in the Lagrangian.

1.1.2 Non-Abelian Gauge Invariance

The idea of non-Abelian gauge theories was formulated by Yang and Mills [20] in 1954. We successfully constructed the QED Lagrangian by imposing the local gauge invariance, $U(1)$. This success encourages us to extend the gauge symmetry from an Abelian gauge case to a non-Abelian case. We will take the isospin symmetry as an example to formulate of the Non-Abelian gauge theories. The Lagrangian for the free protons and neutrons is

$$L = \bar{N}(i\not{\partial} - m)N \quad (1.12)$$

where N represents the isospinor $(p, n)^T$. It is invariant under $SU(2)$ transformations

$$N \rightarrow e^{-i\alpha\sigma/2}N \quad (1.13)$$

where $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ ($\sigma \equiv$ Pauli matrices). and α is an arbitrary phase vector.

The isotropic spin currents $\bar{N}\gamma_\mu\frac{1}{2}\sigma_i N$ are conserved. And the associated charges $T_i = \int d^3x \bar{N}\gamma_\mu\frac{1}{2}\sigma_i N$ generate the algebra of $SU(2)$:

$$[T_i, T_j] = i\epsilon_{ijk}T_k. \quad (1.14)$$

Note that the gauge transformations involving nondiagonal Pauli matrices σ_1, σ_2 result in the mixing between p and n states. However, this would not be a problem,

since there would be no physical difference between proton and neutron for the isospin symmetry. Our selection for p or n totally depends on convention. Therefore it would be natural to redefine p and n locally in an arbitrary way, namely, we require the invariance under non-Abelian gauge transformations:

$$\begin{pmatrix} p \\ n \end{pmatrix} \rightarrow e^{-i\alpha(x)\sigma/2} \begin{pmatrix} p \\ n \end{pmatrix} \quad (1.15)$$

where $\alpha(x)$ is an arbitrary spacetime dependent phase vector. We will encounter the same problem as in the Abelian case discussed above: the derivative $\partial_\mu N$ which occurs in the Lagrangian (1.12) will not transform under local gauge transformations as N itself, and we have to define a proper covariant derivative. To achieve this goal, we introduce a triplet of vector gauge fields W_μ^i , which transforms under an infinitesimal gauge transformation as follows:

$$W_\mu^i \rightarrow W_\mu^i + \epsilon_{ijk}\alpha^j W_\mu^k + \frac{1}{g}\partial_\mu\alpha^i \quad (1.16)$$

This transformation is analogous to eq. (1.9). The second term shows the local rotation of the W^i in the isotropic space. The covariant derivative is defined as

$$D_\mu = \partial_\mu + igW_\mu \quad (1.17)$$

where $W_\mu = \frac{1}{2}\sigma W_\mu$. The Lagrangian of the system is

$$L = \bar{N}(i\cancel{\partial} - m)N - \frac{1}{4}G_{\mu\nu}^i G_i^{\mu\nu}, \quad (1.18)$$

where the $G_{\mu\nu}^i$ are the field strength tensors of the vector fields:

$$G_{\mu\nu}^i = \partial_\nu W_\mu^i - \partial_\mu W_\nu^i - g\epsilon^{ijk}W_{j\nu}W_{k\mu} \quad (1.19)$$

This approach can be generalized to the case where an arbitrary gauge group and an arbitrary fermion representation are involved. The only changes are as follows:

(a) Replace the isospin matrices σ with the corresponding matrices describing the transformation properties of the fermions under the gauge group.

(b) Replace ϵ_{ijk} with the structure constants f_{ijk} of the gauge group.

Note that the fermions can transform as an arbitrary representation of the gauge group, while the vector gauge fields must transform according to the adjoint representation. In the non-Abelian gauge theory, the vector fields interact with each other directly ("they are charged"). This is not the case in the Abelian theory where the vector field is neutral. The Lagrangian of the non-Abelian gauge theory (1.18) describes the interactions of massive fermions with massless gauge bosons.

1.2 Spontaneous Symmetry Breaking

It is known that the weak interactions are mediated by massive vector bosons Z and W^\pm . However, we have seen above that the non-Abelian gauge invariance requires the gauge bosons to be massless. This means we must seek another possibility to introduce masses for the gauge fields in a more subtle way, such that the local gauge invariance is preserved. This can be realized by generating the gauge boson masses via a spontaneous breaking of the gauge symmetry.

Usually the equations of motion for a physical system are symmetric under some symmetry transformations, however, the ground state of the systems is not. For example, the Hamiltonian for an infinitely extended ferromagnet is invariant under rotations in space. However, the ground state breaks the rotational symmetry since the individual spins are always aligned in an arbitrary direction. Similar situations arise in the field theory often, and we will discuss several examples of spontaneous breaking field theories.




Figure 1.2: The potential $V(\phi)$ of the scalar field ϕ in the case $\mu^2 > 0$ (left) and $\mu^2 < 0$ (right)

The simplest example to exhibit the phenomenon of spontaneous symmetry breaking is the ϕ^4 theory. Consider the Lagrangian

$$L = \frac{1}{2}(\partial^\mu \phi \partial_\mu \phi) - \frac{1}{2}\mu^2 \phi^2 - \frac{1}{4}\lambda \phi^4. \quad (1.20)$$

The Lagrangian (1.20) is invariant under the parity transformation P , defined by $\phi \xrightarrow{P} -\phi$. The ground state (vacuum state $|0\rangle$) of the theory is the one where ϕ vanishes everywhere. It is invariant under the parity transformation: $P|0\rangle = |0\rangle$.

If the parameter $\mu^2 > 0$, the potential $V(\phi) = \frac{1}{2}\mu^2 \phi^2 + \frac{1}{4}\lambda \phi^4$ has a unique minimum at $\phi = 0$, as shown in Figure 1.2, which corresponds to the vacuum state.

If the parameter $\mu^2 < 0$, the situation is that of a spontaneously broken symmetry. The potential

$$V(\phi) = -\frac{1}{2}|\mu^2|\phi^2 + \frac{1}{4}|\lambda|\phi^4, \quad (1.21)$$

shown in Figure 1.2, has minima at

$$\langle 0 | \phi | 0 \rangle = \pm \sqrt{-\mu^2/|\lambda|} \equiv \pm v, \quad (1.22)$$

which corresponds to two possible ground states. These vacuums are not invariant under parity transformation since $v \neq -v$. Thus the parity invariance is spontaneously

broken. It is useful to define a new field ϕ' , for which $\langle 0 | \phi' | 0 \rangle = 0$, i.e. $\phi' = \phi - v$. In terms of ϕ we have

$$L = \frac{1}{2}(\partial^\mu \phi' \partial_\mu \phi') - \mu^2 \phi'^2 - \frac{1}{4}\lambda \phi'^4 - \lambda v \phi^3 + \text{const.} \quad (1.23)$$

We see that the Lagrangian (1.20) with $\mu^2 < 0$ describes self-interacting scalar particles with mass $\sqrt{2}|\mu|$.

Next, let us consider the theory of a complex scalar field $\phi = \frac{\sqrt{2}}{2}(\phi_1 + i\phi_2)$.

Consider the Lagrangian

$$\begin{aligned} L &= \frac{1}{2}(\partial^\mu \phi^* \partial_\mu \phi) - \frac{1}{2}\mu^2 \phi^2 - \frac{1}{4}\lambda \phi^4 \\ &= \frac{1}{2}\partial^\mu \phi_1 \partial_\mu \phi_1 + \frac{1}{2}\partial^\mu \phi_2 \partial_\mu \phi_2 - \frac{1}{2}\mu^2(\phi_1^2 + \phi_2^2) - \frac{1}{4}\lambda(\phi_1^2 + \phi_2^2)^2. \end{aligned} \quad (1.24)$$

It is invariant under the phase transformation $\phi \rightarrow e^{-i\theta}\phi$. For $\mu^2 > 0$ this Lagrangian describes a self-interacting scalar complex field of mass μ .

Assume we choose $\mu^2 < 0$ now, then the potential $V(\phi)$ has a minimum at $(\phi_1^2 + \phi_2^2) = 2|\phi|^2 = -\mu^2/\lambda$. So the minimum of the potential occurs along a circle of radius $\sqrt{-\mu^2/\lambda}$ around the origin. Because we could pick any point on the circle as the vacuum state, we are now dealing with an infinite number of possible vacua. Let us take an arbitrary point on the circle as the vacuum, described by the coordinates $v = (v_1, v_2)$. Since the Lagrangian is invariant under phase transformations, we could let this point lie on the positive real axis, namely, $v = (\sqrt{-\mu^2/\lambda})$, which implies $\langle 0 | \phi_1 | 0 \rangle = \sqrt{-\mu^2/\lambda}$, $\langle 0 | \phi_2 | 0 \rangle = 0$.

Now we define $\phi'_1 = \phi_1 - \langle 0 | \phi_1 | 0 \rangle$, then the Lagrangian (1.24) becomes

$$L = \frac{1}{2}\partial^\mu \phi'_1 \partial_\mu \phi'_1 + \frac{1}{2}\partial^\mu \phi_2 \partial_\mu \phi_2 + \mu^2 \phi_1'^2 - \frac{1}{2}\lambda v \phi'_1(\phi_1'^2 + \phi_2^2) - \frac{\lambda}{4}(\phi_1'^2 + \phi_2^2)^2, \quad (1.25)$$

where $v = \sqrt{-\mu^2/\lambda}$. From the Lagrangian (1.25), we see that the field ϕ_1 describes a particle of mass $\sqrt{2|\mu|}$, but ϕ_2 is still massless.

The system described above gives an example of the Goldstone theorem [21–25]: for every spontaneously broken continuous symmetry, the theory contains massless Goldstone bosons (spin-0).

The example discussed above exhibits an invariance under the global gauge group $U(1)$ which is isomorphic to $O(2)$. It could be generalized to involve the gauge group $O(n)$. Let us consider the Lagrangian

$$L = \frac{1}{2}(\partial^\mu \phi_i \partial_\mu \phi_i) - \frac{1}{2}\mu^2 \phi_i \phi_i - \frac{1}{4}\lambda(\phi_i \phi_i)^2, i = 1, 2, \dots, n;$$

$\phi_i \equiv$ real scalar field; summation over i . (1.26)

This Lagrangian is invariant under the group $O(n)$. For $\mu^2 < 0$ the minimum of the potential is at $v = \sqrt{-\mu^2/\lambda}$. The potential $V(\phi)$ exhibits the minimum at $\phi_i \phi_i = -\mu^2/\lambda$, i.e. it arises on the n -dimensional sphere of radius $\sqrt{-\mu^2/\lambda}$ in the n -dimensional space defined by the fields ϕ_i . Because of the $O(n)$ -invariance of the Lagrangian we could select the coordinates of fields so that the vacuum expectation value of the field vector ϕ_i is defined as follows:

$$\langle 0 | \phi | 0 \rangle = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ v \end{bmatrix}. \tag{1.27}$$

Note that the first $(n-1)$ components of $\langle 0 | \phi | 0 \rangle$ are zero, the vacuum remains invariant under the subgroup $O(n-1)$. We see that the vacuum expectation value $\langle 0 | \phi | 0 \rangle$ breaks the $O(n)$ -invariance in a particular way. Let us see the point (1.27)

in the n -dimensional space of the fields ϕ_i . There are $(n - 1)$ linearly independent directions to leave this point, but to stay on the sphere which minimizes the potential. Therefore there must exist $(n - 1)$ massless Goldstone bosons, according to the Goldstone theorem. So the Lagrangian (1.26) describes a massive field of mass $\sqrt{-2\mu^2}$ and $(n - 1)$ massless Goldstone bosons.

The group $O(n)$ has $\frac{1}{2}n(n - 1)$ generators, while the subgroup $O(n - 1)$ has $\frac{1}{2}(n - 1)(n - 2)$ generators. This means $(n - 1)$ generators of $O(n)$ do not leave the vacuum invariant. On the other hand, we have $(n - 1)$ massless Goldstone bosons, namely, the number of massless Goldstone bosons is equal to the number of generators which are broken spontaneously. This feature is a special property of the $O(n)$ model we have discussed above, but a general feature of spontaneously broken theories involving scalar fields. The number of Goldstone bosons corresponds always to the number of the spontaneously broken generators as a consequence of the general Goldstone theorem [26].

1.3 The Higgs Mechanism

As we discussed above, the local gauge invariance requires gauge bosons to be massless only. However, in reality, the observed gauge bosons of weak interactions Z and W^\pm are massive. In order to reconcile this contradiction, we need to incorporate Higgs mechanism [27–29] into the gauge theory, by which spontaneous symmetry breaking generates a mass for a gauge boson.

1.3.1 The Higgs Mechanism in the Abelian Theory

Let us consider the Lagrangian

$$L = \partial^\mu \phi^* \partial_\mu \phi - \mu^2 \phi^* \phi - \lambda(\phi^* \phi)^2, \quad (1.28)$$

which is invariant under the global gauge transformations $\phi \rightarrow e^{-i\alpha}\phi$, i.e., $U(1)$ gauge group. However, we need invariance under the local gauge transformations $\phi \rightarrow e^{-i\alpha(x)}\phi$. In order to achieve this goal, we need to introduce a gauge field A_μ .

Repeating the procedure outlined in Subsection 1.1.1, we arrive at

$$L = (D^\mu \phi)^* D_\mu \phi - \mu^2 \phi^* \phi - \lambda (\phi^* \phi)^2 - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad (1.29)$$

where $F_{\mu\nu} = \partial_\nu A_\mu - \partial_\mu A_\nu$, and $D_\mu = \partial_\mu + igA_\mu$.

We note that the various fields transform under local gauge transformations as follows:

$$\begin{aligned} \phi(x) &\rightarrow e^{-i\alpha(x)}\phi(x), \\ A_\mu(x) &\rightarrow A_\mu(x) + \frac{1}{g}\partial_\mu\alpha(x). \end{aligned} \quad (1.30)$$

For $\mu^2 > 0$ the Lagrangian (1.29) describes the system of a massive scalar field, coupled to a massless gauge field A_μ . If we let $g = e$ (electric charge), then we are dealing with the scalar electrodynamics.

For $\mu^2 < 0$ the gauge symmetry is spontaneously broken, as discussed in Section 1.2. We have known that the Lagrangian (1.28) describes a massive scalar field, accompanied by a massless Goldstone by following the approach outlined in section 1.2. Next, we will investigate what will happen in case of the gauge invariant Lagrangian (1.29). Let us make the substitution

$$\phi_1 = \phi'_1 + \langle 0 | \phi_1 | 0 \rangle = \phi_1 + v, v = \sqrt{-\mu^2/\lambda} \quad (1.31)$$

in eq. (1.29). We therefore expand the scalar field $\phi(x)$ around the vacuum expectation value $\langle 0 | \phi | 0 \rangle$ and arrive at

$$L = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial^\mu \phi_1 \partial_\mu \phi_1 + \frac{1}{2} \partial^\mu \phi_2 \partial_\mu \phi_2 + \frac{1}{2} g^2 v^2 A_\mu A^\mu - gv A_\mu \partial^\mu \phi_2. \quad (1.32)$$

We find that following new terms arise:

$$\frac{1}{2}g^2v^2A_\mu A^\mu, \quad (1.33)$$

$$-gvA_\mu\partial^\mu\phi_2. \quad (1.34)$$

The term (1.33) could be interpreted as the gauge boson mass term where the mass $m_A^2 = g^2v^2$ arises from the nonvanishing vacuum expectation value of ϕ . However, the interpretation of the term (1.34) is sort of vague, since it mixes the gauge field with the Goldstone boson ϕ . In order to clarify this condition, let us consider the gauge transformation $\phi \rightarrow e^{-i\alpha}\phi$ in terms of ϕ'_1 and ϕ_2 .

For an infinitesimal parameter α we have

$$\begin{aligned} \phi &\rightarrow (1 - i\alpha)\phi, \\ \phi_1 &\rightarrow \phi_1 - \alpha\phi_2, \\ \phi_2 &\rightarrow \phi_2 + \alpha\phi_1, \end{aligned} \quad (1.35)$$

and we find

$$\begin{aligned} \phi'_1 &\rightarrow \phi'_1 - \alpha\phi_2, \\ \phi_2 &\rightarrow \phi_2 + \alpha v + \alpha\phi'_1, \end{aligned} \quad (1.36)$$

Thus ϕ_2 undergoes an inhomogenous gauge transformation like the gauge field A_μ . So we could use the freedom of gauge to set $\phi_2 = 0$, in which the mixing term (1.34) and the Goldstone boson vanish.

As we see, the introduction of the gauge field A_μ and the requirement of local gauge invariance fully change the physical condition: the Lagrangian (1.28) describes a massive scalar field accompanied with a massless Goldstone boson, while the Lagrangian (1.29) describes a massive gauge boson A_μ and a massive scalar boson

ϕ . However, the total number of the particle states remains unchanged. Before spontaneous symmetry breaking, the theory had four particle states: two spin-zero particles ϕ and ϕ^* plus two polarization states of the massless gauge boson A_μ , i.e. four states in total. After spontaneous symmetry breaking, we have one scalar particle plus three polarization states of the massive gauge boson A_μ , i.e. still four states. Therefore, we could say that the massless gauge boson "ate" the massless Goldstone boson to become a massive gauge boson.

The Higgs mechanism introduced above is important for the following discussion. In general theories involving massive gauge bosons are non-renormalizable, because of the $k_\mu k_\nu/m^2$ -term in the gauge boson propagator. However, in the original Lagrangian (1.29), the gauge field is formally massless, where no issues with renormalizability occurs. It turns out that spontaneous symmetry breaking would not affect the renormalizability of the theory [13–15]. Note that for the Abelian gauge theory described by the Lagrangian (1.28), the spontaneous generation of the gauge boson mass is not necessary to achieve renormalizability. We would not undermine the renormalizability if we introduce a mass term for the gauge boson, provided that the gauge field A_μ is coupled with a conserved current. However, this is not valid for a non-Abelian theory. The spontaneous generation of gauge boson masses is the only way to ensure renormalizability.

In sum, Higgs mechanism is a remarkable result, suggesting the possibility of establishing spontaneously broken gauge theories in which the interactions are mediated by massive gauge bosons.

1.3.2 The Higgs Mechanism in the Non-Abelian Theory

In order to explore additional complications from spontaneous symmetry breaking of a non-Abelian theory, we choose $SU(2)$ gauge theory as a prototype.

At first, we choose a doublet representation of complex scalar fields, coupled to the gauge fields in a gauge invariant way. Such a Lagrangian is defined by

$$L = -\frac{1}{4}G_{\mu\nu}^i G_i^{\mu\nu} + \left(\partial^\mu\phi + i\frac{g}{2}\sigma^i B^{\mu i}\phi\right)^\dagger \left(\partial_\mu\phi + i\frac{g}{2}\sigma_i B_\mu^i\phi\right) - \mu^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2, \quad (1.37)$$

where the scalar field ϕ represents the $SU(2)$ doublet

$$\phi = \begin{pmatrix} \phi_a \\ \phi_b \end{pmatrix}, \quad (1.38)$$

and the σ_i 's denote the Pauli matrices.

For $\mu^2 > 0$ the Lagrangian (1.37) describes a system of massless gauge fields in interaction with massive scalars of mass μ . Now suppose we choose $\mu^2 < 0$, then the potential $V(\phi) = \mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2$ exhibits its minimum at finite values of ϕ . The manifold of points in the space of fields ϕ_a, ϕ_b for which the minimum of the potential $V(\phi)$ occurs is invariant under $SU(2)$ -transformations. Therefore, we could choose a specific $SU(2)$ frame for which we have $\langle 0|\phi|0\rangle = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v \end{pmatrix}$, where $v = \sqrt{-\mu^2/\lambda}$. From eq. (2.35) we obtain the mass term of the gauge field, which is

$$\begin{aligned} \frac{g^2}{4}[(\sigma^i B_\mu^i)]^\dagger[(\sigma^i B_\mu^i)] &= \frac{g^2}{4}(\phi^\dagger\tau^j\tau^i\phi)(B_\mu^i B^{\mu i}) \\ &= \frac{g^2}{8} \cdot v^2[(B_\mu^1)^2 + (B_\mu^2)^2 + (B_\mu^3)^2]. \end{aligned} \quad (1.39)$$

Thus we could conclude: we obtain massive gauge fields after spontaneous symmetry breaking. The gauge boson mass matrix is $SU(2)$ symmetric, namely, the three gauge bosons are degenerate in mass. This is a special feature of the spontaneous symmetry breaking involving an $SU(2)$ doublet.

Note that the particle content of the theory includes three massive gauge fields and one massive scalar field. Three of the scalar fields (four real fields originally) have been eaten to provide the longitudinal components of the massive gauge fields.

Next, we choose an $SU(2)$ triplet representation of real scalar fields

$$\phi = \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}. \quad (1.40)$$

We require invariance under the gauge transformation

$$\phi \rightarrow e^{iT_i \alpha_i} \phi, \quad (1.41)$$

where the exponential factor is a 3×3 matrix. The operator T_i generates isospin rotations about the i -axis and satisfies the usual $SU(2)$ algebra

$$[T^j, T^k] = i\epsilon_{jkl} T^l. \quad (1.42)$$

The explicit matrix representation is

$$(T^j)_{kl} = -i\epsilon_{jkl}. \quad (1.43)$$

Following the procedure outline in Section 1.2, we have the covariant derivative as follows

$$D_\mu = \partial_\mu - igT_i B_{\mu i}, \quad (1.44)$$

or, in the adjoint representation, the covariant derivative takes the form,

$$(D_\mu)_{kl} = \delta_{kl} \partial_\mu + g\epsilon_{jkl} B_{\mu j}. \quad (1.45)$$

Then the Lagrangian of the theory is

$$L = -\frac{1}{4} G_{\mu\nu}^i G_i^{\mu\nu} + \frac{1}{2} (\partial^\mu \phi_i - g\epsilon_{ijk} B_j^\mu \phi_k) (\partial_\mu \phi_i - g\epsilon_{ilm} B_l^\mu \phi_m) - \frac{1}{2} \mu^2 \phi_i \phi_i - \frac{1}{4} \lambda (\phi_i \phi_i)^2. \quad (1.46)$$

When ϕ is the unique minima of the potential $V(\phi) = \frac{1}{2}\mu^2\phi_i\phi_i + \frac{1}{4}\lambda(\phi_i\phi_i)^2$, the spectrum is that of an ordinary, isospin-conserving gauge field theory: three massive scalar fields, each with mass μ , and three massless gauge fields B_μ . Since each massless gauge boson has two polarization states, then the number of particle states is $3 \times 1 + 3 \times 2 = 9$.

If we choose $\mu^2 < 0$, spontaneous symmetry breaking occurs. We could choose a particular coordinate system of fields such that we have

$$\langle 0 | \phi | 0 \rangle = \begin{pmatrix} 0 \\ 0 \\ v \end{pmatrix}. \quad (1.47)$$

This vector remains invariant under rotation generated by T_3 , i.e. the subgroup $U(1) \subset SU(2)$, generated by the third generator T_3 remains unbroken.

We shift the scalar fields and expand around the v , using

$$\phi \rightarrow \exp\left[\frac{i}{v}(\zeta_1 T_1 + \zeta_2 T_2)\right] \begin{pmatrix} 0 \\ 0 \\ v + \eta \end{pmatrix}. \quad (1.48)$$

We could exploit the gauge invariance of the theory by letting

$$\phi \rightarrow \exp\left[-\frac{i}{v}(\zeta_1 T_1 + \zeta_2 T_2)\right] \phi = \begin{pmatrix} 0 \\ 0 \\ v + \eta \end{pmatrix}. \quad (1.49)$$

With the help of the new gauge, we obtain the Lagrangian

$$L = \frac{1}{2}(\partial_\mu \eta \partial^\mu \eta + 2\mu^2 \eta^2) - \frac{1}{4}G_{\mu\nu}^i G^{\mu\nu i} + \frac{1}{2}g^2 v^2 [B_\mu^1 B^{1\mu} + B_\mu^2 B^{2\mu}] + \dots \quad (1.50)$$

From this Lagrangian, we conclude: η has become a massive Higgs scalar field, with mass $\sqrt{-2\mu^2}$; the Goldstone bosons η_1 and η_2 have disappeared completely, i.e.,

they have been eaten up by gauge fields; the gauge bosons B_1^μ and B_2^μ respectively coupled to the broken generators T_1 and T_2 have acquired a common mass gv ; the gauge boson B_3^μ remains massless, reflecting the invariance of the vacuum under the generator T_3 .

1.4 Standard Model of the Electroweak Interactions

1.4.1 The General Ideas for Building Spontaneous Broken Gauge Theories

After investigating the examples above, we are ready to discuss the general features of spontaneous broken gauge theories. Let us consider a Lagrangian which is invariant under local gauge transformations of a group G . The generators T_i follow the commutation relations

$$[T_i, T_j] = if_{ijk}T_k, \quad (1.51)$$

where $i, j, k = 1, \dots, N$ and f_{ijk} is the structure constant of G . An arbitrary infinitesimal transformation of the group G could be parametrized by $1 - i\epsilon_i T_i$, where ϵ_i 's are infinitesimal parameters.

The scalar field ϕ is assumed to transform under a n -dimensional representation of G . We assume that the field ϕ are real, since a complex field ϕ can always be decomposed into two real ones. For an infinitesimal transformation of G we have:

$$\delta\phi = i\epsilon_i S_i \phi. \quad (1.52)$$

The Lagrangian is defined by

$$L = -\frac{1}{4}G_{\mu\nu}^i G_i^{\mu\nu} + \frac{1}{2}[(\partial^\mu + igS_i A_i^\mu)\phi]^\dagger [(\partial_\mu + igS_j A_{j\mu})\phi] - V(\phi), \quad (1.53)$$

where $V(\phi)$ is a quartic potential in ϕ , invariant under G .

We assume that spontaneous symmetry breaking occurs and the potential exhibits its minimum at $\phi = v$, where v is a n -dimensional vector. The gauge boson mass matrix is then given by

$$(M^2)_{ij} = -g^2(S_i v) \cdot (S_j v). \quad (1.54)$$

In general, there will exist a M -dimensional subgroup G' of G , which preserves an invariance of the vacuum.

Let $T_i(G')$ be the generators of G' , then $T_i(G')v = S_i(G')v = 0$. There exist $(N - M)$ generators of G , for which $T_i v \neq 0$, i.e. one has $(N - M)$ Goldstone bosons. Therefore the $N \times N$ dimensional mass matrix denoted in eq.(1.54) is actually an $(N - M) \times (N - M)$ dimensional matrix, if we leave out all terms for which $S_i v = 0$ because of the S -invariance of the vacuum. The mass matrix (1.54) needs to be diagonalized if we would like to find the massive vector bosons of definite mass. There exist $(N - M)$ massive gauge bosons. The $(N - M)$ Goldstone bosons are absorbed into the longitudinal components of the $(N - M)$ massive gauge bosons.

Up to now, our discussion mainly focuses on scalar and vector fields. Next, we need to incorporate fermion fields into the gauge theory, by adding to the Lagrangian (1.53) the terms

$$L^{\text{fermion}} = \bar{\psi}_L(i\cancel{\partial} - gf_{Li}A_i)\psi_L + \bar{\psi}_R(i\cancel{\partial} - gf_{Ri}A_i)\psi_R - (m\bar{\psi}_R\psi_L + \text{h.c.}) \quad (1.55)$$

and

$$L^{\text{int}} = -G\bar{\psi}_R(R\phi)\psi_L + \text{h.c.} \quad (1.56)$$

where ψ_L , ψ_R stand for the left-handed and right-handed fermion fields. The fields ψ_L and ψ_R transform under G as certain irreducible representations. The matrices f_L , f_R denote the transformation properties of the left-handed and right-handed fermion

fields. In eq. (2.55), we include the Yukawa interaction term of the fermion fields with the scalar fields. The matrices R are constructed so that $\bar{\psi}_R(R\phi)\bar{\psi}_L$ is invariant under the gauge group. We also include a bare mass term ($m\bar{\psi}_R\psi_L + h.c.$), which must be G -invariant.

From the eqs. (1.55) and (1.56) we obtain the fermion mass matrix after the spontaneous symmetry breaking,

$$L^{\text{fermion mass}} = -G\bar{\psi}_R(Rv)\psi_L - m\bar{\psi}_R\psi_L + h.c. \quad (1.57)$$

After introducing the general properties of spontaneous broken gauge theories, we are ready to give the general recipe for building renormalizable gauge theories. The Lagrangian is constructed as follows:

1. Select the gauge group, the representations of left-handed and right-handed fermions and the scalar fields.
2. Couple the gauge fields invariantly to the fermion and scalar fields.
3. Couple the gauge invariant quartic polynomial of the scalar fields so that the potential reaches its minimum for nonvanishing vacuum expectation values v .
4. Construct the gauge invariant Yukawa couplings between the fermions and scalars.

The gauge boson mass matrix has the structure:

$$\frac{1}{2}g^2v^2W_\mu^2,$$

the fermion mass matrix is

$$G \cdot v\bar{\psi}\psi.$$

1.4.2 The Glashow-Weinberg-Salam Theory

We are now ready to write down the spontaneously broken gauge theory that gives experimentally confirmed description of weak and electromagnetic interactions, a model introduced by Glashow, Weinberg, and Salam (GWS) [1–3]. We begin with the doublet of the weak isospin consisting of left-handed electron and its neutrino,

$$\psi_L^e \equiv \begin{pmatrix} \nu_e \\ e \end{pmatrix}_L, \quad (1.58)$$

where the left-handed states are

$$\begin{aligned} \nu_L &= \frac{1}{2}(1 - \gamma_5)\nu_e \\ e_L &= \frac{1}{2}(1 - \gamma_5)e \end{aligned} \quad (1.59)$$

The electron neutrino is known to be nearly massless. It is convenient to idealize it as exactly massless, in which case the right-handed state

$$\nu_R = \frac{1}{2}(1 + \gamma_5)\nu_e \quad (1.60)$$

does not exist. Thus we have only one right-handed fermion,

$$\psi_R^e \equiv e_R = \frac{1}{2}(1 + \gamma_5)e, \quad (1.61)$$

which is an $SU(2)$ -singlet.

Note that we need to have the $U(1)$ -factor in the gauge group $SU(2) \otimes U(1)$ to represent the electric charge. This cannot be an $SU(2)$ generator since the photon couples both to the left-handed and right-handed electron. To incorporate the electromagnetic interaction, we denote the $U(1)$ generator as “weak hypercharge”, Y . Requiring that the Gell-Mann-Nishijima relation for electric charge,

$$Q = I_3 + \frac{1}{2}Y \quad (1.62)$$

be satisfied leads to the assignments

$$Y_L = -1, \quad Y_R = -2. \quad (1.63)$$

By construction, the weak-isopin projection I_3 and the weak hypercharge Y commute:

$$[I_3, Y] = 0. \quad (1.64)$$

Let us take the group of transformations generated by I and Y to be the gauge group $SU(2)_L \otimes U(1)_Y$ for the gauge theory. In order to construct the theory, we introduce the gauge bosons $A_\mu^1, A_\mu^2, A_\mu^3$ for $SU(2)_L$ and B_μ for $U(1)_Y$.

The Lagrangian for the theory might be written as

$$L = L^{\text{gauge}} + L^{\text{fermion}} + L^{\text{scalar}}, \quad (1.65)$$

and the gauge boson part of the Lagrangian is

$$L^{\text{gauge}} = -\frac{1}{4}F_{\mu\nu}^l F_l^{\mu\nu} - \frac{1}{4}G_{\mu\nu} G^{\mu\nu} \quad (1.66)$$

where

$$F_{\mu\nu}^l = \partial_\nu A_\mu^l - \partial_\mu A_\nu^l + g\epsilon_{jkl}A_\mu^j A_\nu^k \quad (1.67)$$

for the $SU(2)_L$ gauge fields and

$$G_{\mu\nu} = \partial_\nu B_\mu - \partial_\mu B_\nu \quad (1.68)$$

for the $U(1)_Y$ gauge field.

We introduce a complex doublet of scalar fields

$$\phi \equiv \begin{pmatrix} \phi^\dagger \\ \phi_0 \end{pmatrix} \quad (1.69)$$

which transforms as an $SU(2)_L$ doublet and has the hypercharge

$$Y_\phi = 1$$

by virtue of the Gell-Mann-Nishijima relation.

Now we are ready to write down the fermion and scalar parts of the Lagrangian:

$$L^{\text{fermion}} = \bar{\psi}_R^e \left(\not{\partial} + \frac{ig'}{2} \not{B} Y \right) \psi_R^e + \bar{\psi}_L^e i \left(\not{\partial} + \frac{ig'}{2} \not{B} Y + \frac{ig}{2} \sigma_i A_i \right) \psi_L^e \quad (1.70)$$

and

$$L^{\text{scalar}} = (D^\mu \phi)^\dagger (D_\mu \phi) - V(\phi^\dagger \phi), \quad (1.71)$$

where the covariant derivative is

$$D_\mu = \partial_\mu + \frac{ig'}{2} B_\mu Y + \frac{ig}{2} \sigma_i A_{i\mu} \quad (1.72)$$

and the most general form of the potential is

$$V(\phi^\dagger \phi) = \mu^2 (\phi^\dagger \phi) + |\lambda| (\phi^\dagger \phi)^2. \quad (1.73)$$

We also add an interaction term, which involves Yukawa couplings of the scalar to the fermions,

$$L^{\text{Yukawa}} = -G_e [\bar{\psi}_R^e (\phi^\dagger \psi_L^e) + (\bar{\psi}_L^e) \psi_R^e], \quad (1.74)$$

which is symmetric under $SU(2)_L \otimes U(1)_Y$ transformations,

Now let us take $\mu^2 < 0$, then the $SU(2)_L \otimes U(1)_Y$ symmetry is spontaneously broken. We choose a $SU(2)$ frame so that the vacuum expectation values of ϕ take the form:

$$\langle 0 | \phi | 0 \rangle = \begin{pmatrix} 0 \\ v/\sqrt{2} \end{pmatrix}, \quad (1.75)$$

where $v = \sqrt{-\mu^2/|\lambda|}$, which breaks both $SU(2)_L$ and $U(1)_Y$ symmetries. However, the electric charge, which is a linear combination of T_3 and Y remains unbroken. The photon will therefore remain massless, while three other gauge bosons will acquire mass.

We next expand the Lagrangian about the minimum of the Higgs potential V by letting

$$\phi = \exp\left(\frac{i\zeta_i\sigma_i}{2v}\right)\begin{pmatrix} 0 \\ (v+\eta)/\sqrt{2} \end{pmatrix} \quad (1.76)$$

and transforming to U-gauge:

$$\phi \rightarrow \exp\left(-\frac{i\zeta_i\sigma_i}{2v}\right)\phi = \begin{pmatrix} 0 \\ (v+\eta)/\sqrt{2} \end{pmatrix}, \quad (1.77)$$

$$\sigma_i A_{i\mu} \rightarrow \sigma_i A'_{i\mu}, \quad (1.78)$$

$$B_\mu \rightarrow B_\mu, \quad (1.79)$$

$$\psi_R^e \rightarrow \psi_R^e, \quad (1.80)$$

$$\psi_L \rightarrow \exp\left(-\frac{i\zeta_i\sigma_i}{2v}\right) \quad (1.81)$$

We now express the Lagrangian in terms of the U-gauge fields (1.77-1.81) and explore the results of spontaneous symmetry breaking. The Yukawa part of Lagrangian has become

$$L^{\text{Yukawa}} = -G_e \frac{v+\eta}{\sqrt{2}} (\bar{e}_R e_L + \bar{e}_L e_R) = -\frac{G_e v}{\sqrt{2}} \bar{e}e - \frac{G_e \eta}{\sqrt{2}} \bar{e}e \quad (1.82)$$

so the electron has acquired a mass

$$m_e = G_e v / \sqrt{2} \quad (1.83)$$

The scalar part of the Lagrangian now becomes

$$L^{\text{scalar}} = \frac{1}{2}(\partial^\mu\eta)(\partial_\mu\eta) - \mu^2\eta^2 + \frac{v^2}{8}[g^2|A_\mu^1 - iA_\mu^2|^2 + (g'B_\mu - gA_\mu^3)^2] + \dots \quad (1.84)$$

We see immediately that the η field has acquired a mass $m_H = \sqrt{-2\mu^2}$; it is the physical Higgs boson. If we define the charged gauge fields

$$W_\mu^\pm \equiv \frac{A_\mu^1 \mp iA_\mu^2}{\sqrt{2}}, \quad (1.85)$$

the term proportional to g^2v^2 is identified as a mass term for the charge gauge bosons:

$$\frac{g^2v^2}{8}(|W_\mu^+|^2 + |W_\mu^-|^2), \quad (1.86)$$

corresponding to charged boson masses

$$M_{W_\pm} = gv/2. \quad (1.87)$$

Then, defining the orthogonal combinations

$$Z_\mu = \frac{-g'B_\mu + gA_\mu^3}{\sqrt{g^2 + g'^2}} \quad (1.88)$$

and

$$A_\mu = \frac{gB_\mu + g'A_\mu^3}{\sqrt{g^2 + g'^2}} \quad (1.89)$$

we see that the neutral boson has acquired a mass

$$M_{Z^0} = \sqrt{g^2g'^2}v/2 = M_W\sqrt{1 + g'^2/g^2} \quad (1.90)$$

and that the gauge field A_μ remains massless.

Next, let us investigate the interactions. We could read off the interactions among the gauge bosons and fermions from L^{fermion} . For the charged gauge bosons we have

$$\begin{aligned} L^{W-f} &= -\frac{g}{\sqrt{2}}(\bar{\nu}_L\gamma_\mu e_L W_\mu^+ \bar{e}_L\gamma^\mu \nu_L W_\mu^-) \\ &= -\frac{g}{2\sqrt{2}}[\bar{n}u\gamma_\mu(1 - \gamma_5)eW_\mu^+ \bar{e}\gamma_\mu(1 - \gamma_5)\nu W_\mu^-], \end{aligned} \quad (1.91)$$

where we identify the coupling constant as

$$\frac{g^2}{8} = \frac{G_f M_W^2}{\sqrt{2}}. \quad (1.92)$$

with G_f is the Fermi constant. Similarly, the neutral gauge boson couplings to fermions are given by

$$L^{0-f} = \frac{gg'}{\sqrt{g^2 + g'^2}} \bar{e} \gamma_\mu e A_\mu - \frac{\sqrt{g^2 + g'^2}}{2} \bar{\nu}_L \gamma_\mu \nu_L Z_\mu + \frac{Z_\mu}{\sqrt{g^2 + g'^2}} + \frac{Z_\mu}{\sqrt{g^2 + g'^2}} \left[-g'^2 \bar{e}_R \gamma_\mu e_R + \frac{(g^2 - g'^2)}{2} \bar{e}_L \gamma_\mu e_L \right]. \quad (1.93)$$

Thus we can identify A_μ as the photon, setting

$$\frac{gg'}{\sqrt{g^2 + g'^2}} = e. \quad (1.94)$$

It is convenient to introduce a weak mixing angle θ_W to parametrize the mixing of the neutral gauge bosons. Defining

$$g' = g \tan \theta_W, \quad \sin \theta_W = \frac{g'}{g^2 + g'^2}, \quad (1.95)$$

we could rewrite eqs. (1.88) and (1.89) as

$$\begin{aligned} Z_\mu &= -B_\mu \sin \theta_W + A_\mu^3 \cos \theta_W, \\ A_\mu &= -A_\mu \cos \theta_W + A_\mu^3 \sin \theta_W. \end{aligned} \quad (1.96)$$

By the virtue of the relation (1.95), the coupling constants of the $SU(2)_L$ and $U(1)_Y$ gauge groups might be written as

$$g = \frac{e}{\sin \theta_W} \geq e, \quad g' = \frac{e}{\cos \theta_W} \geq e \quad (1.97)$$

And the masses of gauge bosons could be rewritten as

$$M_W = \left(\frac{\pi \alpha}{\sqrt{2} G_F} \right)^{\frac{1}{2}} \frac{1}{\sin \theta_W}, \quad (1.98)$$

$$M_Z = \frac{M_W}{\cos \theta_W}. \quad (1.99)$$


It is therefore convenient to express the interaction terms of the Lagrangian (1.93) in terms of the weak mixing angle as

$$\begin{aligned}
L^{0-f} = & e\bar{e}\gamma^\mu e A_\mu - \frac{1}{\sqrt{2}} \left(\frac{G_F M_Z^2}{\sqrt{2}} \right)^{\frac{1}{2}} \bar{\nu}\gamma^\mu(1 - \gamma_5)\nu Z_\mu \\
& - \frac{1}{\sqrt{2}} \left(\frac{G_F M_Z^2}{\sqrt{2}} \right)^{\frac{1}{2}} [2\sin^2\theta_W \bar{e}\gamma^\mu(1 + \gamma_5)e Z_\mu \\
& + (2\sin^2\theta_W - 1)\bar{e}\gamma^\mu(1 - \gamma_5)e Z_\mu]
\end{aligned} \tag{1.100}$$

and the interaction terms of the Lagrangian (1.91) can be rewritten as

$$L^{W-f} = - \left(\frac{G_F M_W^2}{\sqrt{2}} \right)^{\frac{1}{2}} [\bar{\nu}_e \gamma_\mu (1 - \gamma_5) e W_\mu^+ \bar{e} \gamma_\mu (1 - \gamma_5) \nu_e W_\mu^-], \tag{1.101}$$

from which we could derive Feynman rules for the elementary vertices. These are shown below.



A Feynman diagram showing two incoming neutrinos, labeled ν_e , meeting at a vertex. From this vertex, a wavy line representing a Z^0 boson extends to the right. The diagram is associated with the following mathematical expression:

$$Z_\mu^0 - \frac{i}{\sqrt{2}} \left(\frac{G_F M_Z^2}{\sqrt{2}} \right)^{\frac{1}{2}} \bar{\nu}_e \gamma_\mu (1 - \gamma_5) \nu_e$$

A Feynman diagram showing two incoming electrons, labeled e , meeting at a vertex. From this vertex, a wavy line representing a Z^0 boson extends to the right. The diagram is associated with the following mathematical expression:

$$Z_\mu^0 - \frac{i}{\sqrt{2}} \left(\frac{G_F M_Z^2}{\sqrt{2}} \right)^{\frac{1}{2}} \bar{e} \gamma_\mu [R_e (1 + \gamma_5) + L_e (1 - \gamma_5)] e,$$

$$R_e \equiv 2 \sin^2 \theta_W,$$

$$L_e \equiv 2 \sin^2 \theta_W - 1.$$

CHAPTER TWO

Techniques for the Calculation of Electroweak Radiative Corrections at the One-Loop Level

In the previous chapter, We have introduced the minimal theory of electroweak interactions, i.e. $SU(2)_L \otimes U(1)_Y$ theory of electron, proposed by S. L. Glashow [1], S. Weinberg [2], and A. Salam [3], which exhibited the basic motivations and principal features. This theory has been extended to the hadronic degrees of freedom by S. L. Glashow, J. Iliopoulos and L. Maiani [4]. And the Weinberg-Salam-Glashow-Iliopoulos-Maini model is the most comprehensive formulation of a theory of the unified electroweak interaction at present. It is theoretically consistent and confirmed by all experimentally known phenomena of the electroweak origin. After the Weinberg-Salam-Glashow-Iliopoulos-Maini model was proposed, 't Hooft and M. Veltman proved its renormalizability [13–16]. Therefore, the standard model of the electroweak interaction is a calculable quantum field theory capable for precision calculations in high energy physics. Theoretical predictions should have a precision comparable to or even better than the experimental uncertainties. If the experimental precision of the order of 1% the classical level of the theory is no longer sufficient. We have to take into account quantum corrections: the radiative corrections.

In this chapter, we will review the corresponding formulae and techniques for the evaluation of the one loop radiative corrections for the electroweak theory [30–37]. At first with the help of Faddeev-Popov gauge fixing technique, the complete renormalizable Lagrangian for the electroweak SM is given, Next, its renormalization will

be discussed. Then we will introduce the classification and techniques for calculating one loop integrals. At last, we will present some explicit calculations of one-loop radiative correction as illustrations of the method described in this chapter.

2.1 The Model

The classical Lagrangian of the electroweak SM consists of a gauge boson (Yang-Mills), a scalar(Higgs) and a fermion part

$$L^{classical} = L^{\text{gauge}} + L^{\text{scalar}} + L^{\text{fermion}} + L^{\text{Yukawa}}, \quad (2.1)$$

where each of them is separately gauge invariant.

The gauge boson fields includes an isotriplet W_μ^a and an isosinglet B_μ . The isotriplet W_μ^a , $a = 1, 2, 3$ is associated with the generator σ_a (Pauli matrices) of the group $SU(2)_L$, and the isosinglet B_μ is associated with the weak hypercharge Y of the group $U(1)_Y$. The gauge field the Lagrangian is as usual,

$$L^{\text{gauge}} = -\frac{1}{4}F_{\mu\nu}^l F_l^{\mu\nu} - \frac{1}{4}G_{\mu\nu}G^{\mu\nu} \quad (2.2)$$

where

$$F_{\mu\nu}^l = \partial_\nu W_\mu^l - \partial_\mu W_\nu^l + g_2 \epsilon_{jkl} W_\mu^j W_\nu^k \quad (2.3)$$

for the $SU(2)_L$ gauge fields and

$$G_{\mu\nu} = \partial_\nu B_\mu - \partial_\mu B_\nu \quad (2.4)$$

for the $U(1)_Y$ gauge field. The covariant derivative here is given by

$$D_\mu = \partial_\mu - ig_2 \sigma_a W_\mu^a + ig_1 \frac{Y}{2} B_\mu. \quad (2.5)$$

where g_1 is the $U(1)_Y$ gauge coupling and g_2 is the $SU(2)_L$ gauge coupling. The electric charge operator Q is composed of the weak isospin projection I_3 and the

weak hypercharge Y according to the Gell-Mann Nishijima relation.

$$Q = I_3 + \frac{1}{2}Y. \quad (2.6)$$

The scalar Lagrangian is as usual:

$$L^{\text{Scalar}} = (D^\mu \Phi)^\dagger (D_\mu \Phi) - V(\Phi), \quad (2.7)$$

where

$$\Phi(x) = \begin{pmatrix} \phi^\dagger(x) \\ \phi_0(x) \end{pmatrix} \text{ with } Y_\Phi = 1. \quad (2.8)$$

Here, we express the Higgs potential in another way

$$V(\Phi) = \frac{\lambda}{4}(\Phi^\dagger \Phi)^2 - \mu^2 \Phi^\dagger \Phi \quad (2.9)$$

where $\lambda > 0$, $\mu > 0$ such that it gives rise to spontaneous symmetry breaking.

The fermion part is extended to the lepton families (ψ^l) and quark families (ψ^q). The left-handed fermion of each lepton and quark generation are grouped into $SU(2)_L$ doublets:

$$\begin{aligned} \psi_L^l &= \frac{1}{2}(1 - \gamma_5)\psi_l = \begin{pmatrix} \nu_l \\ l \end{pmatrix}_L \\ \psi_L^q &= \frac{1}{2}(1 - \gamma_5)\psi_q = \begin{pmatrix} u_i \\ d_i \end{pmatrix}_L \end{aligned} \quad (2.10)$$

where $l \equiv e, \mu, \tau$, $u_i \equiv u, c, t$ and $d_i \equiv d, s, b$. And the right-handed fermion are grouped into singlets:

$$\begin{aligned} \psi_R^l &= \frac{1}{2}(1 + \gamma_5)\psi^l; \\ (u_R)_i &= \frac{1}{2}(1 + \gamma_5)u_i, (d_R)_i = \frac{1}{2}(1 + \gamma_5)d_i \end{aligned} \quad (2.11)$$

Then the fermion Lagrangian reads off

$$L^{\text{fermion}} = \sum_i (\bar{\psi}_L^l i \not{D} \psi_L^l + \bar{\psi}_L^q i \not{D} \psi_L^q) + \sum_i (\bar{\psi}_L^l i \not{D} \psi_L^l + \bar{u}_{iR} i \not{D} u_{iR} + \bar{d}_{iR} i \not{D} d_{iR}). \quad (2.12)$$

And the Yukawa Lagrangian reads

$$L^{\text{Yukawa}} = - \sum_{ij} [(\bar{\psi}_L^l)_i G_{ij}^l (\psi_R^l)_j \Phi + (\bar{\psi}_L^q)_i G_{ij}^u (u_R)_j \tilde{\Phi} + (\bar{\psi}_L^q)_i G_{ij}^d (d_R)_j \Phi + \text{h.c.}] \quad (2.13)$$

where G_{ij}^l , G_{ij}^u and G_{ij}^d are the Yukawa coupling matrices, $\tilde{\Phi} = \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix}$ is the charge conjugated Higgs field and $\phi^- = (\phi^\dagger)^*$.

From the construction of Higgs part of Lagrangian, we have the vacuum expectation value

$$|\langle 0 | \Phi | 0 \rangle|^2 = \frac{2\mu^2}{\lambda} = \frac{v^2}{2} \neq 0 \quad (2.14)$$

We expand the scalar field around the ground state so that the Higgs field can be expressed as

$$\Phi(x) = \begin{pmatrix} \phi^\dagger(x) \\ \frac{1}{\sqrt{2}}(v(x) + H(x) + i\chi(x)) \end{pmatrix}, \quad (2.15)$$

where the components ϕ^\dagger , H and χ have zero vacuum expectation values. ϕ^\dagger , ϕ^- and χ are unphysical states which can be eliminated by the unitary gauge. The field H is the physical Higgs field with the mass

$$M_H = \sqrt{2}\mu. \quad (2.16)$$

The physical gauge fields W_μ^\pm , Z^0 and A_μ are related to W_μ^a and B_μ by

$$W_\mu^\pm = \frac{1}{\sqrt{2}}(W_\mu^1 \mp iW_\mu^2),$$

$$\begin{pmatrix} Z_\mu \\ A_\mu \end{pmatrix} = \begin{pmatrix} c_w & s_w \\ -s_w & c_w \end{pmatrix} \begin{pmatrix} W_\mu^3 \\ B_\mu \end{pmatrix}. \quad (2.17)$$

where

$$\begin{aligned} c_w &\equiv \cos \theta_W = \frac{g_2}{\sqrt{g_1^2 + g_2^2}}, \\ s_w &\equiv \sin \theta_W = \frac{g_1}{\sqrt{g_1^2 + g_2^2}}. \end{aligned} \quad (2.18)$$

The physical fermion fields are obtained by diagonalizing the corresponding mass matrices

$$\begin{aligned} (f_L)_i &= (U_{f,L})_{ik} (f'_L)_k \\ (f_R)_i &= (U_{f,R})_{ik} (f'_R)_k \end{aligned} \quad (2.19)$$

where $f \equiv \nu_l, l, u_i$ and d_i .

The resulting masses are

$$\begin{aligned} M_Z &= \frac{1}{2} \sqrt{g_1^2 + g_2^2} v, \\ M_W &= M_Z c_W = \frac{1}{2} g_2 v, \\ M_\gamma &= 0, \\ m_{f,i} &= \frac{v}{\sqrt{2}} (U_{f,L})_{ik} G_{km}^f (U_{f,R})_{mi}. \end{aligned} \quad (2.20)$$

By identifying the coupling of the photon field A_μ to the electron with the electrical charge $e = \sqrt{4\pi\alpha}$, we have

$$e = \frac{g_1 g_2}{\sqrt{g_1^2 + g_2^2}}. \quad (2.21)$$

The diagonalization of the fermion mass matrices introduces a unitary quark mixing matrix into the quark-W-boson couplings

$$V_{ij} = (U_{u,L})_{ik} (U_{d,L})_{kj}^\dagger. \quad (2.22)$$

Thus, the relations (2.16), (2.20), (2.21) and (2.22) allow us to replace the set of parameters $\{g_1, g_2, \lambda, \mu^2, G^l, G^u, G^d\}$ with the parameter $\{e, M_W, M_Z, M_h, m_{f,i}, V_{ij}\}$

which are physical. Furthermore, we could express the Lagrangian (2.1) in terms of physical parameters and fields.

Next, we need to apply Faddeev-Popov gauge fixing technique [38,39] to quantize $L^{\text{classical}}$, which requires the specification of a gauge. We choose a renormalizable 't Hooft gauge with the following linear gauge fixings

$$\begin{aligned} F^\pm &= (\xi_1^W)^{-\frac{1}{2}} \partial^\mu W_\mu^\dagger \mp i(\xi_2^W)^{-\frac{1}{2}} \phi^\pm \\ F^Z &= (\xi_1^Z)^{-\frac{1}{2}} \partial^\mu Z_\mu - M_Z (\xi_2^Z)^{\frac{1}{2}} \chi \\ F^\gamma &= (\xi_1^\gamma)^{-\frac{1}{2}} \partial^\mu A_\mu, \end{aligned} \tag{2.23}$$

which lead to the gauge fixing Lagrangian

$$L^{\text{gauge-fixing}} = -\frac{1}{2} [(F^\gamma)^2 + (F^Z)^2 + 2F^+ F^-]. \tag{2.24}$$

L^{fix} includes the unphysical components of the gauge fields. To cancel the unphysical effects, We need to introduce Faddeev Popov ghosts (scalar anti-commuting fields) $\bar{u}^\alpha(x)$, $u^\alpha(x)$ ($\alpha = \pm, \gamma, Z$) with the Lagrangian

$$L^{\text{FP}} = \bar{u}^\alpha(x) \frac{\delta F^\alpha}{\delta \theta^\beta(x)} u^\beta(x), \tag{2.25}$$

where $\frac{\delta F^\alpha}{\delta \theta^\beta(x)}$ is the variation of the gauge fixing operators F^α under infinitesimal gauge transformations characterized by $\theta^\beta(x)$.

The 't Hooft Feynman gauge $\xi^\alpha = 1$ will simplify the problem. At lowest order the poles of the ghost fields, unphysical Higgs fields and longitudinal gauge fields coincide with the poles of the corresponding transverse gauge fields. Moreover, there is no mixing between gauge Higgs and gauge fields.

With the help of $L^{\text{gauge-fixing}}$ and L^{FP} , we obtain the complete renormalizable Lagrangian for the electroweak SM:

$$L^{\text{SM}} = L^{\text{classical}} + L^{\text{gauge-fixing}} + L^{\text{FP}}. \quad (2.26)$$

The corresponding Feynman rules are given in Appendix.A.

2.2 Renormalization in the Electroweak SM

The Lagrangian (2.1) of the minimal $SU(2)_L \otimes U(1)_Y$ model includes a certain number of free parameters $\{e, M_W, M_Z, M_h, m_{f,i}, V_{ij}\}$, which have to be determined experimentally. These parameters could be directly related to experimental quantities (at the tree-level), but this direct relation is no more valid when it comes to higher order corrections. We usually called the parameters of the original Lagrangian bare parameters, which differ from corresponding physical quantities by ultra-violet (UV)-divergent contributions. These divergences would cancel in relations between physical quantities in renormalizable theories. The renormalizability of non-Abelian gauge theories with spontaneous symmetry breaking as proven by 't Hooft [13, 14], which allows meaningful predictions in the electroweak SM.

We are using the counterterm approach to realize the renormalization. Here the UV-divergent bare parameters are expressed by finite renormalized parameters and divergent renormalization constants (counterterms). The bare fields may be replaced by renormalized fields. The counterterms are fixed through renormalization condition. These determine the relation between renormalized and physical parameters and can be chosen arbitrarily. The renormalization procedure could be summarized as follows:

- Choose a set of independent parameters.

- Separate the bare parameters and fields into renormalized parameters, fields and renormalization constants.
- Choose renormalization conditions to fix the counterterms.
- Express physical quantities in terms of the renormalized parameters.
- Choose input data in order to fix the values of the renormalized parameters.
- Compute predictions for physical quantities as functions of the input data. The first three steps in the list specify a renormalization scheme.

In this chapter, we are using on-shell renormalization scheme, in which one chooses counterterms so that the finite renormalized parameters are equal to physical parameters in all orders of perturbation theory. The beauty of on shell renormalization scheme is that all parameters (of the electroweak SM) have clear physical significances and can be measured directly in experiments. In the electroweak SM, we choose the masses of the physical particles M_W , M_Z , M_H , m_f , electric charge e , and the quark mixing matrix V_{ij} as renormalized parameters.

2.2.1 Renormalization Constants and Counterterms

We choose the physical parameters $\{e, M_W, M_Z, M_h, m_{f,i}, V_{ij}\}$ as independent parameters. The renormalized quantities and the renormalization constants are defined as follows (bare quantities are denoted by an subscript 0):

$$\begin{aligned}
 e_0 &= Z_e e = (1 + \delta Z_e) e, \\
 M_{W,0}^2 &= M_W^2 + \delta M_W^2, \\
 M_{Z,0}^2 &= M_Z^2 + \delta M_Z^2, \\
 M_{H,0}^2 &= M_H^2 + \delta M_H^2, \\
 m_{f,i,0} &= m_{f,i} + \delta m_{f,i} \\
 V_{ij,0} &= (U_1 V U_2^\dagger)_{ij} = V_{ij} + \delta V_{ij}.
 \end{aligned} \tag{2.27}$$

where U_1 and U_2 are unitary since $V_{ij,0}$ and V_{ij} are unitary.

The counterterms defined above are sufficient to guarantee all S -matrix elements finite, but it leaves Green function divergent. This is because of the fact that radiative corrections change the normalization of the fields by an infinite amount. In order to get finite Green functions we must renormalize the fields as well. Furthermore, radiative corrections yield nondiagonal corrections to the mass matrices such that the bare fields are no more mass eigenstates. In order to re-diagonalized the mass matrices one has to introduce matrix valued field renormalization constants, allowing to define the renormalized fields in such a way that they are the correct physical mass eigenstates in all orders of the perturbation theory. Therefore, we define renormalized

fields as follows:

$$\begin{aligned}
W_0^\pm &= Z_W^{\frac{1}{2}} W^\pm = (1 + \frac{1}{2} \delta Z_W) W^\pm, \\
\begin{pmatrix} Z_0 \\ A_0 \end{pmatrix} &= \begin{pmatrix} Z_{ZZ}^{\frac{1}{2}} & Z_{ZA}^{\frac{1}{2}} \\ Z_{AZ}^{\frac{1}{2}} & Z_{AA}^{\frac{1}{2}} \end{pmatrix} \begin{pmatrix} Z \\ A \end{pmatrix} = \begin{pmatrix} 1 + \frac{1}{2} Z_{ZZ} & \frac{1}{2} Z_{ZA} \\ \frac{1}{2} Z_{AZ} & 1 + \frac{1}{2} Z_{AA} \end{pmatrix} \begin{pmatrix} Z \\ A \end{pmatrix}, \\
H_0 &= Z_H^{\frac{1}{2}} H = (1 + \frac{1}{2} \delta Z_H) H, \\
f_{i,0}^L &= Z_{ij}^{f,L} f_j^L = (\delta_{ij} + \frac{1}{2} \delta Z_{ij}^{f,L}) f_j^L, \\
f_{i,0}^R &= Z_{ij}^{\frac{1}{2},f,L} f_j^R = (\delta_{ij} + \frac{1}{2} \delta Z_{ij}^{f,R}) f_j^R. \tag{2.28}
\end{aligned}$$

Here we do not discuss the renormalization constants of the unphysical ghost and Higgs fields since they do not affect Green functions of physical particles and the calculation of physical one-loop amplitudes.


In writing $Z = 1 + \delta Z$ we could split the bare Lagrangian L_0 into the basic Lagrangian and the counterterm Lagrangian δL

$$L_0 = L + \delta L. \tag{2.29}$$

L shares the same form as L_0 but depends on renormalized parameters and fields. δL stands for counterterms, which absorbs the divergences and unobservable shifts. The corresponding Feynman rules are list in Appendix.A.

2.2.2 Renormalization Conditions

The renormalization constants described above need to be fixed by imposing renormalization conditions. These consist of two sets: the conditions defining the renormalized parameters and the ones defining the renormalized fields.



$$f_j \xrightarrow{p} \text{circle} \xrightarrow{f_i} = \hat{\Gamma}^{ij}(p)$$

$$= i\delta_{ij}(\not{p} - m_{f,i}) + i[\not{p}\omega_- \hat{\Sigma}_{ij}^{f,L}(p^2) + \not{p}\omega_+ \hat{\Sigma}_{ij}^{f,R}(p^2) + (m_{f,i}\omega_- + m_{f,j}\omega_+) \hat{\Sigma}_{i,j}^{f,S}(p^2)].$$

The propagators are obtained as the inverse of the corresponding two-point functions.

The renormalized mass parameters of the physical particles are fixed in such a way that they are equal to the physical masses. For mass matrices, these conditions must be realized by the corresponding eigenvalues, which might result in complicated expressions. These expressions could be simplified by requiring simultaneously the on-shell conditions for the field renormalization matrices. If the external lines are on their mass shell, the renormalized 1PI two-point functions are diagonal. This determines the nondiagonal elements of field renormalization matrices. The renormalized diagonal elements are fixed so that the residues of the renormalized propagators are equal to one. By this choice of field renormalization, the renormalization conditions for the mass parameter require only the corresponding self-energies. Therefore the renormalization conditions for the two-point functions for on-shell external physical fields are defined as follows:

$$\begin{aligned} \tilde{\mathfrak{R}}\hat{\Gamma}_{\mu\nu}^W \epsilon^\nu(k)|_{k^2=M_W^2} &= 0, & \tilde{\mathfrak{R}}\hat{\Gamma}_{\mu\nu}^{ZZ} \epsilon^\nu(k)|_{k^2=M_Z^2} &= 0, & \tilde{\mathfrak{R}}\hat{\Gamma}_{\mu\nu}^{AZ} \epsilon^\nu(k)|_{k^2=M_Z^2} &= 0 \\ \hat{\Gamma}_{\mu\nu}^{AZ} \epsilon^\nu(k)|_{k^2=0} &= 0, & \hat{\Gamma}_{\mu\nu}^{AA} \epsilon^\nu(k)|_{k^2=0} &= 0, \\ \lim_{k^2 \rightarrow M_W^2} \frac{1}{k^2 - M_W^2} \tilde{\mathfrak{R}}\hat{\Gamma}_{\mu\nu}^W \epsilon^\nu(k) &= -i\epsilon_\mu(k), \\ \lim_{k^2 \rightarrow M_Z^2} \frac{1}{k^2 - M_Z^2} \tilde{\mathfrak{R}}\hat{\Gamma}_{\mu\nu}^{ZZ} \epsilon^\nu(k) &= -i\epsilon_\mu(k), & \lim_{k^2 \rightarrow 0} \frac{1}{k^2} \tilde{\mathfrak{R}}\hat{\Gamma}_{\mu\nu}^{AA} \epsilon^\nu(k) &= -i\epsilon_\mu(k), \end{aligned}$$

$$\begin{aligned}
\Re \hat{\Gamma}^H(k)|_{k^2=M_h^2} &= 0, & \lim_{k^2 \rightarrow M_H^2} \frac{1}{k^2 - M_H^2} \Re \hat{\Gamma}^H(k) &= -i, \\
\tilde{\Re} \hat{\Gamma}_{ij}^f(p) u_j(p)|_{p^2=m_{f,j}^2} &= 0, & \tilde{\Re} \bar{u}_j(p') \hat{\Gamma}_{ij}^f(p')|_{p'^2=m_{f,i}^2} &= 0, \\
\lim_{p^2 \rightarrow m_{f,i}^2} \frac{\not{p} + m_{f,i}}{p^2 - m_{f,i}^2} \tilde{\Re} \hat{\Gamma}_{ii}^f(p) u_i(p) &= i u_i(p), \\
\lim_{p^2 \rightarrow m_{f,i}^2} \bar{u}_i(p') \tilde{\Re} \hat{\Gamma}_{ii}^f(p') \frac{\not{p}' + m_{f,i}}{p'^2 - m_{f,i}^2} &= i u_i(p), \tag{2.30}
\end{aligned}$$

where $\epsilon(k)$, $u(p)$ and $\bar{u}(p')$ are the polarization vectors and spinors of the external fields. $\tilde{\Re}$ only takes the real part of the loop integrals appearing in the self-energies.

From the equations above we get the conditions for the self-energy functions.

$$\begin{aligned}
\tilde{\Re} \hat{\Sigma}_T^W(M_W^2) &= 0, & \Re \hat{\Sigma}_T^{ZZ}(M_Z^2) &= 0, & \Re \hat{\Sigma}_T^{AZ}(M_Z^2) &= 0, \\
\hat{\Sigma}_T^{AZ} &= 0, & \hat{\Sigma}_T^{AA} &= 0, \\
\tilde{\Re} \frac{\partial \hat{\Sigma}_T^W(k^2)}{\partial k^2} \Big|_{k^2=M_W^2} &= 0, & \Re \frac{\partial \hat{\Sigma}_T^{ZZ}(k^2)}{\partial k^2} \Big|_{k^2=M_Z^2} &= 0, & \Re \frac{\partial \hat{\Sigma}_T^{AA}(k^2)}{\partial k^2} \Big|_{k^2=0} &= 0, \tag{2.31}
\end{aligned}$$

$$\Re \hat{\Sigma}^H(M_H^2) = 0, \quad \Re \frac{\partial \hat{\Sigma}_T^H(k^2)}{\partial k^2} \Big|_{k^2=M_H^2} = 0, \tag{2.32}$$

$$\begin{aligned}
m_{f,j} \tilde{\Re} \hat{\Sigma}_{ij}^{f,L}(m_{f,j}^2) + m_{f,j} \tilde{\Re} \hat{\Sigma}_{ij}^{f,L}(m_{f,j}^2) &= 0, \\
m_{f,j} \tilde{\Re} \hat{\Sigma}_{ij}^{f,R}(m_{f,i}^2) + m_{f,i} \tilde{\Re} \hat{\Sigma}_{ij}^{f,S}(m_{f,j}^2) &= 0, \\
\tilde{\Re} \hat{\Sigma}_{ii}^{f,R}(m_{f,i}^2) + \tilde{\Re} \hat{\Sigma}_{ii}^{f,R}(m_{f,i}^2) \\
+ 2m_{f,i}^2 \frac{\partial}{\partial p^2} [\tilde{\Re} \hat{\Sigma}_{ii}^{f,R}(p^2) + \tilde{\Re} \hat{\Sigma}_{ii}^{f,L}(p^2) + 2\tilde{\Re} \hat{\Sigma}_{ii}^{f,S}(p^2)] \Big|_{p^2=m_{f,i}^2} &= 0. \tag{2.33}
\end{aligned}$$

Note that the longitudinal (unphysical) components of the gauge boson self-energies drops out for on-shell external gauge bosons.

For the quark mixing matrix V_{ij} , to the lowest order we have

$$V_{0,ij} = U_{ik}^{u,L} U_{i,0}^{d,L,\dagger}, \tag{2.34}$$

where the matrices $U^{f,L}$ transform the weak interaction eigenstates f'_0 to the lowest order mass eigenstates f_0

$$U_{ij}^{f,L,\dagger} f_{j,0}^L = f_{i,0}^L. \quad (2.35)$$

In the on-shell scheme, the higher order mass eigenstates are related to the bare mass eigenstates in the following way

$$f_i^L = Z_{ij}^{\frac{1}{2},f,L} f_{j,0}^L. \quad (2.36)$$


The renormalized quark mixing matrix is defined through the rotation from the weak interaction eigenstates to the renormalized mass eigenstates. In the one-loop level, the rotation in the fermion wave function renormalization $1 + \frac{1}{2}\delta Z^L$ is given by the anti-Hermitian part δZ^{AH} of δZ^L

$$\delta Z_{ij}^{f,AH} = \frac{1}{2}(\delta Z_{ij}^{f,L} - \delta Z_{ij}^{f,L,\dagger}) \quad (2.37)$$

Therefore the renormalized quark mixing matrix is defined as

$$V_{ij} = \left(\delta_{ik} + \frac{1}{2}\delta Z_{ik}^{u,AH,\dagger} \right) V_{0,kn} \left(\delta_{nj} + \frac{1}{2}\delta Z_{nj}^{d,AH,\dagger} \right). \quad (2.38)$$


At last, the electric charge is defined as the full $ee\gamma$ -coupling for on-shell external particles in the Thomson limit in which all vertex corrections vanish on shell and for zero momentum transfer.



Because of our choice the field renormalization, the correction in the external legs vanish and we have the condition

$$\bar{u}(p)\hat{\Gamma}_\mu^{ee\gamma}u(p)|_{p^2=m_e^2} = ie\bar{u}(p)\gamma u(p), \quad (2.39)$$

for the amputated vertex function



2.2.3 Explicit Form of Renormalization Constants

Next, we will give the explicit expressions of renormalization constants.

From eqs. (2.31) and (2.32), we get for the gauge boson sector

$$\begin{aligned} \delta M_W^2 &= \tilde{\Re}\Sigma_T^W(M_W^2), & \delta Z_W &= \tilde{\Re}\frac{\partial\Sigma_T^W(k^2)}{\partial k^2}|_{k^2=M_W^2}, \\ \delta M_Z^2 &= \Re\Sigma_T^{ZZ}(M_Z^2), & \delta Z_{ZZ} &= \tilde{\Re}\frac{\partial\Sigma_T^{ZZ}(k^2)}{\partial k^2}|_{k^2=M_Z^2}, \\ \delta Z_{AZ} &= -2\Re\frac{\Sigma_T^{AZ}(M_Z^2)}{M_Z^2}, & \delta Z_{ZA} &= -2\Re\frac{\Sigma_T^{AZ}(0)}{M_Z^2}, & \delta Z_{AA} &= -\frac{\Sigma_T^{AZ}(k^2)}{k^2} \\ \delta M_H^2 &= \Re\Sigma^H(M_H^2), & \delta Z_H &= -\Re\frac{\partial\Sigma^H(k^2)}{\partial k^2}|_{k^2=M_H^2}. \end{aligned} \quad (2.40)$$

From eq. (2.33) we obtain for the fermion sector

$$\begin{aligned} \delta m_{f,i} &= \frac{m_{f,i}}{2}\tilde{\Re}[\Sigma_{ii}^{f,L}(m_{f,i}^2) + \Sigma_{ii}^{f,R}(m_{f,i}^2) + \Sigma_{ii}^{f,S}(m_{f,i}^2)], \\ \delta Z_{ij}^{f,L} &= \frac{2}{m_{f,i}^2 - m_{f,j}^2}\tilde{\Re}[m_{f,j}^2\Sigma_{ij}^{f,L}(m_{f,j}^2) + m_{f,i}m_{f,j}\Sigma_{ij}^{f,R}(m_{f,j}^2) \\ &\quad + (m_{f,i}^2 + m^2)f, j\Sigma_{ij}^{f,S}(m_{f,j}^2)], \quad i \neq j \\ \delta Z_{ij}^{f,R} &= \frac{2}{m_{f,i}^2 - m_{f,j}^2}\tilde{\Re}[m_{f,j}^2\Sigma_{ij}^{f,R}(m_{f,j}^2) + m_{f,i}m_{f,j}\Sigma_{ij}^{f,L}(m_{f,j}^2) + 2m_{f,i}m_{f,j}\Sigma_{ij}^{f,S}(m_{f,j}^2)], \\ &\quad i \neq j \end{aligned}$$

$$\begin{aligned}
\delta Z_{ii}^{f,L} &= -\tilde{\Re}\Sigma_{ii}^{f,L}(m_{f,i}^2) - m_{f,i}^2 \frac{\partial}{\partial p^2} \tilde{\Re}[\Sigma_{ii}^{f,L}(p^2) + \Sigma_{ii}^{f,R}(p^2) + \Sigma_{ii}^{f,S}(p^2)]|_{p^2=m_{f,i}^2} \\
\delta Z_{ii}^{f,R} &= -\tilde{\Re}\Sigma_{ii}^{f,R}(m_{f,i}^2) - m_{f,i}^2 \frac{\partial}{\partial p^2} \tilde{\Re}[\Sigma_{ii}^{f,L}(p^2) + \Sigma_{ii}^{f,R}(p^2) + \Sigma_{ii}^{f,S}(p^2)]|_{p^2=m_{f,i}^2}. \quad (2.41)
\end{aligned}$$

The use of $\tilde{\Re}$ guarantees that the renormalized Lagrangian is real. Moreover we have

$$\delta Z_{ij}^\dagger = \delta Z_{ij}(m_i^2 \leftrightarrow m_j^2). \quad (2.42)$$

The renormalization constant for the quark mixing matrix V_{ij} can be derived from eq. (2.38)

$$\delta V_{ij} = \frac{1}{4}[(\delta Z_{ik}^{u,L} - \delta Z_{ik}^{u,L,\dagger}) - V_{ik}(\delta Z_{kj}^{d,L} - \delta Z_{kj}^{d,L,\dagger})]. \quad (2.43)$$

Inserting eq. (2.41) we obtain

$$\begin{aligned}
\delta V_{ij} &= \frac{1}{2} \tilde{\Re} \left\{ \frac{1}{m_{u,i}^2 - m_{u,k}^2} [m_{u,k}^2 \Sigma_{ik}^{u,L}(m_{u,k}^2) + m_{u,i}^2 \Sigma_{ik}^{u,L}(m_{u,i}^2) \right. \\
&\quad + m_{u,i} m_{u,k} (\Sigma_{ik}^{u,R}(m_{u,k}^2) + \Sigma_{ik}^{u,R}(m_{u,i}^2)) \\
&\quad + (m_{u,k}^2 + m_{u,i}^2) (\Sigma_{ik}^{u,S}(m_{u,k}^2) + \Sigma_{ik}^{u,S}(m_{u,i}^2))] V_{kj} \\
&\quad - V_{ik} \frac{1}{m_{d,k}^2 - m_{d,j}^2} [m_{d,j}^2 \Sigma_{kj}^{d,L}(m_{d,j}^2) + m_{d,k}^2 \Sigma_{kj}^{d,L}(m_{d,k}^2) \\
&\quad + m_{d,k} m_{d,j} (\Sigma_{kj}^{d,R}(m_{d,j}^2) + \Sigma_{kj}^{d,R}(m_{d,k}^2)) \\
&\quad \left. + (m_{d,k}^2 + m_{d,j}^2) (\Sigma_{kj}^{d,S}(m_{d,k}^2) + \Sigma_{kj}^{d,S}(m_{d,j}^2))] \right\}. \quad (2.44)
\end{aligned}$$

Next, we will determine the charge renormalization δZ_e from the $ee\gamma$ -vertex. For generalization, we explore the $ff\gamma$ -vertex for arbitrary fermions f . The renormalized vertex function is

$$\hat{\Gamma}_{ij,\mu}^{\gamma ff}(p, p') = -ie\delta_{ij} Q_f \gamma_\mu + ie\hat{\Lambda}_{ij,\mu}^{\gamma ff}(p, p'). \quad (2.45)$$

For on-shell external fermions it can be decomposed as ($k = p' - p$)

$$\hat{\Lambda}_{ij,\mu}^{\gamma ff}(p, p') = \delta_{ij} \left(\gamma_\mu \hat{\Lambda}_V^f(k^2) - \gamma_\mu \gamma_5 \hat{\Lambda}_A^f(k^2) + \frac{(p+p')_\mu}{2m_f} \hat{\Lambda}_S^f(k^2) + \frac{(p'-p)_\mu}{2m_f} \gamma_5 \hat{\Lambda}_S^f(k^2) \right). \quad (2.46)$$

According to eq. (2.39), we obtain

$$\begin{aligned} 0 &= \bar{u}(p) \hat{\Lambda}_{ii,\mu}^{\gamma ff}(p, p) U(p) \\ &= \bar{u}(p) \gamma_\mu u(p) [-Q_f (\delta Z_e + \delta Z_{ii}^{f,V} + \frac{1}{2} \delta Z_{AA}) + \Lambda_V^f(0) + \Lambda_S^f(0) + v_f \frac{1}{2} \delta Z_{ZA}] \\ &\quad - \bar{u}(p) \gamma_\mu \gamma_5 u(p) [-Q_f \delta Z_{ii}^{f,A} + \Lambda_A^f(0) + a_f \frac{1}{2} \delta Z_{ZA}], \end{aligned} \quad (2.47)$$

where

$$\delta Z_{ii}^{f,V} = \frac{1}{2} (\delta Z_{ii}^{f,L} + \delta Z_{ii}^{f,R}), \quad \delta Z_{ii}^{f,A} = \frac{1}{2} (\delta Z_{ii}^{f,L} - \delta Z_{ii}^{f,R}), \quad (2.48)$$

and v_f, a_f are the vector and axialvector couplings of the Z -boson to the fermion f .

From eq. (2.47), we have two conditions

$$0 = -Q_f \left(\delta Z_e + \delta Z_i^{f,V} i + \frac{1}{2} \delta Z_{AA} \right) + \Lambda_V^f(0) + \Lambda_S^f(0) + v_f \frac{1}{2} \delta Z_{ZA}, \quad (2.49)$$

$$0 = -Q_f \delta Z_{ii}^{f,A} + \Lambda_A^f(0) + a_f \frac{1}{2} \delta Z_{ZA}. \quad (2.50)$$

The eq. (2.49) for $f = e$ fixes the charge renormalization constant. The eq. (2.50) is fulfilled because of a Ward identity (derived from gauge invariance). Furthermore the same Ward identity yields

$$\Lambda_V^f(0) + \Lambda_S^f(0) - Q_f \delta Z_{ii}^{f,V} + a_f \frac{1}{2} \delta Z_{ZA} = 0. \quad (2.51)$$

Inserting this equation we finally obtain (using $v_f - a_f = -Q_f \frac{s_W}{c_W}$)

$$\begin{aligned} \delta Z_e &= \frac{1}{2} \delta Z_{AA} - \frac{1}{2} \frac{s_W}{c_W} \delta Z_{ZA} \\ &= \frac{1}{2} \frac{\partial \Lambda_T^{AA}(k^2)}{\partial k^2} \Big|_{k^2=0} - \frac{s_W}{c_W} \frac{\Sigma_T^{AZ}(0)}{M_Z^2}. \end{aligned} \quad (2.52)$$

This result is independent of the fermion species, reflecting electric charge universality.

In the on-shell scheme the weak mixing angle is a derived quantity. It is defined as [40–42]

$$\sin^2 \theta_W = s_W^2 = 1 - \frac{M_W^2}{M_Z^2}, \quad (2.53)$$

using the renormalized gauge boson masses. This definition is process-independent and valid to all orders of perturbation theory. It is convenient to introduce the corresponding counterterms

$$c_{W,0} = c_W + \delta c_W, \quad s_{W,0} = s_W + \delta s_W, \quad (2.54)$$

which are directly related to the counterterms to the gauge boson mass due to eq. (2.53). To one-loop order we have

$$\begin{aligned} \frac{\delta c_W}{c_W} &= \frac{1}{2} \left(\frac{\delta M_W^2}{M_W^2} - \frac{\delta M_Z^2}{M_Z^2} \right) = \frac{1}{2} \tilde{\Re} \left(\frac{\Sigma_T^W(M_W^2)}{M_W^2} - \frac{\Sigma_T^{ZZ}(M_Z^2)}{M_Z^2} \right) \\ \frac{\delta s_W}{s_W} &= -\frac{c_W^2}{s_W^2} \frac{\delta c_W}{c_W} = -\frac{1}{2} \frac{c_W^2}{s_W^2} \tilde{\Re} \left(\frac{\Sigma_T^W(M_W^2)}{M_W^2} - \frac{\Sigma_T^{ZZ}(M_Z^2)}{M_Z^2} \right). \end{aligned} \quad (2.55)$$

We have now got all renormalization constants in terms of unrenormalized self-energies. In the next section, we will introduce the methods to calculate to one-loop radiative corrections.

2.3 One-Loop Integrals

Perturbative calculations at one-loop level involve complicated integrals over the loop momentum (scalar, vector and tensor integrals). In this section, we will introduce the basic modern tools for the calculation of loop diagrams [30–32, 43], in which all one-loop integrals can be reduced to the scalar ones.

2.3.1 Scalar One-loop Integrals for $N \leq 4$

We first introduce the basic scalar one-loop integrals A_0 , B_0 , C_0 and D_0 , which were derived in [43].

We begin by introducing the scalar one-point function

$$\begin{aligned}
A_0(m) &= \frac{(2\pi\mu)^{4-n}}{i\pi^2} \int d^d q \frac{1}{q^2 - m^2 + i\epsilon} \\
&= -m^2 \left(\frac{m^2}{4\pi\mu^2} \right)^{\frac{d-4}{2}} \Gamma\left(1 - \frac{D}{2}\right) \\
&= m^2 \left(\Delta - \log \frac{m^2}{\mu^2} + 1 \right), \tag{2.56}
\end{aligned}$$

where the UV-divergence is contained in

$$\Delta = \frac{2}{4-d} - \gamma_E + \log 4\pi \tag{2.57}$$

and γ_E is the Euler's constant.

The scalar two-point function is given by

$$\begin{aligned}
B_0(p_{10}, m_0, m_1) &= \frac{(2\pi\mu)^{4-n}}{i\pi^2} \int d^d q \frac{1}{[q^2 - m_0^2 + i\epsilon][(q + p_{10})^2 - m_1^2 + i\epsilon]} \\
&= \Delta + 2 - \log \frac{m_0 m_1}{\mu} + \frac{m_0^2 - m_1^2}{p_{10}^2} \log \frac{m_1}{m_0} - \frac{m_0 m_1}{p_{10}^2} \left(\frac{1}{r} - r \right) \log r \tag{2.58}
\end{aligned}$$

where r and $\frac{1}{r}$ are determined from

$$x^2 + \frac{m_0^2 + m_1^2 - p_{10}^2 - i\epsilon}{m_0 m_1} x + 1 = (x + r) \left(x + \frac{1}{r} \right). \tag{2.59}$$

For the field renormalization constants, the derivative of B_0 with respect to p_{10}^2 is required. It is given by

$$\begin{aligned}
\frac{\partial}{\partial p_{10}^2} B_0(p_{10}, m_0, m_1) &= -\frac{m_0^2 - m_1^2}{p_{10}^4} \log \frac{m_1}{m_0} + \frac{m_0 m_1}{p_{10}^4} \left(\frac{1}{r} - r \right) \log r \\
&\quad - \frac{1}{p_{10}^2} \left(1 + \frac{r^2 + 1}{r^2 - 1} \log r \right). \tag{2.60}
\end{aligned}$$

The scalar three-point reads

$$C_0(p_{10}, p_{20}, m_0, m_1, m_2) = \frac{(2\pi\mu)^{4-n}}{i\pi^2} \int d^d q \frac{1}{D_0 D_1 D_2}, \tag{2.61}$$

where

$$D_0 = q^2 - m_0^2 + i\epsilon; \quad D_1 = (q + p_1)^2 - m_1^2 + i\epsilon; \quad D_2 = (q + p_2)^2 - m_2^2 + i\epsilon \quad (2.62)$$

In order to compute three-point function, we need to introduce two Feynman parameters. The general result for scalar three-point function valid for all real momenta and physical masses was calculated by [43]. It can be also expressed into symmetric form

$$\begin{aligned} C_0(p_{10}, p_{20}, m_0, m_1, m_2) &= - \int_0^1 dx \int_0^x dy [p_{21}^2 x^2 + p_{10}^2 y^2 + (p_{20}^2 - p_{10}^2 - p_{21}^2)xy \\ &\quad + (m_1^2 - m_2^2 - p_{21}^2)x + (m_0^2 + m_1^2 + p_{21}^2 - p_{20}^2)y + m_2^2 - i\epsilon]^{-1} \\ &= \frac{1}{\alpha} \sum_{i=0}^2 \left\{ \sum_{\sigma=\pm} \left[Li_2 \left(\frac{y_{0i} - 1}{y_{i\sigma}} \right) - Li_2 \left(\frac{y_{0i}}{y_{i\sigma}} \right) + \eta \left(1 - x_{i\sigma}, \frac{1}{y_{i\sigma}} \right) \log \frac{y_{0i} - 1}{y_{i\sigma}} \right. \right. \\ &\quad \left. \left. - \eta \left(-x_{i\sigma}, \frac{1}{y_{i\sigma}} \right) \log \frac{y_{0i}}{y_{i\sigma}} \right] - [\eta(-x_{i+}, -x_{i-}) - \eta(y_{i+}, y_{i-}) - 2\pi i \theta(-p_{jk}^2) \right. \right. \\ &\quad \left. \left. \theta(-\Im(y_{i+}y_{i-})) \right] \log \frac{1 - y_{i0}}{y_{i\sigma}} \right\}, \end{aligned} \quad (2.63)$$

where

$$\begin{aligned} p_{ij} &= p_i - p_j, \quad p_{i0} = p_i, \\ y_{0i} &= \frac{1}{2\alpha p_{jk}^2} [p_{jk}^2 (p_{jk}^2 - p_{ki}^2 - p_{ij}^2 + 2m_i^2 - m_j^2 - m_k^2), \\ &\quad - (p_{ki}^2 - p_{ij}^2)(m_k^2 - m_k^2) + \alpha(p_{jk}^2 - m_j^2 + m_k^2)], \\ x_{i\pm} &= \frac{1}{2p_{jk}^2} [p_{jk}^2 - m_j^2 + m_k^2 \pm \alpha_i], \quad y_{i\pm} = y_{0i} - x_{i\pm}, \\ \alpha &= \kappa(p_{10}^2, p_{21}^2, p_{20}^2), \\ \alpha_i &= \kappa(p_{ij}^2, m_j^2, m_k^2)(1 + i\epsilon p_{jk}^2), \end{aligned} \quad (2.64)$$

and κ is the Kallen function

$$\kappa(x, y, z) = \sqrt{x^2 + y^2 + z^2 - 2(xy + yz + zx)}. \quad (2.65)$$

The Spence function $Li_2(x)$ is defined as

$$Li_2(x) = - \int_0^1 \frac{dt}{t} \log(1 - xt), \quad |arg(1 - x)| < \pi. \quad (2.66)$$

The η -function is defined as

$$\eta(a, b) = 2i\pi[\theta(-\Im a)\theta(-\Im b)\theta(-\Im ab) - \theta(\Im a)\theta(-\Im b)\theta(-\Im ab)] \quad (2.67)$$

All η -functions in eq. (2.63) vanish if α and all the masses m_i are real.

Next, let us investigate the scalar four-point function $D_0(p_{10}, p_{20}, p_{30}, m_0, m_1, m_2, m_3)$, which can be expressed in terms of 16 dilogarithms [44] instead of 24 dilogarithms of the result calculated by [43].

The scalar four-point integral can be expressed in the symmetric form

$$D_0(p_{10}, p_{20}, p_{30}, m_0, m_1, m_2, m_3) = \frac{(2\pi\mu)^{4-n}}{i\pi^2} \int d^d q \frac{1}{D_0 D_1 D_2 D_3}, \quad (2.68)$$

where

$$\begin{aligned} D_0 &= q^2 - m_0^2 + i\epsilon, & D_1 &= (q + p_1)^2 - m_1^2 + i\epsilon, \\ D_2 &= (q + p_2)^2 - m_2^2 + i\epsilon, & D_3 &= (q + p_3)^2 - m_3^2 + i\epsilon \end{aligned} \quad (2.69)$$

We first give some variables and functions before we exhibit the result. We define

$$k_{ij} = \frac{m_i^2 + m_j^2 - p_{ij}^2}{m_i m_j}, \quad 1 \leq i < j \leq 4. \quad (2.70)$$

The quantities r_{ij} and \tilde{r}_{ij} are defined by

$$\begin{cases} x^2 + k_{ij}x + 1 = (x + r_{ij})(x + 1/r_{ij}), \\ x^2 + (k_{ij} - i\epsilon)x + 1 = (x + \tilde{r}_{ij})(x + 1/\tilde{r}_{ij}). \end{cases} \quad (2.71)$$

Note that for real k_{ij} the r_{ij} 's lie either on the real axis or on the complex unit circle.

In addition,

$$\begin{aligned}
P(y_0, y_1, y_2, y_3) &= \sum_{0 \leq i < j \leq 3} k_{ij} y_i y_j + \sum_{j=0}^3 y_j^2, \\
Q(y_0, y_1, 0, y_3) &= (1/r_{02} - r_{02})y_0 + (k_{12} - r_{02}k_{01})y_1 + (k_{23} - r_{02}k_{03})y_3, \\
Q(y_0, 0, y_2, y_3) &= (1/r_{13} - r_{13})y_3 + (k_{12} - r_{13}k_{23})y_2 + (k_{01} - r_{13}k_{03})y_0.
\end{aligned} \tag{2.72}$$

and $x_{1,2}$ is determined by

$$\begin{aligned}
ax^2 + bx + c + i\epsilon d = \frac{r_{02}r_{13}}{x} \left\{ \left[P\left(1, \frac{x}{r_{13}}, 0, 0\right) - i\epsilon \right] \left[P\left(0, 0, \frac{x}{r_{02}}, x\right) - i\epsilon \right] \right. \\
\left. - \left[P\left(0, \frac{x}{r_{13}}, \frac{1}{r_{02}}, 0\right) - i\epsilon \right] \left[P\left(1, 0, 0, x\right) - i\epsilon \right] \right\},
\end{aligned} \tag{2.73}$$

where

$$\begin{cases} a = k_{23}/r_{13} + r_{02}k_{01} - k_{03}r_{02}/r_{13} - k_{12}, \\ b = (r_{13} - 1/r_{13})(r_{02} - 1/r_{02}) + k_{01}k_{23} - k_{03}k_{12}, \\ c = k_{01}/r_{02} + r_{13}k_{23} - k_{03}r_{13}/r_{02} - k_{12}, \\ d = k_{12} - r_{02}k_{01} - r_{13}k_{23} + r_{02}r_{13}k_{03}. \end{cases} \tag{2.74}$$

Furthermore, we introduce

$$\gamma_{kl} = \text{sgn} \Re[a(x_k - x_l)], \quad k, l = 1, 2, \tag{2.75}$$

$$\begin{cases} x_{k0} = x_k, & s_0 = \tilde{r}_{03} \\ x_{k1} = x_k/r_{13}, & s_1 = \tilde{r}_{01} \\ x_{k2} = x_k r_{02}/r_{13}, & s_2 = \tilde{r}_{12} \\ x_{k3} = x_k r_{02}, & s_3 = \tilde{r}_{23} \end{cases} \tag{2.76}$$

and

$$x_{kj}^{(0)} = \lim_{\epsilon \rightarrow 0} x_{kj} \quad \text{as} \quad r_{ij} = \lim_{\epsilon \rightarrow 0} \tilde{r}_{ij} \quad (2.77)$$

At last, we introduce

$$\tilde{\eta}(a, \tilde{b}) = \begin{cases} \eta(a, b) & \text{for } b \text{ not real,} \\ 2\pi i [\theta(-Ima)\theta(-Im\tilde{b}) - \theta(Ima)\theta(Im\tilde{b})] & \text{for } b < 0, \\ 0 & \text{for } b > 0, \end{cases} \quad (2.78)$$

with $b = \lim_{\epsilon \rightarrow 0} \tilde{b}$.

Then we have the result for real r_{02}

$$\begin{aligned} D_0(p_{10}, p_{20}, p_{30}, m_0, m_1, m_2, m_3) &= \frac{1}{m_1 m_2 m_3 m_4 a (x_1 - x_2)} \\ &\times \left\{ \sum_{j=0}^3 \sum_{k=1}^2 (-1)^{j+k} \left[Li_2(1 + s_j x_{kj}) + \eta(-x_{kj}, s_j) \log(1 + s_j x_{kj}) \right. \right. \\ &+ Li_2\left(1 + \frac{x_{kj}}{s_j}\right) + \eta\left(-x_{kj}, \frac{1}{s_j}\right) \log\left(1 + \frac{x_{kj}}{s_j}\right) \left. \right] \\ &+ \sum_{k=1}^2 (-1)^{k+1} \left[\tilde{\eta}\left(-x_{kj}, \frac{1}{s_j}\right) \left[\log(r_{02} x_k) + \log\left(Q\left(\frac{1}{x_k^{(0)}}, 0, 0, 1\right) - i\epsilon\right) \right. \right. \\ &+ \log\left(\frac{\bar{Q}(0, 0, 1, r_{02} x_k^{(0)})}{d} + i\epsilon \gamma_{k,3-k} \operatorname{sgn}(r_{02} Im\tilde{r}_{13}) \right) \left. \right] \\ &- \tilde{\eta}\left(-x_k, \frac{1}{\tilde{r}_{13}}\right) \left[\log\left(\frac{x_k}{r_{13}}\right) + \log\left(Q\left(\frac{r_{13}}{x_k^{(0)}}, 1, 0, 0\right) - i\epsilon\right) \right. \\ &+ \log\left(\frac{\bar{Q}(1, 0, 0, x_k^{(0)})}{d} + i\epsilon \gamma_{k,3-k} \operatorname{sgn}(Im\tilde{r}_{13}) \right) \left. \right] \\ &- \left[\tilde{\eta}\left(-x_k, \frac{\tilde{r}_{02}}{\tilde{r}_{13}}\right) + \tilde{\eta}\left(\tilde{r}_{02}, \frac{1}{\tilde{r}_{13}}\right) \right] \left[\log\left(\frac{r_{02} x_k}{r_{13}}\right) + \log\left(Q\left(\frac{r_{13}}{x_k^{(0)}}, 1, 0, 0\right) - i\epsilon\right) \right. \\ &+ \log\left(\frac{\bar{Q}(1, 0, 0, r_{02} x_k^{(0)})}{d} + i\epsilon \gamma_{k,3-k} \operatorname{sgn}(r_{02} Im\tilde{r}_{13}) \right) \left. \right] \\ &+ \left. \eta\left(\tilde{r}_{02}, \frac{1}{\tilde{r}_{13}}\right) \tilde{\eta}\left(-x_k, -\frac{\tilde{r}_{02}}{\tilde{r}_{13}}\right) \right\}. \quad (2.79) \end{aligned}$$

In the case that $|r_{ij}| = 1$ for all r_{ij} , the result can be written as

$$\begin{aligned}
D_0(p_{10}, p_{20}, p_{30}, m_0, m_1, m_2, m_3) &= \frac{1}{m_1 m_2 m_3 m_4 a(x_1 - x_2)} \\
&\left\{ \sum_{j=0}^3 \sum_{k=1}^2 (-1)^{j+k} \left[Li_2(1 + s_j x_{kj}) + \eta(-x_{kj}, s_j) \log(1 + s_j x_{kj}) \right. \right. \\
&+ Li_2\left(1 + \frac{x_{kj}}{s_j}\right) + \eta\left(-x_{kj}, \frac{1}{s_j}\right) \log\left(1 + \frac{x_{kj}}{s_j}\right) \left. \right] \\
&+ \sum_{k=1}^2 (-1)^{k+1} \left[\eta\left(-x_k, \frac{1}{r_{13}}\right) \left[\log\left(\frac{r_{13}}{x_k^{(0)}} P\left(1, \frac{x_k^{(0)}}{r_{[13]}}\right), 0, 0\right) - \frac{x_k^{(0)}}{r_{13}} \epsilon b \gamma_{k,3-k}\right) \right] \\
&+ \eta(-x_k, r_{02}) \left[\log\left(\frac{1}{r_{02} x_k^{(0)}} P(0, 0, 1, r_{02} x_k^{(0)}) - r_{02} x_k^{(0)} \epsilon b \gamma_{k,3-k}\right) + \log(r_{02} x_k^{(0)}) \right] \\
&- \left[\eta\left(-x_k, \frac{r_{02}}{r_{13}}\right) + \eta\left(r_{02}, \frac{1}{r_{13}}\right) \right] \left[\log\left(\frac{r_{[13]}}{r_{02} x_k^{(0)}} P\left(0, 1, \frac{r_{02} x_k^{(0)}}{r_{13}}\right), 0\right) - \frac{r_{02} x_k^{(0)}}{r_{13}} \epsilon b \gamma_{k,3-k}\right) \right. \\
&\left. + \log\left(\frac{r_{02} x_k^{(0)}}{r_{13}}\right) \right] + (1 - \gamma_{k,3-k} \text{sgn}(b)) \eta\left(-x_k, -\frac{r_{02}}{r_{13}}\right) \eta\left(r_{02}, \frac{1}{r_{13}}\right) \left. \right\}.
\end{aligned} \tag{2.80}$$

2.3.2 Tensor Integral Reduction

We have introduced the four basic scalar integrals A_0 , B_0 , C_0 and D_0 for calculations of the perturbative quantum field theory. Besides the scalar integrals we also need tensor integrals in the perturbative calculations, of which the evaluations could be very complicated in practice. In order to deal with these complications, we follow one particular procedure [30,31] in which these tensor structures could be reduced to linear combinations of scalar integrals.

In general, the one-loop integrals in d -dimensions are classified with the the number N of propagators in the denominator and the number P of integration momenta in the numerators. According to power counting, the integrals with $P + D - 2N \geq 0$ are UV-divergent. The divergencies can be regulated by evaluating the integrals in general dimensions $d \neq 4$ (dimensional regularization [15]). The

UV-divergences would be cancelled in the procedure of the renormalization. For renormalizable theories we have $P \leq N$ and therefore a finite number of divergent integrals.

We define the general one-loop tensor integral as


$$T_{\mu_1 \dots \mu_P}^N(p_1, \dots, p_{N_1}, m_0, \dots, m_{N_1}) = \frac{(2\pi\mu)^{4-d}}{i\pi^2} \int d^d q \frac{q_{\mu_1} \dots q_{\mu_P}}{D_0 D_1 \dots D_{N-1}} \quad (2.81)$$

where the denominator factors

$$D_0 = q^2 - m_0^2 + i\epsilon, \quad D_i = (q + p_i)^2 - m_i^2 + i\epsilon, \quad i = 1, \dots, N-1, \quad (2.82)$$

arising from the propagators in the Feynman diagram. Moreover we introduce

$$p_{i0} = p_i \quad \text{and} \quad p_{ij} = p_i - p_j. \quad (2.83)$$



Apparently the tensor integrals are invariant under permutations of the propagators D_i , $i \neq 0$ and totally symmetric in the Lorentz indices μ_k . $i\epsilon$ is an infinitesimal imaginary part which regulates singularities of the integrand. Its specific choice guarantees causality. The parameter μ has mass dimension and play a role to keep the dimension of the integrals fixed for varying d . Conventionally T^N is denoted by the

N th character of alphabet, i.e. $T^1 \equiv A, T^2 \equiv B, \dots$, and the scalar integrals carry a subscript 0.

Lorentz covariance of the integrals allows to decompose tensor integrals into tensors constructed from the external momenta p_i , and the metric tensor $g_{\mu\nu}$ with totally symmetric coefficient functions $T_{i_1 \dots i_P}^N$. Formally we introduce an artificial momentum p_0 to write terms containing $g_{\mu\nu}$ in a compact way

$$T_{\mu_1 \dots \mu_P}^N(p_1, \dots, p_{N_1}, m_0, \dots, m_{N_1}) = \sum_{i_1, \dots, i_P=0}^{N-1} T_{i_1 \dots i_P}^N p_{i_1} p_{i_1 \mu_1} \dots p_{i_P \mu_P}, \quad (2.84)$$

the $g_{\mu\nu}$ terms are recovered by omitting terms containing an odd number of p_0 's and replacing the products of even numbers of p_0 's by the corresponding totally symmetric tensor constructed from the $g_{\mu\nu}$, for example,

$$\begin{aligned} p_{0\mu_1} p_{0\mu_2} &\rightarrow g_{\mu_1 \mu_2} \\ p_{0\mu_1} p_{0\mu_2} p_{0\mu_3} p_{0\mu_4} &\rightarrow g_{\mu_1 \nu_1} g_{\mu_3 \nu_4} + g_{\mu_1 \nu_3} g_{\mu_2 \nu_4} = g_{\mu_1 \nu_4} g_{\mu_2 \nu_3}. \end{aligned} \quad (2.85)$$

The explicit Lorentz decompositions for the lowest order integral are easily derived. The two various two point functions can be written as

$$\begin{aligned} B_\mu &= p_{1\mu} B_1, \\ B_{\mu\nu} &= g_{\mu\nu} B_{00} + p_{1\mu} p_{1\nu} B_{11}, \end{aligned} \quad (2.86)$$

the three point functions reads

$$\begin{aligned} C_\mu &= p_{1\mu} C_1 + p_{2\mu} C_2 = \sum_{i=1}^2 p_{i\mu} C_i, \\ C_{\mu\nu} &= g_{\mu\nu} C_{00} + \sum_{i,j=1}^2 p_{i\mu} p_{j\nu} C_{ij}, \\ C_{\mu\nu\rho} &= \sum_{i=1}^2 (g_{\mu\nu} p_{i\rho} + g_{\nu\rho} p_{i\mu} + g_{\mu\rho} p_{i\nu}) C_{00i} + \sum_{i,j,k=1}^2 p_{i\mu} p_{j\nu} p_{k\rho} C_{ijk}, \end{aligned} \quad (2.87)$$

And the various D functions could be expressed as

$$\begin{aligned}
D_\mu &= \sum_{i=1}^3 p_{i\mu} D_i, & D_{\mu\nu} &= g_{\mu\nu} D_{00} + \sum_{i,j=1}^3 p_{i\mu} p_{j\nu} D_{ij}, \\
D_{\mu\nu\rho} &= \sum_{i=1}^3 (g_{\mu\nu} p_{i\rho} + g_{\nu\rho} p_{i\mu} + g_{\mu\rho} p_{i\nu}) D_{00i} + \sum_{i,j,k=1}^3 p_{i\mu} p_{j\nu} p_{k\rho} D_{ijk} \\
D_{\mu\nu\rho\sigma} &= (g_{\mu\nu} g_{\rho\sigma} + g_{\mu\rho} g_{\nu\sigma} + g_{\mu\sigma} g_{\nu\rho}) D_{0000} \\
&+ \sum_{i,j=1}^3 (g_{\mu\nu} p_{i\rho} p_{j\sigma} + g_{\nu\rho} p_{i\mu} p_{j\sigma} + g_{\mu\rho} p_{i\nu} p_{j\sigma} \\
&+ g_{\mu\sigma} p_{i\nu} p_{j\rho} + g_{\nu\sigma} p_{i\mu} p_{j\rho} + g_{\rho\sigma} p_{i\mu} p_{j\nu}) D_{00ij} \\
&+ \sum_{i,j,k=1}^3 p_{i\mu} p_{j\nu} p_{k\rho} p_{l\sigma} D_{ijkl}
\end{aligned} \tag{2.88}$$

For a general tensor integral with $N \geq 5$, the terms involving $g_{\mu\nu}$ should be omitted since the four dimensional space is spanned by four Lorentz vectors. Furthermore, the decomposition (2.84) should contain at most four Lorentz vectors. Therefore, the decomposition (2.84) arrives at

$$T_{\mu_1 \dots \mu_P}^N(p_1, \dots, p_{N_1}, m_0, \dots, m_{N_1}) = \sum_{i_1, \dots, i_P=0}^4 T_{i_1 \dots i_P}^N p_{i_1 \mu_1} \dots p_{i_P \mu_P}, \tag{2.89}$$

where $\{p_1, \dots, p_4\}$ is any set of four linear independent Lorentz vectors out of $\{p_1, \dots, p_{N-1}\}$.

The symmetry of the tensor integrals under exchange of the propagators gives rise to relations between the scalar coefficient functions. Exchanging the arguments $(p_i, m_i) \leftrightarrow (p_j, m_j)$ together with the corresponding indices $i \leftrightarrow j$ leaves the scalar coefficient functions invariant, for example,

$$\begin{aligned}
C_1(p_1, p_2, m_0, m_1, m_2) &= C_1(p_2, p_1, m_0, m_2, m_1), \\
C_{00}(p_1, p_2, m_0, m_1, m_2) &= C_{00}(p_2, p_1, m_0, m_2, m_1), \\
C_{12}(p_1, p_2, m_0, m_1, m_2) &= C_{12}(p_2, p_1, m_0, m_2, m_1).
\end{aligned} \tag{2.90}$$

All one-loop tensor integrals could be expressed iteratively in terms of the scalar ones $T_0^N(A_0, B_0, C_0, D_0 \dots)$, using the Lorentz decomposition of the tensor integrals. We will derive the general procedure for the tensor integral.

The product of the integration momentum q_μ with an external momentum could be written in terms of the denominators

$$qp_k = \frac{1}{2}[D_k - D_0 - f_k], \quad f_k = p_k^2 - m_k^2 + m_0^2. \quad (2.91)$$

Multiplying eq. (2.81) with p_k and substituting eq. (2.91) yields

$$\begin{aligned} R_{\mu_1 \dots \mu_{P-1}}^{N,k} &= T_{\mu_1 \dots \mu_P}^N p_k^{\mu_P} \\ &= \frac{1}{2} \frac{(2\pi\mu)^{4-d}}{i\pi^2} \int d^d q \left[\frac{q_{\mu_1} \dots q_{\mu_{P-1}}}{D_0 \dots D_{k-1} D_{k+1} \dots D_{N-1}} \right. \\ &\quad \left. - \frac{q_{\mu_1} \dots q_{\mu_{P-1}}}{D_1 \dots D_{N-1}} - f_k \frac{q_{\mu_1} \dots q_{\mu_{P-1}}}{D_1 \dots D_{N-1}} \right] \\ &= \frac{1}{2} [T_{\mu_1 \dots \mu_{P-1}}^{N-1}(k) - T_{\mu_1 \dots \mu_{P-1}}^{N-1}(0) - f_k T_{\mu_1 \dots \mu_{P-1}}^N], \end{aligned} \quad (2.92)$$

where the argument k in $T_{\mu_1 \dots \mu_{P-1}}^{N-1}(k)$ implies that the propagator D_k was eliminated. Note that $T_{\mu_1 \dots \mu_{P-1}}^{N-1}(0)$ has an external momentum in its first propagator. So we need to perform a shift of the integration momentum to recover it to the form (2.81). All tensor integrals on the right-hand side of eq. (2.92) have one Lorentz index less than the original tensor integral. In two of them one propagator is cancelled.

For $P \geq 2$, contracting eq. (2.81) with $g_{\mu\nu}$ and using

$$g^{\mu\nu} q_\mu q_\nu = q^2 = D_0 + m_0^2, \quad (2.93)$$

yields

$$\begin{aligned}
R_{\mu_1 \dots \mu_{P-2}}^{N,00} &= T_{\mu_1 \dots \mu_P}^N g^{\mu_{P-1} \mu_P} \\
&= \frac{(2\pi\mu)^{4d}}{i\pi^2} \int d^d q \left[\frac{q_{\mu_1} \dots q_{\mu_{P-2}}}{D_1 \dots D_N} + m_0^2 \frac{q_{\mu_1} \dots q_{\mu_{P-2}}}{D_0 \dots D_N} \right] \\
&= [T_{\mu_1 \dots \mu_{P-2}}^{N-1}(0) + m_0^2 T_{\mu_1 \dots \mu_{P-2}}^N].
\end{aligned} \tag{2.94}$$

Plugging the Lorentz decomposition (2.84) for the tensor integrals in eqs. (2.92) and (2.94) we obtain a set of linear equations for the corresponding coefficient functions. This set decomposes into disjoint set of $N - 1$ equations for each tensor integral. If the inverse of the matrix

$$X_{N-1} = \begin{pmatrix} p_1^2 & p_1 p_2 & \dots & p_1 p_{N-1} \\ p_2 p_1 & p_2^2 & \dots & p_2 p_{N-1} \\ \vdots & \vdots & \ddots & \vdots \\ p_{N-1} & p_{N-1} p_2 & \dots & p_{N-1}^2 \end{pmatrix} \tag{2.95}$$

exists, these can be solved for the invariant functions $T_{i_1 \dots i_P}^N$. In this way all tensor integrals are reduced iteratively to scalar integrals T_0^L with $L \leq N$.

If the matrix X_{N-1} becomes singular, the reduction algorithm fails. If this is due to the linear dependence of the momenta we can leave out the linear dependent vectors of the set $\{p_1, \dots, p_{N-1}\}$ in the Lorentz decomposition bringing in a smaller matrix X_M . If X_M is nonsingular the reduction algorithm works again. This occurs at the edge of phase space where some of the momenta p_i become collinear.

Now we exhibit the results for reduction of arbitrary N -point integrals depending on $M \leq N - 1$ linear independent Lorentz vectors in d dimensions for nonsingular

matrix X_M . Inserting the Lorentz decomposition of T_N , $R^{N,K}$ and $R^{N,00}$

$$\begin{aligned}
R_{\mu_1 \dots \mu_{P-1}}^{N,K} &= T_{\mu_1 \dots \mu_P}^N p_k^{\mu_P} = \sum_{i_1, \dots, i_{P-1}=0}^M R_{i_1 \dots i_{P-1}}^{N,K} p_{i_1 \mu_1} \dots p_{i_{P-1} \mu_{P-1}}, \\
R_{\mu_1 \dots \mu_{P-1}}^{0,0} &= T_{\mu_1 \dots \mu_P}^N g^{\mu_{P-1} \mu_P} = \sum_{i_1, \dots, i_{P-2}=0}^M R_{i_1 \dots i_{P-2}}^{N,00} p_{i_1 \mu_1} \dots p_{i_{P-2} \mu_{P-2}},
\end{aligned} \tag{2.96}$$

into eqs. (2.92) and (2.94), these equations could be solved for $T_{i_1 \dots i_P}^N$:

$$\begin{aligned}
T_{00 i_1 \dots i_{P-2}}^N &= \frac{1}{D + P - 2 - M} \left[R_{i_1 \dots i_{P-2}}^{N,00} - \sum_{k=1}^M R_{k i_1 \dots i_{P-2}}^{N,k} \right], \\
T_{k i_1 \dots i_{P-1}}^N &= (X_M^{-1})_{kk'} \left[R_{i_1 \dots i_{P-1}}^{N,k'} - \sum_{r=1}^{P-1} \delta_{i_r}^{k'} T_{00 i_1 \dots i_{r-1} i_{r+1} \dots i_{P-1}}^N \right].
\end{aligned} \tag{2.97}$$

Using the eq. (2.92) and (2.94), the R 's can be expressed in terms of $T_{i_1 \dots i_{P-1}}^N$, and

$T_{i_1 \dots i_q}^{N-1}$, with $q < P$:

$$\begin{aligned}
R_{i_1 \dots i_q}^{N,00} \underbrace{M \dots M}_{P-2-q} &= m_0^2 T_{i_1 \dots i_q}^N \underbrace{M \dots M}_{P-2-q} + (-1)^{P-q} \left[\tilde{T}_{i_1 \dots i_q}^{N-1}(0) + \binom{P-2-q}{1} \sum_{k_1=1}^{M-1} \tilde{T}_{i_1 \dots i_q k_1}^{N-1}(0) \right. \\
&\quad + \binom{P-2-q}{2} \sum_{k_1, k_2=1}^{M-1} \tilde{T}_{i_1 \dots i_q k_1 k_2}^{N-1}(0) + \dots \\
&\quad \left. + \binom{P-2-q}{P-2-q} \sum_{k_1, \dots, k_{P-2-q}=1}^{M-1} \tilde{T}_{i_1 \dots i_q k_1 \dots k_{P-2-q}}^{N-1}(0) \right],
\end{aligned} \tag{2.98}$$

$$\begin{aligned}
R_{i_1 \dots i_q}^{N,k} \underbrace{M \dots M}_{P-1-q} &= \frac{1}{2} \left\{ T_{i_1 \dots i_q}^N \underbrace{M \dots M}_{P-1-q}(k) \theta(k | i_1, \dots, i_q, M, \dots, M) \right. \\
&\quad - f_k T_{i_1 \dots i_q}^N \underbrace{M \dots M}_{P-1-q} - (-1)^{P-1-q} \left[\tilde{T}_{i_1 \dots i_q}^{N-1}(0) \right. \\
&\quad + \binom{P-1-q}{1} \sum_{k_1=1}^{M-1} \tilde{T}_{i_1 \dots i_q k_1}^{N-1}(0) \\
&\quad + \binom{P-1-q}{2} \sum_{k_1, k_2=1}^{M-1} \tilde{T}_{i_1 \dots i_q k_1 k_2}^{N-1}(0) + \dots \\
&\quad \left. \left. + \binom{P-1-q}{P-1-q} \sum_{k_1, \dots, k_{P-1-q}=1}^{M-1} \tilde{T}_{i_1 \dots i_q k_1 \dots k_{P-1-q}}^{N-1}(0) \right] \right\}
\end{aligned} \tag{2.99}$$

where $i_1, \dots, i_q \neq M$ and

$$\theta(k|i_1, \dots, i_{P-1}) = \begin{cases} 1 & i_r \neq k, \quad r = 1, \dots, P-1, \\ 0 & \text{else.} \end{cases} \quad (2.100)$$

The indices \tilde{i} denotes the i -th momentum of the corresponding N -point function T^N but to the $(i-1)$ -th momentum of the $(N-1)$ -point function $T^{N-1}(K)$ if $i > k$. In sum, with the reduction algorithm above all one-loop integrals can be reduced to the scalar ones as long as the matrices X_M are nonsingular.

Next we take the reduction of tensor two-point integrals as an example to illustrate the reduction algorithm describe above.

We start with

$$B_0(p_{10}, m_0, m_1) = \frac{(2\pi\mu)^{4-n}}{i\pi^2} \int d^d q \frac{1}{D_0 D_1} = p_{10\mu} B_1(P_{10}^2, m_0^2, m_1^2). \quad (2.101)$$

Using the relation $q^2 = D_0 - m_0^2$ with (for convenience, $p_{10} \rightarrow p$)

$$qp = \frac{1}{2}(D_1 - D_0 - f), \quad f = p^2 - m_1^2 + m_2^2, \quad (2.102)$$

we derive the following relations:

$$p^2 B_1(p^2, m_0^2, m_1^2) = \frac{1}{2}[A_0(m_0) - A_0(m_1) - f B_0(p, m_0^2, m_1^2)]. \quad (2.103)$$

Therefore we have

$$B_1(p^2, m_1, m_0) = \frac{1}{2p^2}[A_0(m_0) - A_0(m_1) - (m_0^2 - m_1^2 - p^2)B_0(p, m_0^2, m_1^2)]. \quad (2.104)$$

The rank two tensor integral can be reduced as follows:

$$B_{\mu\nu}(p^2, m_0^2, m_1^2) = g_{\mu\nu} B_{00}(p^2, m_0^2, m_1^2) + p_\mu p_\nu B_{11}(p^2, m_0^2, m_1^2) \quad (2.105)$$

Multiplying eq.(2.105) by $g_{\mu\nu}$ and p_ν yields

$$\begin{cases} p^2 B_{11}(p^2, m_0^2, m_0^2) + dB_{00}(p^2, m_0, m_1) = A_0(m_1) - m_0^2 B_0(p^2, m_0^2, m_1^2) \\ p^2 B_{11}(p^2, m_0^2, m_0^2) + B_{00}(p^2, m_0, m_1) = \frac{1}{2}[A_0(m_1) - f B_0(p^2, m_0^2, m_1^2)] \end{cases} \quad (2.106)$$

After a simple calculation, we have

$$\begin{aligned} B_0(p^2, m_0^2, m_1^2) &= \frac{1}{\Delta} - \int_0^1 dx \log\left(\frac{\chi}{\mu^2}\right) \rightarrow \frac{1}{\Delta}, \\ B_1(p^2, m_0^2, m_1^2) &= -\frac{1}{2} \frac{1}{\Delta} - \int_0^1 x dx \log\left(\frac{\chi}{\mu^2}\right) \rightarrow -\frac{1}{2} \frac{1}{\Delta}, \\ B_{11}(p^2, m_0^2, m_1^2) &= \frac{1}{3} \frac{1}{\Delta} - \int_0^1 dx x^2 \log\left(\frac{\chi}{\mu^2}\right) \rightarrow \frac{1}{3} \frac{1}{\Delta}, \\ B_{22}(p^2, m_0^2, m_1^2) &= -\frac{1}{2} \left(\frac{1}{\Delta} + 1\right) \int_0^1 dx \chi + \frac{1}{2} \int_0^1 dx \chi \log\left(\frac{\chi}{\mu^2}\right) \\ &\rightarrow -\frac{1}{4} \left(\frac{1}{3} p^2 - m_0^2 - m_1^2\right) \frac{1}{\Delta}, \end{aligned} \quad (2.107)$$

where

$$\begin{aligned} \chi(x) &= -p^2 x^2 + (p^2 - m_1^2 + m_0^2)x - m_0^2, \\ \Delta &= \frac{2}{4-d} - \gamma_E + \log(4\pi). \end{aligned} \quad (2.108)$$

Using these relations, we get

$$dB_{22}(p^2, m_0, m_1) = 4B_{22}(p^2, m_0^2, m_1^2) + \frac{K^2}{6}, \quad K^2 = p^2 - 3(m_0^2 + m_1^2). \quad (2.109)$$

Furthermore we have

$$\begin{cases} p^2 B_{11}(p^2, m_0^2, m_0^2) + 4B_{00}(p^2, m_0, m_1) = A_0(m_1) - m_0^2 B_0(p^2, m_0^2, m_1^2) - \frac{K^2}{6} \\ p^2 B_{11}(p^2, m_0^2, m_0^2) + B_{00}(p^2, m_0, m_1) = \frac{1}{2}[A_0(m_1) - f B_0(p^2, m_0^2, m_1^2)]. \end{cases} \quad (2.110)$$

At this moment we introduce a X_1 -matrix (nonsingular)

$$\begin{pmatrix} p^2 & 4 \\ p^2 & 1 \end{pmatrix} \quad (2.111)$$

and the vector b

$$b = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = \begin{pmatrix} A_0(m_1^2) - m_0^2 B_0(p^2, m_0^2, m_1^2) - \frac{\kappa^2}{6} \\ \frac{1}{2} [A_0(m_1) + f B_1(p^2, m_0, m_1)] \end{pmatrix} \quad (2.112)$$

Therefore, $B_{00}(p^2, m_0^2, m_1^2)$ and $B_{11}(p^2, m_0^2, m_1^2)$ can be obtained by using the inverse of matrix X_1

$$\begin{pmatrix} B_{11}(p^2, m_0^2, m_1^2) \\ B_{00}(p^2, m_0^2, m_1^2) \end{pmatrix} = X_1^{-1} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}. \quad (2.113)$$

Now the tensor two-point integrals have been reduced to the scalar integrals A_0 and B_0 and their explicit expressions are listed as follows:

$$\begin{aligned} B_1(p^2, m_0^2, m_1^2) &= \frac{1}{2p^2} [A_0(m_0) - A_0(m_2) + (m_1^2 - m_0^2 - p^2) B_0(p^2, m_0^2, m_1^2)], \\ B_{11}(p^2, m_0^2, m_1^2) &= \frac{p^2 - 3(m_0^2 + m_1^2)}{18p^2} \\ &\quad + \frac{\Delta m^2 - p^2}{3p^4} A_0(m_0) - \frac{\Delta m^2 - 2p^2}{3p^4} A_0(m_1) \\ &\quad + \frac{\kappa(-p^2, -m_0^2, -m_1^2) + 3p^2 m_0^2}{3p^4} B_0(p^2, m_0, m_1), \end{aligned}$$

$$\begin{aligned} B_{22}(p^2, m_0^2, m_1^2) &= -\frac{p^2 - 3(m_0^2 + m_1^2)}{18} - \frac{\Delta m^2 - p^2}{12p^2} A_0(m_0) + \frac{\Delta m^2 + p^2}{12p^2} A_0(m_2) \\ &\quad - \frac{\kappa(-p^2, -m_0^2, -m_1^2)}{12p^2} B_0(p^2, m_0^2, m_1^2), \end{aligned} \quad (2.114)$$

with $\Delta m^2 = m_1^2 - m_0^2$.

2.4 Example for One-loop Radiative Correction Calculations

The discussion above gave us a reduction method to compute one-loop tensor integrals, which is a powerful tool for perturbative calculations for the Standard Model.

As illustrations of the reduction method we will present the detailed calculation of the one-loop amplitude for the decay of the W -boson into massless fermions.

$$W^+(k) \rightarrow f_i(p_i)\bar{f}_j(p_j). \quad (2.115)$$




Figure 2.1. Born Diagram to $W \rightarrow f_i\bar{f}_j$

At the tree level only one Feynman diagram contributes to the amplitude

$$\mathcal{M}_0 = -\frac{eV_{ij}}{\sqrt{2}s_w}\bar{u}(p_1)\epsilon(k)\frac{1}{2}(1-\gamma_5)v(p_2) = \frac{eV_{ij}}{\sqrt{2}s_w}\mathcal{M}_1^-, \quad (2.116)$$

with $\mathcal{M}_1^- = \bar{u}(p_1)\epsilon(k)\frac{1}{2}(1-\gamma_5)v(p_2)$, which leads to the lowest order decay width

$$\Gamma_0 = \frac{\alpha}{6} \frac{M_W}{2s_W^2} |V_{ij}|^2. \quad (2.117)$$

There are six loop digrams and one counterterm diagram at one-loop order (see Feynman diagrams below)




Figure 2.2. One-loop diagrams to $W \rightarrow f_i \bar{f}_j$

These six diagrams could be grouped into two generics, the first two loop diagrams corresponding one generic diagram and the rest four corresponding to another generic diagram. So we first compute the two generic diagrams. The amplitude for the first one is written as follows

$$\begin{aligned}
\delta\mathcal{M}_1 &= i\mu^{4-D} \int \frac{d^d q}{2\pi^d} \frac{1}{(q^2 - M^2)(q + p_1)^2(q - p_2)^2} \\
&\bar{u}(p_1)\gamma^\nu(g_1^- \omega_- + g_1^+ \omega_+)(\not{q} + \not{p}_1)\not{\epsilon}(g_3^- \omega_- + g_3^+ \omega_+) \\
&(\not{q} - \not{p}_2)\gamma_\mu(g_2^- \omega_- + g_2^+ \omega_+)v(p_2),
\end{aligned} \tag{2.118}$$

where g^\pm represent the generic left- and right-handed fermion-fermion-vector couplings

$$g_f^+ = -\frac{s_w}{c_w} Q_f, g_f^- = \frac{I^3 - Q_f^2 s_w^2}{s_w c_w}, \quad (2.119)$$

and

$$\omega_\pm = \frac{1}{2}(1 \pm \gamma_5). \quad (2.120)$$

Algebraic simplification and decomposition into tensor integral gives

$$\begin{aligned} \delta\mathcal{M}_1 &= i\mu^{4-D} \int \frac{d^d q}{2\pi^d} \frac{1}{(q^2 - M^2)(q + p_1)^2(q - p_2)^2} \\ &\bar{u}(p_1)[-2(\not{q} - \not{p}_2)\not{\epsilon}(\not{q} + \not{p}_1) + (4 - d)\not{q}\not{\epsilon}\not{q}](g_1^- g_3^- g_2^- \omega_- g_1^+ g_2^+ g_3^+ \omega_+)v(p_2) \\ &= -\frac{1}{16\pi^2} \bar{u}(p_1)[(2 - d)C_{\mu\nu}\gamma^\mu\not{\epsilon}\gamma^\nu + 2C_\mu(\not{p}_2\not{\epsilon}\gamma^\mu - \gamma^\mu\not{\epsilon}\not{p}_1) + 2C_0\not{p}_2\not{\epsilon}\not{p}_1] \\ &(g_1^- g_2^- g_3^- \omega_- + g_1^+ g_2^+ g_3^+ \omega_+)v(p_2). \end{aligned} \quad (2.121)$$

Inserting the Lorentz decomposition (2.87) yields

$$\begin{aligned} \delta\mathcal{M}_1 &= -\frac{1}{16\pi^2}(g_1^- g_3^- g_2^- \mathcal{M}_1^- + g_1^+ g_3^+ g_2^+ \mathcal{M}_1^+) \\ &[(2 - d)^2 C_{00} - 2k^2(C_{12} + C_1 + C_2 + C_0)]. \end{aligned} \quad (2.122)$$

Finally the amplitude for the first generic diagram arrives at

$$\begin{aligned} \delta\mathcal{M}_1 &= -\frac{1}{16\pi^2} \left(g_1^- g_3^- g_2^- \mathcal{M}_1^- + g_1^+ g_3^+ g_2^+ \mathcal{M}_1^+ \right) \\ &\left[-2k^2 C_0(0, k^2, 0, M, 0, 0) \left(1 + \frac{M^2}{k^2} \right)^2 \right. \\ &\quad \left. - B_0(k^2, 0, 0) \left(3 + 2\frac{M^2}{k^2} \right) \right. \\ &\quad \left. + 2B_0(0, M, 0) \left(2 + \frac{M^2}{k^2} \right) - 2 \right]. \end{aligned} \quad (2.123)$$

Similarly the amplitude for the second generic diagram is obtained

$$\begin{aligned}
\delta\mathcal{M}_2 &= -i\mu^{4-d} \int \frac{d^d q}{(2\pi)^d} \frac{\bar{u}(p_1)\gamma^\nu(g_1^-\omega_- + g_1^+\omega_+)(-\not{q})\gamma_\rho(g_2^-\omega_- + g_2^+\omega_+)v(p_2)}{q^2[(q+p_1)^2 - M_1^2][(q-p_2)^2 - M_2^2]} \\
&g_3[g_{\rho\mu}(p_1 + 2p_2 - q)_\nu - g_{\mu\nu}(2p_1 + p_2 + q)_\rho + g_{\nu\rho}(2q + p_1 - p_2)_\mu]\epsilon^\mu \\
&= \frac{1}{16\pi^2}g_3(g_1^-g_2^-\mathcal{M}_1^- + g_1^+g_2^+\mathcal{M}_1^+)[4(d-1)C_{00} - 2k^2(C_{12} + C_1 + C_2)] \\
&= \frac{1}{16\pi^2}g_3(g_1^-g_2^-\mathcal{M}_1^- + g_1^+g_2^+\mathcal{M}_1^+) \\
&\left[2\left(M_1^2 + M_2^2 + \frac{M_1^2M_2^2}{k^2}\right)C_0(0, k^2, 0, 0, M_1, M_2) - \left(1 + \frac{M_1^2 + M_2^2}{k^2}\right) \right. \\
&\left. B_0(K^2, M_1, M_2) + \left(2 + \frac{M_1^2}{k^2}\right)B_0(0, 0, M_1) + \left(2 + \frac{M_2^2}{k^2}\right)B_0(0, 0, M_2) \right]
\end{aligned} \tag{2.124}$$

For convenience we define the generic vertex function as follows:

$$\begin{aligned}
\mathcal{V}_a(m_1^2, m_0^2, m_2^2, M_0, M_1, M_2) &= B_0(m_0^2, M_1, M_2) - 2 \\
&\quad - (M_0^2 - m_1^2 - M_1^2)C_1 - (M_0^2 - m_2^2 - M_2^2)C_2 \\
&\quad - 2(m_0^2 - m_1^2 - m_2^2)(C_1 + C_2 + C_0), \\
\mathcal{V}_b^-(m_1^2, m_0^2, m_2^2, M_0, M_1, M_2) &= 3B_0(m_0^2, M_1, M_2) + 4M_0^2C_0 \\
&\quad + (4m_1^2 + 2m_2^2 - 2m_0^2 + M_0^2 - M_1^2)C_1 \\
&\quad + (4m_2^2 + 2m_1^2 - 2m_0^2 + M_0^2 - M_2^2)C_2.
\end{aligned} \tag{2.125}$$

so that amplitudes for two generic diagrams can be expressed as

$$\begin{aligned}
\delta\mathcal{M}_1 &= -\frac{1}{16\pi^2}(g_1^-g_3^-g_2^-\mathcal{M}_1^- + g_1^+g_3^+g_2^+\mathcal{M}_1^+)\mathcal{V}_a(0, k^2, 0, M, 0, 0) \\
\delta\mathcal{M}_2 &= \frac{1}{16\pi^2}g_3(g_1^-g_2^-\mathcal{M}_1^- + g_1^+g_2^+\mathcal{M}_1^+)\mathcal{V}_b^-(0, k^2, 0, 0, M_1, M_2).
\end{aligned} \tag{2.126}$$

Next, we only need to insert the actual couplings and masses of the six actual diagrams into these two generic diagrams and add the counterterm diagram. Finally we obtain the virtual one loop corrections to the invariant amplitude for $W \rightarrow f_i \bar{f}_j$

$$\begin{aligned}
\delta\mathcal{M} = & -\frac{e}{\sqrt{2}s_W} \frac{\alpha}{4\pi} V_{ij} \mathcal{M}_1^- \\
& \left\{ Q_{f_i} Q_{f_j} \mathcal{V}_a(0, M_W^2, 0, \lambda, 0, 0) + g_{f_i}^- g_{f_j}^- \mathcal{V}_a(0, M_W^2, 0, M_Z, 0, 0) \right. \\
& + Q_{f_i} \mathcal{V}_b(0, M_W^2, 0, 0, \lambda, M_W) - Q_{f_j} \mathcal{V}_b(0, M_W^2, 0, 0, M_W, \lambda) \\
& + \frac{c_W}{s_W} g_{f_i}^- \mathcal{V}_b(0, M_W^2, 0, 0, M_Z, M_W) - \frac{c_W}{s_W} g_{f_j}^- \mathcal{V}_b(0, M_W^2, 0, 0, M_W, M_Z) \\
& \left. + \frac{1}{2} \delta Z_{ii}^{f_i, L\dagger} + \frac{1}{2} \delta Z_{jj}^{f_j, L\dagger} + \frac{1}{2} \delta Z_W + \delta Z_e - \frac{\delta s_W}{s_W} \right\}. \tag{2.127}
\end{aligned}$$

$\delta\mathcal{M}$ contains infrared divergences and they are regulated with a photon mass λ .

CHAPTER THREE

Spinor Techniques

3.1 Introduction

The cross sections of bremsstrahlung processes at high energies in gauge theories are of interest for both theoretical and experimental physicists due to the development of experiments in colliders. In these processes, light leptons and quarks could be viewed as massless as long as the electromagnetic and strong interactions are concerned. When the Feynman rules of the theory are determined, the rest of the work is reduced to the calculation of the amplitudes of Feynman diagrams. The amplitudes can be computed with the standard manipulations for squaring matrix elements and summing over the polarizations of the particles.

Although the standard approach is straightforward, it becomes impractical when both the number of external lines and the number of diagrams involved become large. To be specific, after writing down the amplitude M of the corresponding Feynman diagrams, we usually have an analytic expression for the cross section $\sum |M|^2$ with a spin and/or color sum or average. The result is usually a function of Minkowski products of the particle four-momenta.

The calculation of the cross section is facilitated by two considerations. First, the polarization vector ϵ^μ of external spin-1 particles could be summed covariantly. For a massive vector boson with mass m and momentum q^μ we have the spin sum

$$\sum \epsilon^\mu \epsilon^{*\nu} = -g^{\mu\nu} + \frac{q^\mu q^\nu}{m^2}, \quad (3.1)$$

while for a massless vector boson with momentum k^μ we have in the axial gauge:

$$\sum \epsilon^\mu \epsilon^{*\nu} = -g^{\mu\nu} + \frac{p^\mu k^\nu + k^\mu p^\nu}{p \cdot k}, \quad (3.2)$$

where p^μ is a four-vector different from k^μ . Second, a product of spinor sandwiches can be expressed as a trace over a string of Dirac matrices, using spinor projection operators. For a massive spin- $\frac{1}{2}$ particle with mass m and momentum p^μ we have the spin sum

$$\begin{cases} \sum \bar{u}(p)u(p) = \not{p} + m, \\ \sum \bar{v}(p)v(p) = \not{p} - m. \end{cases} \quad (3.3)$$

For a massless fermion with momentum p^μ and helicity $\lambda = \pm 1$, we have

$$u_\lambda(p)\bar{u}_\lambda(p) = \omega_\lambda \not{p}, \quad \omega_\lambda = \frac{1}{2}(1 + \lambda\gamma_5). \quad (3.4)$$

The above method has some appealing features. The unobserved spins and polarizations do not arise in the final result; the arbitrary overall complex phases of the spinors and polarization vectors cancel. The algebraic calculation of the trace expressions is straightforward and can be performed for any amplitude. However, the last condition limits the complexity of the problems that can be dealt with. Since we have to square the amplitude before we can use eqs. (3.1)-(3.4), both the number of traces and their length increases very fast with the order of perturbation theory. It is very easy to make mistakes.

In order to handle this problem, spinor product methods were proposed by three groups independently: the CALKUL approach [46–50], “Chinese magic” polarization scheme [51], and Kleiss and Stirling Spinor Technique [52]. The essential idea is that the gauge invariance allows one to use a set of polarization vectors which

eliminate radiation from one entire side of a charged line and simplify considerably the calculation. In this chapter, we will introduce these three approaches respectively.

3.2 The CALKUL Approach

First, we introduce the helicity amplitude approach [46–49] proposed the CALKUL collaboration (F. A. Berends, P. De Causmaecker, R. Gastmans, R. Kleiss, W. Troost and T. T. Wu).

For a massless fermion of four-momentum q , there are two possible helicity states $u_+(q)$ and $u_-(q)$ specified by

$$u_{\pm}(q) = \frac{1}{2}(1 \pm \gamma_5)u_{\pm}(q), \quad (3.5)$$

$$\bar{u}_{\pm}(q) = \bar{u}_{\pm}(q)\frac{1}{2}(1 \mp \gamma_5). \quad (3.6)$$

By the normalization

$$u_+(q)\bar{u}_+(q) + u_-(q)\bar{u}_-(q) = \not{q}, \quad (3.7)$$

it follows that

$$u_{\pm}(q)\bar{u}_{\pm}(q) = \frac{1}{2}(1 \pm \gamma_5)\not{q}. \quad (3.8)$$

For the anti-fermion of momentum q , the relations are similar:

$$v_{\pm}(q) = \frac{1}{2}(1 \mp \gamma_5)v_{\pm}(q), \quad (3.9)$$

$$\bar{v}_{\pm}(q) = \bar{v}_{\pm}(q)\frac{1}{2}(1 \pm \gamma_5), \quad (3.10)$$

$$v_{\pm}(q)\bar{v}_{\pm}(q) = \frac{1}{2}(1 \mp \gamma_5)\not{q}. \quad (3.11)$$

It is convenient to apply the Dirac bracket notation.

$$\begin{aligned}
|q, +\rangle &\equiv u_+(q), & |q, +\rangle &\equiv v_-(q), \\
|q, -\rangle &\equiv u_-(q), & |q, -\rangle &\equiv v_+(q), \\
\langle q, +| &\equiv \bar{u}_+(q), & \langle q, -| &\equiv \bar{v}_-(q), \\
\langle q, -| &\equiv \bar{u}_-(q), & \langle q, -| &\equiv \bar{v}_+(q).
\end{aligned} \tag{3.12}$$

Then we have

$$\begin{aligned}
|q, +\rangle \langle q, +| &= \frac{1}{5}(1 + \gamma_5)\not{q}, \\
|q, -\rangle \langle q, -| &= \frac{1}{5}(1 - \gamma_5)\not{q},
\end{aligned} \tag{3.13}$$

and

$$\langle q, +|q, +\rangle = \langle q, -|q, -\rangle = 0. \tag{3.14}$$

More generally, for arbitrary q_1 and q_2 satisfying $q_1^2 = q_2^2 = 0$,

$$\langle q_1, \pm|q_2, \pm\rangle = 0. \tag{3.15}$$

Products like $\langle q_1, +|q_2, -\rangle$ are not well defined since each state can carry an arbitrary phase, but the norm is

$$|\langle q_1, +|q_2, -\rangle|^2 = \text{tr} \left[\frac{1}{2}(1 - \gamma_5)\not{q}_2 \frac{1}{2}(1 + \gamma_5)\not{q}_1 \right] = 2(q_1 q_2). \tag{3.16}$$

Next, let us discuss the polarization vectors of the gauge boson. Assume a massless gauge boson with four-momentum k is radiated from a charged line for which q_+ and q_- are the momenta of the outgoing antifermion and fermion. A massless boson has two polarization states, and the polarizations can be constructed as follows:

$$\left\{ \begin{array}{l} \epsilon_\mu^\parallel = N[(q_+ k)q_{-\mu} - (q_- k)q_{+\mu}], \\ \epsilon_\mu^\perp = N\epsilon_{\mu\alpha\beta\gamma}q_+^\alpha q_-^\beta k^\gamma, \end{array} \right. \tag{3.17}$$

where

$$N = [2(q_+q_-)(q_+k)(q_-k)]^{\frac{1}{2}}. \quad (3.18)$$

Alternatively, from eq. (3.17), we can introduce the circular polarization vectors

$$\epsilon_\mu^\pm = \sqrt{\frac{1}{2}}(\epsilon_\mu^\parallel \pm i\epsilon_\mu^\perp), \quad (3.19)$$

with which we will work from now on.

Using the identity

$$i\gamma^\mu \epsilon_{\mu\alpha\beta\gamma} = (\gamma_\alpha\gamma_\beta\gamma_\gamma - \gamma_\alpha g_{\beta\gamma} + \gamma_\beta g_{\alpha\gamma} - \gamma_\gamma g_{\alpha\beta})\gamma_5, \quad (3.20)$$

we have

$$\not{\epsilon}^\pm = -\frac{1}{2\sqrt{2}}N[\not{k}\not{q}_-k\not{q}_+(1 \pm \gamma_5) - \not{q}_-\not{q}_+\not{k}(1 \mp \gamma_5)]. \quad (3.21)$$

which leads to great simplifications. There are several reasons for advantages resulting from eq. (3.21):

- If the gauge boson line is next to the the external fermion or antifermion line, only one of the terms on the right-hand side of eq. (3.21) gives a non-zero contribution due to the Dirac equations for massless fermions.
- When there is a gauge boson line next to the external fermion or antifermion line, either a factor $1 + \gamma_5$ or a factor $1 - \gamma_5$ occurs. This factor ensures for every other real boson line attached to this fermion line, only one of the two terms on the right-hand side of eq. (3.21) survives.
- When the gauge boson line is next to the external fermion line, there is a cancellation of the denominator. Assume we have an outgoing electron with momentum q_- , and the emission of a photon with momentum k . If the vertex for the photon

emission is next to the outgoing electron line, then the amplitude contains a factor

$$\begin{aligned}\bar{u}(q_-)\not{\epsilon}^\pm\frac{\not{q}_- + \not{k}}{2(q_-k)} &= -\frac{1}{2\sqrt{2}}N\bar{u}(q_-)\not{k}\not{q}_-\not{q}_+(1 \pm \gamma_5)\frac{q_- + k}{q_-k} \\ &= -\frac{1}{2\sqrt{2}}N\bar{u}(q_-)\not{q}_+(\not{q}_- + \not{k})(1 \mp \gamma_5).\end{aligned}\quad (3.22)$$


Therefore the denominator $2(q_-k)$ is cancelled. In the upcoming example, we will see this denominator cancellation makes a large contribution for the simplicity of the calculation.

In order to illustrate the nice features of the CALKUL helicity amplitude method, we will exhibit the explicit computation the amplitude for the single bremsstrahlung process in QED.

For the reaction

$$e^+(p_+) + e^-(p_-) \rightarrow \mu^+(q_+) + \mu^-(q_-) + \gamma(k), \quad (3.23)$$

for which the Feynman diagrams are shown below



Applying the Feynman rules, we have

$$\begin{aligned}
M_1 &= \frac{ie^2}{2(q_+q_-)} \bar{v}(p_+) \gamma_\mu \frac{\not{p}_- - \not{k}}{2(p_-k)} \not{\epsilon} u(p_-) \bar{u}(q_-) \gamma^\mu v(q_+), \\
M_2 &= \frac{ie^2}{2(q_+q_-)} \bar{v}(p_+) \gamma_\mu \frac{-\not{p}_+ + \not{k}}{-2(p_+k)} \not{\epsilon} u(p_-) \bar{u}(q_-) \gamma^\mu v(q_+), \\
M_3 &= \frac{ie^2}{2(p_+p_-)} \bar{v}(p_+) \gamma_\mu u(p_-) \bar{u}(q_-) \not{\epsilon} \frac{\not{q}_- + \not{k}}{2(q_-k)} \gamma^\mu v(q_+), \\
M_4 &= \frac{ie^2}{2(p_+p_-)} \bar{v}(p_+) \gamma_\mu u(p_-) \bar{u}(q_-) \gamma^\mu \frac{-\not{q}_+ - \not{k}}{2(q_+k)} \not{\epsilon} v(q_+). \tag{3.24}
\end{aligned}$$

When the photon is radiated from the electron line, it is convenient to choose

$$\begin{aligned}
\epsilon_p^\pm &= N_p [\not{p}_+ \not{p}_- \not{k} (1 \mp \gamma_5) - \not{k} \not{p}_+ \not{p}_- (1 \pm \gamma_5)], \\
N_p^{-1} &= 4[(p_+p_-)(p_+k)(p_-k)]^{\frac{1}{2}}. \tag{3.25}
\end{aligned}$$

but for radiation from the muon line, it is advantageous to take

$$\begin{aligned}
\epsilon_q^\pm &= N_q [\not{q}_- \not{q}_+ \not{k} (1 \mp \gamma_5) - \not{k} \not{q}_- \not{q}_+ (1 \pm \gamma_5)], \\
N_q^{-1} &= 4[(q_+q_-)(q_+k)(q_-k)]^{\frac{1}{2}}. \tag{3.26}
\end{aligned}$$

The two choices are related by a simple phase factor. Let us consider the photon to be moving along the z -axis. Because the diagrams M_1 and M_2 together form a gauge-invariant set, and so do M_3 and M_4 , we can make gauge transformations so that the polarization vector ϵ_p^\pm and ϵ_q^\pm only have components in the xy -plane. Since they have the same norm, they can differ at most by a phase factor and terms proportional k ,

$$\begin{aligned}
e_q &= e^{\pm i\phi} \epsilon_p^\pm + \beta_\pm k, \\
e^{\pm i\phi} &= -(\epsilon_p^\mp \epsilon_q^\pm) = -N_p N_q \text{tr}[\not{p}_+ \not{p}_- \not{k} \not{q}_- \not{q}_+ \not{k} (1 \mp \gamma_5)]. \tag{3.27}
\end{aligned}$$

Now we are ready to calculate the helicity amplitudes which are denoted by $M(\lambda_1(e^+), \lambda_2(e^-), \lambda_3(\mu^+), \lambda_4(\mu^-), \lambda_5(k))$. Due to helicity conservation, electron and

positron helicities must be opposite, as well as the muon helicities. For for all λ_5 ,

$$\begin{aligned} M(+, +, +, +, \lambda_5) &= M(-, -, +, +, \lambda_5) \\ &= M(+, +, -, -, \lambda_5) = M(-, -, -, -, \lambda_5) = 0. \end{aligned} \quad (3.28)$$

Consider the non-vanishing helicity amplitudes next. Let us introduce the notation

$$\begin{aligned} s &= (p_+ + p_-)^2, \quad s' = (q_+ + q_-)^2, \\ t &= (p_+ - q_+)^2, \quad t' = (p_- - q_-)^2, \\ u &= (p_+ - q_-)^2, \quad u' = (p_- - q_+)^2, \end{aligned} \quad (3.29)$$

with

$$s + s' + t + t' + u + u' = 0. \quad (3.30)$$

For the amplitude $M(+, -, +, -, +)$, only the diagrams M_1 and M_4 contribute.

$$\begin{aligned} &M(+, -, +, -, +) \\ &= \frac{ie^3}{s'} N_p e^{i\phi} \bar{v}(p_+) \gamma_\mu \frac{\not{p}_- - \not{k}}{-2(p_- k)} \not{p}_+ \not{p}_- \not{k} (1 - \gamma_5) u(p_-) \bar{u}(q_-) \gamma^\mu \frac{1 - \gamma_5}{2} v(q_+) \\ &+ \frac{ie^3}{s} N_p \bar{v}(p_+) \gamma_\mu \frac{1 - \gamma_5}{2} u(p_-) \bar{u}(q_-) \gamma^\mu \frac{-\not{q}_+ - \not{k}}{2(q_+ k)} \not{q}_- \not{q}_+ \not{k} (1 - \gamma_5) v(q_+) \\ &= -\frac{ie^3}{2s'} N_p e^{i\phi} \bar{v}(p_+) \gamma_\mu (\not{q}_+ + \not{q}_-) \not{p}_+ (1 - \gamma_5) u(p_-) \bar{u}(q_-) \gamma^\mu (1 - \gamma_5) v(q_+) \\ &- \frac{ie^3}{2s'} N_q \bar{v}(p_+) \gamma_\mu (1 - \gamma_5) u(p_-) \bar{u}(q_-) \gamma^\mu (\not{p}_+ + \not{p}_-) \not{q}_- (1 - \gamma_5) v(q_+). \end{aligned} \quad (3.31)$$

To simplify this result further, we rewrite the first term for example

$$\begin{aligned}
& -\frac{ie^3}{2s}N_p e^{i\phi}\bar{v}(p_+)\gamma_\mu(\not{q}_+ + \not{q}_-)\not{p}_+(1-\gamma_5)u(p_-)\bar{u}(q_-)\gamma^\mu(1-\gamma_5)v(q_+) \\
& = -\frac{ie^3}{2s}N_p e^{i\phi}\frac{\bar{v}(p_+)\gamma_\mu(\not{q}_+ + \not{q}_-)\not{p}_+(1-\gamma_5)u(p_-)\bar{u}(q_-)\gamma^\mu(1-\gamma_5)v(q_+)}{\bar{v}(q_+)\not{q}_-(1-\gamma_5)v(p_+)} \\
& \times \bar{v}(q_+)\not{q}_-(1-\gamma_5)v(p_+) \\
& = \frac{4ie^3}{s}N_p e^{i\phi}\frac{\bar{u}(q_-)\not{p}_+\not{q}_-\not{q}_+(\not{q}_+ + \not{q}_-)\not{p}_+(1-\gamma_5)u(p_-)}{\bar{v}(q_+)\not{q}_-(1-\gamma_5)v(p_+)} \\
& = \frac{16ie^3}{s}N_p(p_+q_-)(q_+q_-)\frac{\bar{u}(q_-)\not{p}_+(1-\gamma_5)u(p_-)\bar{v}(p_+)\not{q}_-(1-\gamma_5)v(q_+)}{\text{tr}[\not{q}_+\not{q}_-(1-\gamma_5)\not{p}_+\not{q}_-(1-\gamma_5)]} \\
& = \frac{ie^3}{s}N_p\bar{u}(q_-)\not{p}_+(1-\gamma_5)u(p_-)\bar{v}(p_+)\not{q}_-(1-\gamma_5)v(q_+). \tag{3.32}
\end{aligned}$$

Performing similar manipulations with the last term in eq. (3.31) we have

$$\begin{aligned}
& -\frac{ie^3}{2s'}N_q\bar{v}(p_+)\gamma_\mu(1-\gamma_5)u(p_-)\bar{u}(q_-)\gamma^\mu(\not{p}_+ + \not{p}_-)\not{q}_-(1-\gamma_5)v(q_+) \\
& = \frac{ie^3}{s}N_q\bar{u}(q_-)\not{p}_+(1-\gamma_5)u(p_-)\bar{v}(p_+)\not{q}_-(1-\gamma_5)v(q_+). \tag{3.33}
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
& M(+, -, +, -, +) \\
& = -ie^3\left[\frac{N_q}{s}(\epsilon_q^-\epsilon_q^+) + \frac{N_p}{s'}(\epsilon_p^-\epsilon_q^+)\right]\bar{u}(q_-)\not{p}_+(1-\gamma_5)u(p_-)\bar{v}(p_-)\not{q}_-(1-\gamma_5)v(q_+) \\
& \simeq 4e^3\frac{u}{(ss')^{\frac{1}{2}}}[s'N_q(\epsilon_q^-\epsilon_q^+) + sN_p(\epsilon_p^-\epsilon_q^+)]. \tag{3.34}
\end{aligned}$$

For the helicity amplitude $M(+, -, -, +, +)$, only the diagrams M_1 and M_3 survive and we have, performing similar manipulations

$$\begin{aligned}
& M(+, -, -, +, +) \\
& = ie^3\left[\frac{N_q}{s}(\epsilon_q^-\epsilon_q^+) + \frac{N_p}{s'}(\epsilon_p^-\epsilon_q^+)\right]\bar{u}(q_-)\not{q}_+\not{p}_+(1-\gamma_5)u(p_-)\bar{v}(p_+)(1+\gamma_5)v(q_+) \\
& \simeq 4e^3[s'N_q(\epsilon_q^-\epsilon_q^+) + sN_p(\epsilon_p^-\epsilon_q^+)]\frac{t}{(ss')^{\frac{1}{2}}}. \tag{3.35}
\end{aligned}$$

In a completely analogous way, we have

$$\begin{aligned}
M(-, +, +, -, +) &\simeq 4e^3 [s' N_q(\epsilon_q^- \epsilon_q^+) + s N_p(\epsilon_p^- \epsilon_q^+)] \frac{t'}{(ss')^{\frac{1}{2}}}, \\
M(-, +, -, +, +) &\simeq 4e^3 [s' N_q(\epsilon_q^- \epsilon_q^+) + s N_p(\epsilon_p^- \epsilon_q^+)] \frac{u'}{(ss')^{\frac{1}{2}}}.
\end{aligned} \tag{3.36}$$

By parity conjugation, we have

$$\begin{aligned}
M(+, -, +, -, -) &\simeq 4e^3 [s' N_q(\epsilon_q^- \epsilon_q^+) + s N_p(\epsilon_p^- \epsilon_q^+)] \frac{u'}{(ss')^{\frac{1}{2}}}, \\
M(+, -, -, +, -) &\simeq 4e^3 [s' N_q(\epsilon_q^- \epsilon_q^+) + s N_p(\epsilon_p^- \epsilon_q^+)] \frac{t'}{(ss')^{\frac{1}{2}}}, \\
M(-, +, +, -, -) &\simeq 4e^3 [s' N_q(\epsilon_q^- \epsilon_q^+) + s N_p(\epsilon_p^- \epsilon_q^+)] \frac{t}{(ss')^{\frac{1}{2}}}, \\
M(-, +, -, +, -) &\simeq 4e^3 [s' N_q(\epsilon_q^- \epsilon_q^+) + s N_p(\epsilon_p^- \epsilon_q^+)] \frac{u}{(ss')^{\frac{1}{2}}}.
\end{aligned} \tag{3.37}$$

From eqs. (3.34), (3.36) and (3.37), we obtain the unpolarized squared matrix element

$$|M|^2 = 8e^6 |s' N_q(\epsilon_q^+ \epsilon_q^-) + s N_p(\epsilon_p^+ \epsilon_q^-)|^2 \frac{t^2 + t'^2 + u^2 + u'^2}{ss'}. \tag{3.38}$$

Introducing the vectors

$$v_q = \frac{q_+}{(q_+ k)} - \frac{q_-}{(q_- k)}, \quad v_p = \frac{p_+}{(p_+ k)} - \frac{p_-}{(p_- k)}, \tag{3.39}$$

we have

$$\begin{aligned}
|s' N_q(\epsilon_q^+ \epsilon_q^-)|^2 &= -\frac{1}{8} v_q^2, \\
|s' N_p(\epsilon_p^+ \epsilon_q^-)|^2 &= -\frac{1}{8} v_p^2, \\
\Re[s s' N_q N_p(\epsilon_q^+ \epsilon_q^-)(\epsilon_q^+ \epsilon_p^-)] &= \frac{1}{8} v_q v_p.
\end{aligned} \tag{3.40}$$

Furthermore we obtain the spin averaged matrix element

$$|M|^2 = -e^6 (v_q - v_p)^2 \frac{t^2 + t'^2 + u^2 + u'^2}{ss'}. \tag{3.41}$$

As it is shown, through the introduction of explicit polarization vectors for the radiated gauge boson, it is feasible to compute the various helicity amplitudes for single bremsstrahlung in massless QED in a simple and covariant way. For each amplitude, only a few diagrams contribute, rendering the calculation very easy. Although we discussed only the process $e^+e^- \rightarrow \mu^+\mu^-\gamma$, the outline technique could be applied to all bremsstrahlung process. The introduction of the polarization vectors for the radiated photons bring in relatively simple expressions for the various helicity amplitudes in all cases in which the fermions are massless. And the introduction of analogous helicity vectors for the external gauge bosons result in similar simplifications in $SU(N)$ gauge theories.

3.3 The "Chinese Magic" Polarization Scheme

The massless spinors with momentum p and helicity λ , $u_{\pm}(p)$, $v_{\pm}(p)$, $\bar{u}_{\pm}(p)$, $\bar{v}_{\pm}(p)$, satisfy the relations

$$\not{p}u_{\pm}(p) = \not{p}v_{\pm}(p) = \not{p}\bar{u}_{\pm}(p) = \not{p}\bar{v}_{\pm}(p) = 0, \quad p^2 = 0, \quad (3.42)$$

$$(1 \mp \gamma_5)u_{\pm} = (1 \pm \gamma_5)v_{\pm} = \bar{u}_{\pm}(1 \pm \gamma_5) = \bar{v}_{\pm}(1 \mp \gamma_5) = 0, \quad (3.43)$$

with the normalization

$$\bar{u}_{\pm}(p)\gamma_{\mu}u_{\pm}(p) = \bar{v}_{\pm}(p)\gamma_{\mu}v_{\pm}(p) = 2p_{\mu}. \quad (3.44)$$

We use the Dirac convention

$$\begin{aligned} u_{\pm}(p) &= v_{\mp} = |p_{\pm}\rangle \\ \bar{u}_{\pm}(p) &= \bar{v}_{\mp} = \langle p_{\pm}|, \end{aligned} \quad (3.45)$$

and

$$|p_{-}\rangle = |p_{+}\rangle^c, \quad (3.46)$$

where $|\psi\rangle^c$ denotes the charge conjugation of the spinor $|\psi\rangle$. The following relations hold for massless momenta p and q ,

$$|p_{\pm}\rangle \langle p_{\pm}| = \frac{1}{2}(1 \pm \gamma_5)\not{p}, \quad (3.47)$$

$$\langle p_+|q_+\rangle = \langle p_-|q_-\rangle = 0, \quad (3.48)$$

$$\langle p_-|q_+\rangle = -\langle q_-|p_+\rangle, \quad (3.49)$$

$$\langle p_-|p_+\rangle = \langle p_+|p_-\rangle = 0. \quad (3.50)$$

For simplicity we let

$$\langle p_-|q_+\rangle = \langle pq\rangle; \quad (3.51)$$

then

$$\langle q_-|p_+\rangle = -\langle pq\rangle,$$

$$\langle q_+|p_-\rangle = \langle pq\rangle^*,$$

$$\langle p_+|q_-\rangle = -\langle pq\rangle^*. \quad (3.52)$$

and

$$|\langle pq\rangle|^2 = 2(pq). \quad (3.53)$$

The scalar $\langle pq\rangle$ is called the spinor inner-product [53] which play a vital role in the "Chinese magic" polarization scheme. Next, we introduce properties of the spinor inner-product for the following discussions.

In general, for any massless spinor $\langle A_{\pm}|$, we have

$$\langle A_{\mp}|B_{\pm}\rangle = -\langle B_{\mp}|A_{\pm}\rangle,$$

$$\langle A_+|\gamma_{\mu}|B_+\rangle = \langle B_-|\gamma_{\mu}|A_-\rangle, \quad (3.54)$$

where $\langle A_{\pm} | = \overline{|A_{\pm}\rangle}$, and if $|A_{\pm}\rangle$ has the form

$$|A_{\pm}\rangle = \begin{cases} k_1 \cdots k_n |q_{\pm}\rangle & (n \text{ even}) \\ k_1 \cdots k_n |q_{\mp}\rangle & (n \text{ odd}), \end{cases} \quad (3.55)$$

then

$$|A_{\pm}\rangle^c = \begin{cases} k_1 \cdots k_n |q_{\mp}\rangle & (n \text{ even}) \\ -k_1 \cdots k_n |q_{\pm}\rangle & (n \text{ odd}), \end{cases} \quad (3.56)$$

where q is a momentum with $q^2 = 0$ and k_i ($i = 1, \dots, n$) is any momentum with or without $k^2 = 0$. Thus we have

$$\begin{aligned} \langle p_- | k_1 \cdots k_n | q_+ \rangle &= -\langle q_- | k_n \cdots k_1 | p_+ \rangle \quad (n \text{ even}), \\ \langle p_+ | k_1 \cdots k_n | q_- \rangle &= -\langle q_+ | k_n \cdots k_1 | p_- \rangle \quad (n \text{ even}), \\ \langle p_+ | k_1 \cdots k_n | q_+ \rangle &= -\langle q_- | k_n \cdots k_1 | p_- \rangle \quad (n \text{ even}). \end{aligned} \quad (3.57)$$

The matrix $|B_+\rangle \langle A_+|$ can be expanded into a linear combination of 1 , γ_{μ} , γ_5 , $\gamma_{\mu}\gamma_5$ and $\gamma_{\mu}\gamma_{\nu}$ ($\mu \neq \nu$):

$$2 |B_+\rangle \langle A_+| = \langle A_+ | \gamma_{\mu} | B_+ \rangle \gamma^{\mu} \frac{1}{2} (1 - \gamma_5). \quad (3.58)$$

Furthermore, we have

$$\begin{aligned} \langle A_+ | \gamma_{\mu} | B_+ \rangle &= \langle B_- | \gamma_{\mu} | A_- \rangle \\ \langle A_+ | \gamma_{\mu} | B_+ \rangle \langle C_- | \gamma^{\mu} | D_- \rangle &= 2 \langle A_+ | D_- \rangle \langle C_- | B_+ \rangle. \end{aligned} \quad (3.59)$$

The circular polarization vectors ϵ^\pm of a gauge boson with momentum k , in which the sign \pm represents the helicity, satisfying the relations below,

$$\begin{aligned}(\epsilon^+ k) &= (\epsilon^- k) = (\epsilon^+)^2 = (\epsilon^-)^2 = 0, \\(\epsilon^+)^* &= \epsilon^-, \\(\epsilon^+ \epsilon^-) &= -1.\end{aligned}\tag{3.60}$$

The polarization vectors are defined by referring to another momentum ($q^2 = 0$) so that [53]

$$\epsilon_\mu^+(k, q) = \frac{\langle q_- | \gamma_\mu | k_- \rangle}{\sqrt{2} \langle qk \rangle},\tag{3.61}$$

$$\epsilon_\mu^-(k, q) = \frac{\langle q_+ | \gamma_\mu | k_+ \rangle}{\sqrt{2} \langle qk \rangle^*},\tag{3.62}$$

and

$$\not{\epsilon}^+(k, q) = \frac{\sqrt{2}}{\langle qk \rangle} [|k_- \rangle \langle q_- | + |q_+ \rangle \langle k_+ |],\tag{3.63}$$

$$\not{\epsilon}^-(k, q) = \frac{\sqrt{2}}{\langle qk \rangle^*} [|k_+ \rangle \langle q_+ | + |q_- \rangle \langle k_- |].\tag{3.64}$$

Furthermore we have

$$(\epsilon^+ q) = (\epsilon^- q) = 0,\tag{3.65}$$

and when the reference momentum q is transformed into p the polarization vectors change only by an additional term proportional to k :

$$\epsilon^\pm(k, q) = \epsilon^\pm(k, p) + \beta^\pm(p, q, k)k,\tag{3.66}$$

where

$$\beta^+(p, q, k) = \frac{\sqrt{2} \langle pq \rangle}{\langle pk \rangle \langle qk \rangle}, \quad \beta^-(p, q, k) = \beta^+(p, q, k)^*.\tag{3.67}$$

Due to the gauge-invariance of the theories the amplitudes vanish when the polarization vector of an external gauge boson is set equal to its momentum,

$$[M]_{\epsilon=k} = 0, \tag{3.68}$$

so the reference momentum could be chosen in an arbitrary way without changing the amplitude.

The polarization vector defined above differs from that of the CALKUL collaboration by a term proportional to k and a phase factor. Thus, all the attractive features in the CALKUL approach [46, 48] are preserved in the "Chinese magic" polarization scheme. Especially some of the helicity amplitudes vanish when an external boson line is attached to an external fermion line. This can be achieved in the "Chinese magic" scheme by choosing the reference momentum in the following way: since the helicity signs appearing in the bra and ket must be the same for the same fermion line, and we let it be the helicity of the line. When the boson has the same (opposite) helicity as that of the fermion line attached, the reference momentum is chosen to be the incoming (outgoing) momentum of the line. Moreover, the polarization vector is now expressed in terms of the spinors, therefore it gives the factorization of the amplitude in a natural way.

As an illustration of the helicity amplitude method in "Chinese magic" scheme we will present the explicit calculation of helicity amplitudes for the single bremsstrahlung process. Consider the process

$$e^+(p') + e^-(p) \rightarrow \mu^+(q') + \mu^-(q) + \gamma(k), \tag{3.69}$$

for which the Feynman diagrams are shown below, and the helicity amplitude is written as $M(\lambda(e^+), \lambda(e^-), \lambda(\mu^+), \lambda(\mu^-), \lambda(\gamma))$. For the helicity amplitude $M(-, +, +, -, +)$,


the polarization vector of the photon should be chosen to be

$$\not{\epsilon}_q^+ = \not{\epsilon}^+(k, q) = \frac{\sqrt{2}}{\langle qk \rangle} [|k_- \rangle \langle q_-| + |q_+ \rangle \langle k_+|] \quad (3.70)$$

for the diagrams (1) and (2), and

$$\not{\epsilon}_p^+ = \not{\epsilon}^+(k, p) = \frac{\sqrt{2}}{\langle pk \rangle} [|k_- \rangle \langle p_-| + |p_+ \rangle \langle k_+|] \quad (3.71)$$

for the diagrams (3) and (4).



Note that [49, 53] the diagrams (1) + (2) and (3) + (4) form two independent gauge invariant subsets of the Feynman diagrams, distinguished by the photon attachment to different fermion lines, namely,

$$[M_{(1)} + M_{(2)}]_{\epsilon=k} = [M_{(3)} + M_{(4)}]_{\epsilon=k} = 0, \quad (3.72)$$

Because of eqs. (3.29) and (3.35) we have

$$M(e^+e^- \rightarrow \mu^- \mu^- \gamma) = [M_{(1)} + M_{(2)}]_{\epsilon=\epsilon_q} = [M_{(3)} + M_{(4)}]_{\epsilon=\epsilon_p}. \quad (3.73)$$

By eq. (3.22), the calculation is straightforward

$$\begin{aligned}
M_{(1)}(-, +, +, -, +) &= 0, \\
M_{(2)}(-, +, +, -, +) &= 2ie^3 \frac{\langle pq \rangle^2}{\langle p'p \rangle \langle q'q \rangle} \beta(p', p, k), \\
M_{(3)}(-, +, +, -, +) &= 0, \\
M_{(4)}(-, +, +, -, +) &= -2ie^3 \frac{\langle pq \rangle^2}{\langle p'p \rangle \langle q'q \rangle} \beta(q', q, k), \tag{3.74}
\end{aligned}$$

where

$$\beta(p, q, k) = \frac{\sqrt{2} \langle pq \rangle}{\langle pk \rangle \langle qk \rangle}. \tag{3.75}$$

Therefore the calculation arrives at

$$\begin{aligned}
M(-, +, +, -, +) &= [M_{(1)} + M_{(2)} + M_{(3)} + M_{(4)}](-, +, +, -, +) \\
&= 2ie^3 \frac{\langle pq \rangle^2}{\langle p'p \rangle \langle q'q \rangle} [\beta(p', p, k) - \beta(q', q, k)]. \tag{3.76}
\end{aligned}$$

As for the norm we have

$$\begin{aligned}
|\beta(p, q, k)|^2 &= -\frac{1}{2} v^2(p, q, k), \\
|\beta(p', p, k) - \beta(q', q, k)|^2 &= -\frac{1}{2} [v(p', p, k) - v(q', q, k)]^2, \tag{3.77}
\end{aligned}$$

where

$$v(p, q, k) = \frac{p}{(pk)} - \frac{q}{(qk)}, \tag{3.78}$$

and

$$|M(-, +, +, -, +)|^2 = -2e^6 \frac{t'^2}{s s'} (v_p - v_q)^2, \tag{3.79}$$

where

$$\begin{aligned}
v_p &\equiv v(p', pk), & v_q &\equiv v(q', q, k) \\
s &= (p' + p)^2 = 2(p'p), & s &= (q' + q)^2 = 2(q'q), \\
t &= (p' - q')^2 = -2(p'q'), & t' &= (p - q)^2 = -2(pq), \\
u &= (p' - q)^2 = -2(p'q), & u' &= (p - q')^2 = -2(pq').
\end{aligned} \tag{3.80}$$

In a complete analogous way, we obtain

$$\begin{aligned}
|M(-, +, -, +, +)|^2 &= -2e^6 \frac{u'^2}{ss'} (v_p - v_q)^2, \\
|M(+, -, -, +, +)|^2 &= -2e^6 \frac{t^2}{ss'} (v_p - v_q)^2, \\
|M(+, -, +, -, +)|^2 &= -2e^6 \frac{u^2}{ss'} (v_p - v_q)^2.
\end{aligned} \tag{3.81}$$

By parity conjugation, we have

$$\begin{aligned}
|M(+, -, -, +, -)|^2 &= -2e^6 \frac{t'^2}{ss'} (v_p - v_q)^2, \\
|M(+, -, +, -, -)|^2 &= -2e^6 \frac{u'^2}{ss'} (v_p - v_q)^2, \\
|M(-, +, +, -, -)|^2 &= -2e^6 \frac{t^2}{ss'} (v_p - v_q)^2, \\
|M(-, +, -, +, -)|^2 &= -2e^6 \frac{u^2}{ss'} (v_p - v_q)^2.
\end{aligned} \tag{3.82}$$

Therefore, the averaged norm for polarized scattering is finally

$$\overline{|M|^2} = -2e^6 \frac{t^2 + u^2 + t'^2 + u'^2}{ss'} (v_p - v_q)^2. \tag{3.83}$$

This result is identical with that obtained with CALKUL method [49], but the calculation is simplified.

We have shown that we could simplify the calculation of the various helicity amplitudes for the single bremsstrahlung process $e^+e^- \rightarrow \mu^+\mu^-\gamma$ in the massless

QED by introduction the explicit helicity vectors in the "Chinese magic" polarization scheme. This technique described here could be utilized for all bremsstrahlung process. Furthermore the introduction of polarization vectors for the external gauge boson leads to simplifications in massless gauge theories.

3.4 Kleiss and Stirling Spinor Technique

Besides the CALKUL approach and "Chinese magic" polarization scheme, let us introduce another spinor product method, Kleiss and Stirling spinor technique [52].

First we derive expressions for the spinor products. We begin the discussion by establishing a convention for the overall complex phase of the spinors. Let us choose two four-vectors k_0^μ and k_1^ν with the following properties:

$$k_0 \cdot k_0 = 0, \quad k_1 \cdot k_1 = -1, \quad k_0 \cdot k_1 = 0. \quad (3.84)$$

Next we define the basic spinor $u_-(k_0)$ as follows:

$$u_-(k_0)\bar{u}_-(k_0) = \omega_- \not{k}_0, \quad (3.85)$$

where

$$\omega_\pm = \frac{1}{2}(1 \pm \gamma_5). \quad (3.86)$$

The spinor $u_-(k_0)$ is therefore the negative-helicity state of a massless fermion with momentum k_0 . The corresponding positive-helicity state is chosen to be

$$u_+(k_0) = \not{k}_1 u_-(k_0). \quad (3.87)$$

From these two spinors we could construct spinors for any other lightlike momentum p as follows:

$$u_\lambda(p) = \frac{\not{p} u_{-\lambda}(k_0)}{\sqrt{2p \cdot k_0}}. \quad (3.88)$$

From eqs. (3.84)-(3.88) we can derive some useful relations for spinor sandwiches for the following discussion. Let Γ be an arbitrary string of γ matrices, and let Γ^R be the same string in the reversed order. Then we find, for arbitrary lightlike momenta and helicities:

$$\bar{u}_{\lambda_1}(p_1)\Gamma u_{\lambda_2}(p_2) = \lambda_1\lambda_2\bar{u}_{-\lambda_2}(p_2)\Gamma^R u_{-\lambda_1}(p_1). \quad (3.89)$$

The second useful identity (Chisholm identity) is

$$\{\bar{u}_{\lambda_1}(p_1)\gamma^\mu u_{\lambda_2}(p_2)\}\gamma_\mu = 2u_\lambda(p_2)\bar{u}_\lambda(p_1) + 2u_{-\lambda}(p_1)\bar{u}_{-\lambda}(p_2). \quad (3.90)$$

Next we discuss the spinor products themselves. For massless fermions with momenta p_1 and p_2 there are two non-zero products:

$$\begin{aligned} s(p_1, p_2) &\equiv \bar{u}_+(p_1)u_-(p_2) = -s(p_2, p_1), \\ t(p_1, p_2) &\equiv \bar{u}_-(p_1)u_+(p_2) = [s(p_2, p_1)]^*. \end{aligned} \quad (3.91)$$

Using eqs. (3.85) and (3.88), we could evaluate $s(p_1, p_2)$:

$$\begin{aligned} s(p_1, p_2) &= \frac{\bar{u}_-(k_0)\not{p}_1\not{p}_2 u_+(k_0)}{\sqrt{4(p_1 \cdot k_0)(p_2 \cdot k_0)}} = \frac{\text{tr}(\omega_- \not{k}_0 \not{p}_1 \not{p}_2 \not{k}_1)}{\sqrt{4(p_1 \cdot k_0)(p_2 \cdot k_0)}} \\ &= \frac{[(p_1 \cdot k_0)(p_2 \cdot k_1) - (p_1 \cdot k_1)(p_2 \cdot k_0) - i\epsilon_{\mu\nu\rho\sigma}k_0^\mu k_0^\nu p_1^\rho p_2^\sigma]}{\sqrt{4(p_1 \cdot k_0)(p_2 \cdot k_0)}}. \end{aligned} \quad (3.92)$$

In a practical calculation we can specify k_0^μ and k_1^μ such that the form of $s(p_1, p_2)$ becomes compact. For instance, we take

$$\begin{aligned} p_i^\mu &= (p_i^0, p_i^x, p_i^y, p_i^z), \\ k_0^\mu &= (0, 0, 0, 0), \\ k_1^\mu &= (0, 0, 1, 0). \end{aligned} \quad (3.93)$$

This leads to

$$s(p_1, p_2) = (p_1^y + ip_1^z) \left[\frac{p_2^0 - p_2^x}{p_1^0 - p_1^x} \right]^{\frac{1}{2}} - (p_2^y + ip_2^z) \left[\frac{p_2^0 - p_1^x}{p_2^0 - p_2^x} \right]^{\frac{1}{2}} \quad (3.94)$$

and $t(p_1, p_2)$ is obtained by complex conjugation. Furthermore, if we have calculated the spinor product, we immediately obtain the vector product as well:

$$d(p_1, p_2) \equiv 2p_1 \cdot p_2 = |s(p_1, p_2)|^2. \quad (3.95)$$

Analogous to the CALKUL and "Chinese magic" scheme, the spinor products play a more fundamental role than vector products in practical calculations.

We now proceed to construct the polarization vectors of the external gauge bosons. Let us consider polarization vectors with states of definite helicities, denoted by ϵ_{\pm}^{μ} and satisfying

$$\begin{aligned} \epsilon_{\lambda} \cdot k &= 0, & \epsilon_{\lambda} \cdot \epsilon_{\lambda} &= 0, \\ \epsilon_{-\lambda}^{\mu} &= (\epsilon_{\lambda}^{\mu})^*, & \epsilon_{\lambda} \cdot \epsilon_{-\lambda} &= -1, \end{aligned} \quad (3.96)$$

where k_{μ} is the photon momentum, and $\lambda = \pm 1$. As long as the polarization vectors ϵ_{λ}^{μ} obey the relations (3.96), it is an acceptable choice. We use the following convention:

$$\epsilon_{\lambda}^{\mu} = N \bar{u}_{\lambda}(k) \gamma^{\mu} u_{\lambda}(p), \quad N = [4(p \cdot k)]^{\frac{1}{2}}, \quad (3.97)$$

where p^{μ} is an arbitrary vector not collinear to k^{μ} or k_0^{μ} .

Spinors for the massive particle with four momentum p (with $p^2 = m^2$) could also be defined in an analogous way in which the spinor for massless particles [54]:

$$\begin{aligned} u(p, \lambda) &= \frac{1}{\sqrt{2p \cdot k}} (\not{p} + m) u_{-\lambda}(k), \\ v(p, \lambda) &= \frac{1}{\sqrt{2p \cdot k}} (\not{p} - m) u_{\lambda}(k). \end{aligned} \quad (3.98)$$

where k^{μ} is an arbitrary lightlike vector.

The definition (3.98) can be rewritten in terms of massless spinor as follows

$$\begin{aligned} u(p, \lambda) &= u_\lambda(p_k) + \frac{m}{\sqrt{2p \cdot k}} u_{-\lambda}(k), \\ v(p, \lambda) &= u_{-\lambda}(p_k) - \frac{m}{\sqrt{2p \cdot k}} u_\lambda(k), \end{aligned} \quad (3.99)$$

where

$$p_k = p - k \frac{m^2}{2p \cdot k}, \quad p_k^2 = 0 \quad (3.100)$$

is the light-cone projection of p obtained with the help of the auxiliary vector k .

Therefore we could obtain the explicit expressions of spinor products for the massive spinors by eq. (3.98):

$$\left\{ \begin{aligned} \bar{u}(p_1, \lambda_1) u(p_2, \lambda_2) &= S(p_1, m_1, \lambda_1, p_2, m_2, \lambda_2), \\ \bar{u}(p_1, \lambda_1) v(p_2, \lambda_2) &= S(p_1, m_1, \lambda_1, p_2, -m_2, -\lambda_2), \\ \bar{v}(p_1, \lambda_1) u(p_2, \lambda_2) &= S(p_1, -m_1, -\lambda_1, p_2, m_2, \lambda_2), \\ \bar{v}(p_1, \lambda_1) v(p_2, \lambda_2) &= S(p_1, -m_1, -\lambda_1, p_2, -m_2, -\lambda_2). \end{aligned} \right. \quad (3.101)$$

where

$$S(p_1, m_1, \lambda_1, p_2, m_2, \lambda_2) = \delta_{\lambda_1, -\lambda_2} s_{\lambda_1}(p_{1k}, p_{2k}) + \delta_{\lambda_1, \lambda_2} \left(m_1 \sqrt{\frac{2p_2 k}{2p_1 k}} + m_2 \sqrt{\frac{2p_1 k}{2p_2 k}} \right). \quad (3.102)$$

Up to now we have introduced a calculational tool for helicity amplitudes, which brings out simplifications in gauge theories. And this approach shares many essential features with CALKUL helicity amplitude method and "Chinese magic" polarization scheme. This formalism can be directly programmed into a computer due to its symbolic properties, which plays important roles in perturbative calculations in high energy physics.


CHAPTER FOUR

Numerical Computations for the Five Point Function

The one loop integrals play an important role in calculating radiative corrections in particle physics since they are manageable for fast MC event generator execution for arbitrary masses and kinematics for high energy scattering process. It has been demonstrated that n -point functions at one-loop level can be reduced to scalar one loop functions [30,31]. Considering representations of the scalar four-point function for arbitrary masses and momenta relevant to most high energy collider applications have been given and they fit MC implementation well, it is natural to attempt expressing higher point-function ($n \geq 5$) in terms of the 1, 2, 3 and 4-point functions. When the problem becomes reducing higher-point functions into expressions in terms of the 1,2,3 and 4-point functions, we are most concerned about the numerical stability. To solve this problem, B. F. L. Ward presented an approach to evaluate higher point loop integrals using Chinese magic in the virtual loop integration variable [55], which is called "magic spinor product methods in loop integrals". Based on this method, we developed a program to compute the general five-point function numerically. In this chapter, we first introduce the physics content of "magic spinor product methods in loop integrals" and then we compare our results with those from LoopTools [56], which is a package for computation of one-loop integrals based on the FF package by G. J. van. Oldenborgh [57]. By comparison, it shows the result from our program is accurate and numerically stable.

4.1 Magic Spinor Product Approach in Loop Integrals

Here we use the conventions of Refs. [54,58] which are derived from Refs. [51,52]. The five-point function which we analyze is shown in Feynman diagram (c) as follows. It has many applications in collider precision physics. For example, together with Feynman diagrams (a) and (b) it generates a gauge invariant contribution to the ISR for $e^+e^- \rightarrow f\bar{f} + \gamma$, $f \neq e$. We focus on the application of Chinese magic in the loop integral in Feynman diagram (c) to illustrate the possible simplifications here.



Applying the Feynman rules, we have

$$\begin{aligned}
M_{\lambda_1 \lambda_2 \lambda'_1 \lambda'_2 \lambda_\gamma}^{(1c)} &= (2\pi)^4 \delta(p_1 + p_2 - p'_1 - p'_2 - k) \mathcal{C} \\
&\times \int \frac{d^4 q}{(2\pi)^4} \frac{\bar{v}_{\lambda_2} \gamma^\beta (\not{q} + \not{p}_1 - \not{k} + m_1) \not{\epsilon}_{\lambda_\gamma}^* (\not{q} + \not{p}_1 + m_1) \gamma^\alpha u_{\lambda_1}}{[(q + p_1 - k)^2 - m_1^2 + i\epsilon][(q + p_1)^2 - m_1^2 + i\epsilon]} \\
&\frac{\bar{u}'_{\lambda'_1} \gamma_\alpha (\not{q} + \not{p}'_1 + m_2) \gamma_\beta v'_{\lambda'_2}}{[(q + p_1 + p_2 - k)^2 - M_{V_2}^2 + i\epsilon][(q + p'_1)^2 - m_2^2 + i\epsilon](q^2 - M_{V_1}^2 + i\epsilon)} \\
&+ \dots, \tag{4.1}
\end{aligned}$$

where we define massless limit coupling factor

$$\begin{aligned}
\mathcal{C} &= \mathcal{C}(\lambda_i, \lambda'_j) \\
&= Q_1 e G^2 G'^2 (v'_1 + a'_1 \lambda_2)(v_1 - a_1 \lambda_1)(v'_2 + a'_2 \lambda_2)(v_2 + a_2 \lambda'_1) \tag{4.2}
\end{aligned}$$

with the couplings $Q_1 e$, G , and G' for the γ , V_1 and V_2 respectively. In order to obtain the loop integral in terms of Chinese magic, we take the following kinematics

$$\begin{aligned}
p_1 &= (E, p\hat{z}) \\
p_2 &= (E, -p\hat{z}) \\
-p_4 &= (E', p'(\cos\theta'_1 \hat{z} + \sin\theta'_1 \hat{x})) \equiv p'_1 \\
k &= (k_0, k(\cos\theta_\gamma \hat{z} + \sin\theta_\gamma(\cos\phi_\gamma \hat{x} + \sin\phi_\gamma \hat{y}))) \\
p_1 + p_2 &= -p_4 - p_3 + k = (\sqrt{s}, \vec{0}) \\
-p_3 &\equiv p'_2, \tag{4.3}
\end{aligned}$$

with $k^0 = k$, $\sqrt{s} = 2E$. Besides, we introduce the alternative notations $p'_1 = -p_4$, $p'_2 = -p_3$. Now we introduce the two sets of magic polarization vectors associated to the two incoming lines

$$\begin{aligned}
(\epsilon_\sigma^\mu(\beta))^* &= \frac{\bar{u}_\sigma(k) \gamma^\mu u_\sigma(\beta)}{\sqrt{2} \bar{u}_{-\sigma} u_\sigma(\beta)}, \\
(\epsilon_\sigma^\mu(\zeta))^* &= \frac{\bar{u}_\sigma(k) \gamma^\mu u_\sigma(\zeta)}{\sqrt{2} \bar{u}_{-\sigma} u_\sigma(\zeta)}, \tag{4.4}
\end{aligned}$$

with $\beta^2 = 0$ and $\zeta^2 = 0$. We choose the basis of the four-dimensional momentum space as follows:

$$\begin{aligned}
\ell_1 &= (E, E\hat{z}) \\
\ell_2 &= (E, -E\hat{z}) \\
\ell_3 &= E \frac{\langle \ell_2 + |\gamma^\mu| \ell_1 + \rangle}{\sqrt{2} \langle \ell_2 - |\ell_1 + \rangle} = -\frac{E}{\sqrt{2}}(\hat{x} + i\hat{y}) \\
\ell_4 &= E \frac{\langle \ell_2 - |\gamma^\mu| \ell_1 - \rangle}{\sqrt{2} \langle \ell_2 + |\ell_1 - \rangle} = \frac{E}{\sqrt{2}}(\hat{x} - i\hat{y})
\end{aligned} \tag{4.5}$$

where we use the Dirac notations in Refs. [50, 52, 55]. Note that all four of the basis four-vector are lightlike so that they can attend in Chinese magic.

We define the loop momentum as

$$q = \alpha_i \ell_i \tag{4.6}$$

with summation over repeated indices. The coefficient α_i 's are dertermined as

$$\begin{aligned}
\alpha_1 &= \frac{q\ell_2}{2E^2} = \frac{1}{s}(D_3 - D_2 - s + 2p_2k + M_{V_2}^2), \\
\alpha_2 &= \frac{q\ell_1}{2E^2} = \frac{1}{s}(D_1 - D_0 - M_{V_1}^2), \\
\alpha_3 &= \frac{q\ell_4}{E^2} = -\frac{q\ell_3^*}{E^2} = -\alpha_4^*, \\
\alpha_4 &= -\frac{i}{\sqrt{2s}}[c_j D_j + c_5 M_{V_1}^2 + c_6(M_{V_2}^2 + 2p_2k - s) + c_7(2kp_1)],
\end{aligned} \tag{4.7}$$

where we define the denominators as

$$\begin{aligned}
D_0 &= q^2 - M_{V_1}^2 + i\epsilon \\
D_1 &= (q + p_1)^2 - m_1^2 + i\epsilon \\
D_2 &= q + p_1 - k^2 - m_1^2 + i\epsilon \\
D_3 &= q + p_1 + p_2 - k^2 - M_{V_2}^2 + i\epsilon \\
D_4 &= (q - p_4)^2 - m_2^2 + i\epsilon
\end{aligned} \tag{4.8}$$

such that the expansion coefficients $\{c_j\}$ are

$$\begin{aligned}
c_0 &= \csc \phi_\gamma \left(\frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 E'_1} - \frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 \sqrt{s}} + \frac{\csc \theta_\gamma}{\sqrt{s}} - \frac{\cot \theta'_1 e^{i\phi_\gamma} - \cot \theta_\gamma}{\beta_1 \sqrt{s}} \right), \\
c_1 &= \csc \phi_\gamma \left(\frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 \sqrt{s}} - \frac{\csc \theta'_1}{\sqrt{s}} + \frac{\cot \theta'_1 e^{i\phi_\gamma} - \cot \theta_\gamma}{\beta_1 \sqrt{s}} + \frac{\csc \theta_\gamma}{k^0} \right), \\
c_2 &= \csc \phi_\gamma \left(\frac{-\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 \sqrt{s}} + \frac{\csc \theta_\gamma}{\sqrt{s}} + \frac{\cot \theta'_1 e^{i\phi_\gamma} - \cot \theta_\gamma}{\beta_1 \sqrt{s}} - \frac{\csc \theta_\gamma}{k^0} \right), \\
c_3 &= \csc \phi_\gamma \left(\frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 \sqrt{s}} - \frac{\csc \theta_\gamma}{\sqrt{s}} - \frac{\cot \theta'_1 e^{i\phi_\gamma} - \cot \theta_\gamma}{\beta_1 \sqrt{s}} \right), \\
c_4 &= -\csc \phi_\gamma \frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 E'_1}, \\
c_5 &= \csc \phi_\gamma \left(\frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 E'_1} - \frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 \sqrt{s}} + \frac{\csc \theta_\gamma}{\sqrt{s}} - \frac{\cot \theta'_1 e^{i\phi_\gamma} - \cot \theta_\gamma}{\beta_1 \sqrt{s}} \right), \\
c_6 &= \csc \phi_\gamma \left(\frac{\csc \theta'_1 e^{i\phi_\gamma}}{\beta'_1 \sqrt{s}} - \frac{\csc \theta_\gamma}{\sqrt{s}} - \frac{\cot \theta'_1 e^{i\phi_\gamma} - \cot \theta_\gamma}{\beta_1 \sqrt{s}} \right), \\
c_7 &= -\csc \phi_\gamma \frac{\csc \theta_\gamma}{k^0}, \tag{4.9}
\end{aligned}$$

with $\beta = \frac{p}{E}$ and $\beta'_1 = \frac{p'}{E'}$. Therefore the $\{c_j\}$ are determined by the kinematics that we choose. Note that the Chinese magic now carries over to the loop variable via the identity

$$\begin{aligned}
\not{q} &= \alpha_j \not{\ell}_j \\
&= \sum_{j=1}^2 \alpha_j (|\ell_{j+}\rangle \langle \ell_{j+}| + |\ell_{j-}\rangle \langle \ell_{j-}|) + \alpha_3 \frac{\sqrt{2}E}{\langle p_2 - | p_1 + \rangle} (|\ell_{2-}\rangle \langle \ell_{1-}| + |\ell_{1-}\rangle \langle \ell_{2+}|) \\
&\quad + \alpha_4 \frac{\sqrt{2}E}{\langle p_2 + | p_1 - \rangle} (|\ell_{2+}\rangle \langle \ell_{1+}| + |\ell_{1-}\rangle \langle \ell_{2-}|) \\
&\equiv \sum_{j=1}^2 \alpha_j (|p_{j+}\rangle \langle p_{j+}| + |p_{j-}\rangle \langle p_{j-}|) + \alpha_3 \frac{\sqrt{2}E}{\langle p_2 - | p_1 + \rangle} (|p_{2-}\rangle \langle p_{1-}| + |p_{1-}\rangle \langle p_{2+}|) \\
&\quad + \alpha_4 \frac{\sqrt{2}E}{\langle p_2 + | p_1 - \rangle} (|p_{2+}\rangle \langle p_{1+}| + |p_{1-}\rangle \langle p_{2-}|) \\
&\equiv \sum_{j=1}^2 \alpha_j (|p_{j+}\rangle \langle p_{j+}| + |p_{j-}\rangle \langle p_{j-}|) + \tilde{\alpha}_3 (|p_{2-}\rangle \langle p_{1-}| + |p_{1-}\rangle \langle p_{2+}|) \\
&\quad + \tilde{\alpha}_4 (|p_{2+}\rangle \langle p_{1+}| + |p_{1-}\rangle \langle p_{2-}|), \tag{4.10}
\end{aligned}$$

where we let $\ell_1 \equiv p_1$, $\ell_2 \equiv p_2$ since for the numerator algebra we work in the massless limit. And we define as well

$$\begin{aligned}\tilde{\alpha}_3 &\equiv \alpha_3 \frac{\sqrt{2}E}{\langle p_2 - |p_1+\rangle} = -\frac{\alpha_3}{\sqrt{2}}, \\ \tilde{\alpha}_4 &\equiv \alpha_4 \frac{\sqrt{2}E}{\langle p_2 + |p_1-\rangle} = -\frac{\alpha_4}{\sqrt{2}}.\end{aligned}\tag{4.11}$$

Next, we introduce the representation (4.10) into the numerator, N , of the integrand in eq (4.1). Then we have the reduction

$$\begin{aligned}N &= \frac{4\sqrt{2}}{\langle k - |p_1+\rangle} \{ (A_1 \langle p_2 + |p_1'-\rangle \langle p_2' - |p_2+\rangle \\ &\quad + A_2 \langle p_2 + |p_1'-\rangle \langle p_2' - |p_1+\rangle) \times (A_3 \langle p_2 + |p_1'-\rangle \langle p_1' - |p_1+\rangle \\ &\quad + A_4 \langle p_1 + |p_1'-\rangle \langle p_1' - |p_1+\rangle) \\ &\quad + \tilde{\alpha}_4 (A_1 \langle p_2 + |p_1-\rangle \langle p_2' - |p_2+\rangle + A_2 \langle p_2 + |p_1-\rangle \langle p_2' - |p_1+\rangle) \\ &\quad \times (A_3 \langle p_2 + |p_1'-\rangle \langle p_2 - |p_1+\rangle \\ &\quad + A_4 \langle p_1 + |p_1'-\rangle \langle p_2 - |p_1+\rangle) \}\end{aligned}\tag{4.12}$$

from the standard identities

$$\begin{aligned}\not{\epsilon}_{\lambda_\gamma}^* &= \frac{\sqrt{2}}{\langle k - \lambda_\gamma | \ell_1 \lambda_\gamma \rangle} [|\ell_1 \lambda_\gamma\rangle \langle k \lambda_\gamma| + |k - \lambda_\gamma\rangle \langle \ell_1 - \lambda_\gamma|], \\ \gamma^\rho \langle \ell_1 \lambda | \gamma_\rho | \ell_2 \lambda \rangle &= 2[|\ell_1 - \lambda\rangle \langle \ell_2 - \lambda| + |\ell_2 \lambda\rangle \langle \ell_1 \lambda|], \\ \ell_1 &= |\ell_1+\rangle \langle \ell_1+| + |\ell_1-\rangle \langle \ell_1-|,\end{aligned}\tag{4.13}$$

where we defined

$$\begin{cases} A_1 = \tilde{\alpha}_4 \langle p_1 + |k-\rangle + \alpha_2 \langle p_2 + |k-\rangle, \\ A_2 = (1 + \alpha_1) \langle p_1 + |k-\rangle + \tilde{\alpha}_3 \langle p_2 + |k-\rangle, \\ A_3 = \alpha_2 \langle p_1 - |p_2+\rangle, \\ A_4 = \tilde{\alpha}_4 \langle p_1 - |p_2+\rangle, \end{cases}\tag{4.14}$$

for the magic choice $\beta = p_1$. Note that the Chinese magic trick has eliminated all but one set of the terms with three factors of the virtual momentum expansion coefficients. Moreover, in the numerator of the propagator before or after the real emission vertex, this trick has annihilated the terms associated with $\not{p}_1 \not{k}$ as well as half of the terms in the respective virtual momentum expansion in the former case. Compared to the traditional approach of taking traces on the fermion lines, the Chinese magic trick has a large fraction of the terms on the right-hand side of eq. (4.12). Specifically speaking, We need to compute $2\Re\mathcal{M}_B^*\mathcal{M}^{(1c)}$ in the usual method of taking traces of fermion lines, where \mathcal{M}_B is the respective Born amplitude that would interfere with the one-loop amplitude to generate the one-loop correction of the respective cross section. In the Chinese magic representation, we find that only radiation from the antifermion (p_2) incoming line contributes.

$$\begin{aligned}
\mathcal{M}_{B+-+--+} &= (2\pi)^4 \delta(p_1 + p_2 - p'_1 - p'_2 - k) \\
&\times \frac{2\sqrt{2}ieQ_1 G_j^2 (v'_j - a'_j)(v_j - a_j) \langle p'_2 - |p_1+\rangle}{\langle k - |p_1+\rangle \langle k - |p_2+\rangle (s' - M_{V_j}^2 + i\epsilon)} \\
&\times [\langle p_1 - |p_2+\rangle \langle p_2 + |p'_1-\rangle - \langle p_1 - |k+\rangle \langle k + |p'_1-\rangle].
\end{aligned} \tag{4.15}$$

Therefore the calculation for $2\Re\mathcal{M}_B^*\mathcal{M}^{(1c)}$ just involves multiplying N in eq. (4.12) by the complex conjugate of the simple expression and taking twice the real part.

Assume that the traditional approach of taking traces of fermion lines is applied, we need the trace of two sets of terms with 10 Dirac gamma matrices multiplied by a factor with the trace of 6 Dirac gamma matrices. If so, we will have $2 \cdot 9 \cdot 7 \cdot 5 \cdot 4 \times 5 \cdot 4 = 2520 \times 20 = 50400$ terms. Each terms requires Passarino-Veltman (PV) reduction of three, two, and one five-point tensor integrals. By comparison,

we can appreciate the great simplification that eq. (4.12) represents. We see that this form of N in eq. (4.12) has efficiently reduced the problem of reduction of the five-point function with three, two, one tensor indices in the PV scheme to the question of a single scalar five-point and lower four, three, and two point functions with the coefficients already explicitly written in terms of the center of mass (CMS) kinematic variables which are important to efficient MC generations. Furthermore, if we introduce the result of N in eq. (4.12) into the integral in eq. (4.1), we have the integrals

$$\int \frac{d^4q}{(2\pi)^4} \frac{D_i D_j D_k; D_i D_j; D_j; 1}{D_0 D_1 D_2 D_3 D_4}, \quad i, j, k = 0, \dots, 4 \quad (4.16)$$

all of which are determined from the lower point functions when the results for the representation of the scalar five-point function in terms of four-point functions used in Refs. [43, 59–61]. Then we obtain an advantage: no calculation of wave functions at complex momenta is required here. Our work results rigorously from Lagrangian quantum field theory and thus it could be used as a cross check on other possible approaches.

So far we have simplified considerably the computation and have removed the Gram determinant-type factors in the tensor integral reductions. But our calculation still depends on the Gram determinant-type denominator factors in the results in Refs. [43, 59–61] for the representation of the five-point function in terms of scalar four-point functions. In order to achieve a better numerical stability, we need to replace the representation of the scalar five-point function.

Let us begin with the identity

$$\begin{aligned}
q^2 &= D_0 + M_{V_1}^2 - i\epsilon = (\alpha_i \ell_i)^2 \\
&= 2\alpha_1 \alpha_2 \ell_1 \ell_2 + 2\alpha_3 \alpha_4 \ell_3 \ell_4 = s\alpha_1 \alpha_2 + \frac{s}{2}\alpha_3 \alpha_4.
\end{aligned} \tag{4.17}$$

Dividing by $D_0 \cdots D_4$ and integrating over d^4q we have the following representation of the required scalar five-point function

$$\begin{aligned}
&E_0(\bar{p}_1, \bar{p}_2, \bar{p}_3, \bar{p}_4, \bar{m}_0, \bar{m}_1, \bar{m}_2, \bar{m}_3, \bar{m}_4) \\
&= \frac{1}{C_{E_0}} \left\{ -D_0(0) + \frac{1 + \beta_1^2}{2s\beta_1^2} \left[C_0(13) - C_0(12) - C_0(03) + C_0(02) \right. \right. \\
&+ (M_{V_2}^2 - s + 2p_2k) \left(D_0(1) - D_0(0) \right) - M_{V_1}^2 \left(D_0(3) - D_0(2) \right) \left. \right] \\
&- \frac{1 - \beta_1^2}{4s\beta_1^2} \left[\Delta r_{1,0} \left(D_0(1) - D_0(0) \right) + 2\Delta \bar{p}_{1,0} \left(D_{11}(1)\bar{p}(1)_1 - D_{11}(0)\bar{p}(0)_1 \right. \right. \\
&+ D_{12}(1)\bar{p}(1)_2 - D_{12}(0)\bar{p}(0)_2 + D_{13}(1)\bar{p}(1)_3 - D_{13}(2)\bar{p}(2)_3 - D_0(3)\bar{p}(3)_4 \\
&+ D_0(2)\bar{p}(2)_4 \left. \right) + 2(M_{V_2}^2 - s + 2p_2k) \left(D_0(3) - D_0(2) \right) \left. \right] \\
&- \frac{1}{4} \left[\sum_{j=0}^4 |c_j|^2 \left(C_0(j, j+1) + \Delta r_{j, j+1} D_0(j) + 2\Delta \bar{p}_{j, j+1} (D_{11}(j)\bar{p}(j)_1 \right. \right. \\
&+ D_{12}(j)\bar{p}(j)_2 + D_{13}(j)\bar{p}(j)_3 - D_0(j)\bar{p}(j)_4) \left. \right) \\
&+ 2 \left(\sum_{i>j}^4 \Re(c_i c_j^*) C_0(ij) + \sum_{j=0}^4 \Re(c_j (c_5^* M_{V_1}^2 + c_6^* (M_{V_2}^2 - s + 2p_2k) + c_7^* (2kp_1))) \right. \\
&\left. \left. D_0(j) \right) \right] \left. \right\} \tag{4.18}
\end{aligned}$$

where $\bar{p} = p_1$, $\bar{p}_2 = p_1 - k$, $\bar{p}_3 = p_1 + p_2 - k$, $\bar{p} = p'_1$, $\bar{m}_0 = M_{V_1}$, $\bar{m}_1 = m_1$, $\bar{m} = m_1$, $\bar{m}_3 = M_{V_2}$, $\bar{m}_4 = m_2$, where the coefficient C_{E_0} is

$$\begin{aligned}
C_{E_0} = & M_{V_1}^2 - i\epsilon + \frac{1 + \beta_1^2}{2\beta_1^2 s} M_{V_1}^2 (M_{V_2}^2 - s + 2p_2 k) + \frac{1 - \beta_1^2}{4\beta_1^2 s} (M_{V_1}^4 \\
& + (M_{V_2}^2 - s + 2p_2 k)^2) + \frac{1}{2} \Re[c_5 c_6^* M_{V_1}^2 (M_{V_2}^2 - s + 2p_2 k) \\
& + c_5 c_7^* M_{V_1}^2 (2kp_1) + c_6 c_7^* (M_{V_2}^2 - s + 2p_2 k) (2kp_1)] \\
& + \frac{1}{4} [|c_5|^2 M_{V_1}^4 + |c_6|^2 (M_{V_2}^2 - s + 2p_2 k)^2 + |c_7|^2 (2kp_1)^2]. \tag{4.19}
\end{aligned}$$

We have used a mixture of notations from [31, 59–61] such that

$$D_j = (q - \bar{p}_j)^2 - \bar{m}_j^2 + i\epsilon = q^2 + 2q\bar{p}_j + \bar{p}_j^2 - \bar{m}_j^2 + i\epsilon \equiv q^2 + 2q\bar{p}_j + r_j,$$

$$\Delta_{i,j} \equiv r_i - r_j,$$

$$\Delta \bar{p}_{i,j} \equiv \bar{p}_i - \bar{p}_j$$

$D_0(j) \equiv$ four-point scalar function obtained from five-point scalar function

by omitting denominator D_j ,

$C_0(i, j) \equiv$ three-point scalar function obtained from five-point scalar function

by omitting denominator D_i and D_j , $i \neq j$ (4.20)

where we also use the Passarino-Veltman [31] notation of the four-point one-tensor integral, $D_\mu(j)$, obtained from the five-point one tensor integral function by omitting the denominator D_j , with

$$D_\mu(j) \equiv D_{11}(j) \bar{\bar{p}}(j)_1 + D_{12}(j) \bar{\bar{p}}(j)_2 + D_{13}(j) \bar{\bar{p}}(j)_3 - D_0(j) \bar{\bar{p}}(j)_4,$$

where the four-vectors $\{\bar{\bar{p}}(j)_j\}$ are determined according to Ref. [31]. Note that $\bar{\bar{p}}(j)_4$ is only nonzero if it is necessary to shift the q -integration variable by it to reach the standard form of Passarino-Veltman representation. This expression for E_0 have no

problem with Gram determinant type denominators.

$$\begin{aligned}
\mathcal{M}^{(1a)}_{+--+} &= 0, \text{ by Chinese magic trick} \\
\mathcal{M}^{(1b)}_{+--+} &= (2\pi)^4 \delta(p_1 + p_2 - p'_1 - p'_2 - k) \\
&\quad \times \frac{4\sqrt{2}\mathcal{C}}{\langle k - |p_{1+}\rangle \langle k - |p_{2+}\rangle} \int \frac{d^4q}{(2\pi)^4} \frac{N'}{D_0 D_1 D_3 D_4}, \tag{4.21}
\end{aligned}$$

where the numerator N' is given by

$$\begin{aligned}
N' &= (\langle p'_2 - |p_{1+}\rangle a_1 + \langle p'_2 - |p_{2+}\rangle b_1) \\
&\quad \times (\langle p_1 - |p_{2+}\rangle \langle p_2 + |p'_{1-}\rangle - \langle p_1 - |k+\rangle \langle k + |p'_{1-}\rangle) \\
&\quad + (\langle p'_2 - |p_{1+}\rangle \bar{a}_1 + \langle p'_2 - |p_{2+}\rangle \bar{b}_1) \\
&\quad \times [-2p_1(p_2 - k)\tilde{\alpha}_4 + \alpha_2 \langle p_1 - |k+\rangle \langle k + |p_{2-}\rangle] \tag{4.22}
\end{aligned}$$

with

$$\begin{aligned}
a_1 &= (1 + \alpha_1)(2p_1 p'_1) + \alpha_3 \langle p_2 + |p'_{-}\rangle \langle p'_1 - |p_{1+}\rangle, \\
b_1 &= \alpha_2 \langle p_2 + |p'_{1-}\rangle \langle p'_1 - |p_{1+}\rangle + \tilde{\alpha}_4(2p_1 p'_1), \\
\bar{a}_1 &= \langle p_1 - |p_{2+}\rangle [(1 + \alpha_1) \langle p_1 + |p'_{1-}\rangle + \tilde{\alpha}_3 \langle p_2 + |p'_{1-}\rangle], \\
\bar{b}_1 &= \langle p_1 - |p_{2+}\rangle [\alpha_2 \langle p_2 + |p'_{1-}\rangle + \tilde{\alpha}_4 \langle p_1 + |p'_{1-}\rangle]. \tag{4.23}
\end{aligned}$$

As we see this method gives a considerable reduction to the known scalar functions while computing $2\Re\mathcal{M}_B^* \mathcal{M}^{1b}$ compared to the traditional method of tracing over fermion lines.

In sum, we have exhibited that the application of Chinese magic technique in the virtual loop momentum could reduce significantly the volume of algebra required for efficient and stable physical calculations of higher point virtual corrections with the general mass scale. Furthermore, we could construct computer realizations of the


method described above for evaluating the five-point function. In the next section, we will exhibit our results and compare them with those from LoopTools.

4.2 Numerical Results for the Five-Point Function E_0

Based on the magic spinor product method, we have developed a computer program to calculate the five-point function E_0 . In our program, the LoopTools package is included. The five-point function E_0 is calculated with the help of eqs. (4.9), (4.18) and (4.19). Scalar three-point functions $C_0(i, j)$, scalar four-point functions $D_0(j)$ and tensor four-point functions $D_{\mu\nu}(j)$ are calculated by LoopTools. To be specific, we choose

$$\left\{ \begin{array}{l} \sqrt{s} = 500 \text{ GeV}, \\ m_1 = m_e = 0.510999 \times 10^{-3} \text{ GeV}, \\ m_2 = m_\mu = 0.1056583 \text{ GeV}, \\ M_{V_1} = M_{V_2} = 91 \text{ GeV}, \end{array} \right.$$

and the kinematics is determined by eq. (4.3).






Figure 4.1: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 0^\circ$




Figure 4.2: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 15^\circ$

then by eq. (4.8) and (4.20), we have the expressions of $D_0(j)$ and $C_0(i, j)$ which give inputs of the corresponding three- and four-point functions in Looptools so that we can compute $D_0(j)$, $C_0(i, j)$ and $D_{\mu\nu}(i, j)$ via Looptools. Besides, we have computed E_0 via Looptools directly as a comparison. We choose $(\phi_\gamma, \theta_\gamma, \theta'_1)$ as the variables. The percent Differences of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{looptools}(\phi_\gamma, \theta_\gamma)$ are shown in Figure (4.1)-(4.13).




Figure 4.3: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 30^\circ$




Figure 4.4: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 45^\circ$




Figure 4.5: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 60^\circ$




Figure 4.6: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{\text{LoopTools}}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 75^\circ$




Figure 4.7: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{\text{LoopTools}}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 90^\circ$




Figure 4.8: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{\text{LoopTools}}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 105^\circ$




Figure 4.9: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 120^\circ$




Figure 4.10: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 135^\circ$




Figure 4.11: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{LoopTools}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 150^\circ$




Figure 4.12: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{\text{LoopTools}}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 165^\circ$




Figure 4.13: Percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{\text{LoopTools}}(\phi_\gamma, \theta_\gamma)$ for $\sqrt{s} = 500$ GeV, $M_{V_1} = 91$ GeV, $M_{V_1} = 91$ GeV and $\theta'_1 = 180^\circ$

As we see, the result from magic spinor product method agrees with that from LoopTools in overall, except several regions. The percentage difference of $E_0(\phi_\gamma, \theta_\gamma)$ and $E_0^{\text{LoopTools}}(\phi_\gamma, \theta_\gamma)$ is insensitive with ϕ_γ , but sensitive with θ'_1 . When $\theta'_1 > 90^\circ$, our result mainly fits to that from LoopTool in the region ($0^\circ < \phi_\gamma < 180^\circ$, $0^\circ < \theta_\gamma < 20^\circ$). However, when $\theta'_1 < 75^\circ$, our result agrees greatly with that from Looptools.

CHAPTER FIVE

Quantum Chromodynamics

5.1 Introduction to Quantum Chromodynamics

We now come to the other constituent of the standard model of particle physics, Quantum Chromodynamics (QCD) [6–11]. Quantum Chromodynamics is a non-Abelian gauge theory of strong interactions. The gauge symmetry of QCD is $SU(3)$ color. The choice of gauge group must rely on three facts: (a) the group must admit complex representations in order to distinguish a quark from antiquark; (b) the group must have completely antisymmetric color singlet to solve the statistical puzzle for the lowest lying baryons of spin 1/2 and 3/2; (c) the number of colors for each kind of quarks must agree with the data on the total hadronic e^+e^- annihilation cross section and on the $\pi^0 \rightarrow 2\gamma$. These requirements make the $SU(3)_C$ be the unique choice. The quanta of $SU(3)_C$ is called gluon. Since the $SU(3)_C$ symmetry is unbroken, the gauge boson, gluon, must be massless. Therefore if A_a^μ denotes the gluon field (a is the color index), ψ_i^α the quark field with flavor index i and color index α , the QCD Lagrangian is

$$\mathcal{L}^{\text{classical}} = -\frac{1}{4}F_{\mu\nu a}F_a^{\mu\nu} + \bar{\psi}^i(i\not{D}_{ij} - m\delta_{ij})\psi^j \quad (5.1)$$

where

$$F_a^{\mu\nu} = \partial^\mu A_a^\nu - \partial^\nu A_a^\mu + gf_{abc}A_b^\mu A_c^\nu, \quad (5.2)$$

$$D^\mu = \partial^\mu - igA_a^\mu T_a. \quad (5.3)$$

Note that last term $gf_{abc}A_b^\mu A_c^\nu$ implies self-interactions of gluons, while there are no such self-interactions in Abelian gauge theory. T_a is the generators of the triplet representation of $SU(3)_C$, following the commutation relations

$$[T_a, T_b] = if_{abc}T_c, \quad (5.4)$$

where f_{abc} are completely antisymmetric structure constants.

In order to quantize the theory one needs a gauge fixing term to be added to eq. (5.1). Usually, the gauge fixing term is chosen as follows:

$$\mathcal{L}^{\text{gauge fixing}} = -\frac{1}{2\alpha}(\partial^\mu A_\mu^a)^2. \quad (5.5)$$

The introduction of such a term requires the addition of the Faddeev-Popov ghost interactions in turn,

$$\mathcal{L}^{\text{FP}} = (\partial^\mu \chi^{a*})D_\mu^{ab}\chi^b, \quad (5.6)$$

where D_μ^{ab} refer to the adjoint representation of $SU(3)_C$. Here we choose a pair of the ghost fields χ^a and χ^{a*} . It is also possible to choose two real fields χ_1^a and χ_2^a instead of χ^a and χ^{a*} . By setting

$$\chi^a = \frac{1}{\sqrt{2}}(\chi_1^a + i\chi_2^a) \quad (5.7)$$

with the Grassmann property

$$(\chi_1^a)^2 = (\chi_2^a)^2 = 0, \quad (\text{no summation on } a), \quad (5.8)$$

we can rewrite the Faddeev-Popov ghost term as

$$\mathcal{L}^{\text{FP}} = i(\partial^\mu \chi_1^a)D_\mu^{ab}\chi_2^b. \quad (5.9)$$

Therefore, we obtain the complete Lagrangian of the theory

$$\begin{aligned}
\mathcal{L} = & -\frac{1}{4}(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)(\partial^\mu A^{a\nu} - \partial^\nu A^{a\mu}) - \frac{1}{2\alpha}(\partial^\mu A_\mu^a)^2 \\
& + i(\partial^\mu \chi_1^a)(\partial_\mu \chi_2^a) + \bar{\psi}^i(i\not{\partial} - m)\psi^i - \frac{g}{2}f^{abc}(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)A^{b\mu}A^{c\nu} \\
& - \frac{g^2}{4}f^{abe}cdeA_\mu^aA_\nu^bA^{c\mu}A^{d\nu} - igf^{abc}(\partial^\mu \chi_1^a)\chi_2^bA_\mu^c + g\bar{\psi}^iT_{ij}^a\gamma^\mu\psi^jA_\mu^a.
\end{aligned} \tag{5.10}$$

Accordingly we obtain Feynman rules for the Lagrangian of quantum chromodynamics (see Appendix B). This theory is renormalizable [26].

We are here using the counterterm approach to realize renormalization again with similar procedure describe in the Section (2.2). We redefine the fields A_μ^a , χ_1^a , χ_2^a and ψ by

$$A_\mu^a = Z_3^{\frac{1}{2}}A_{\mu R}^a, \quad \chi_{1,2}^a = \tilde{Z}_3^{\frac{1}{2}}\chi_{1,2R}^a, \quad \psi = Z_2^{\frac{1}{2}}\psi_R, \tag{5.11}$$

and the parameters g , α and m by

$$g = Z_g g_R, \quad \alpha = Z_3 \alpha_R, \quad m = Z_m m_R, \tag{5.12}$$

where the constants Z_3 , \tilde{Z}_3 and Z_2 denote the gauge field, ghost field and quark field renormalization constants, respectively, while the constants Z_g and Z_m car called the coupling-constant and mass renormalization constants.

Inserting eqs. (5.11) and (5.12) into eq. (5.10), we have

$$\mathcal{L} = \mathcal{L}^R + \mathcal{L}^C \tag{5.13}$$

where the renormalized Lagrangian \mathcal{L}^R is precisely equal to \mathcal{L} if the quantities $\{A_\mu^a, \chi_{1,2}^a, \psi, g, \alpha\}$ are replaced by the renormalized ones, $\{A_{\mu R}^a, \chi_{1,2R}^a, \psi_R, g_R, \alpha_R\}$. The counterterm Lagrangian \mathcal{L}^C is given by

$$\begin{aligned}
\mathcal{L}^C = & (Z_3 - 1) \frac{1}{2} A_R^{a\mu} \delta_{ab} (g_{\mu\nu} \partial^2 - \partial_\mu \partial_\nu) A_R^{b\nu} + (\tilde{Z}_3 - 1) \chi_{1R}^a \delta_{ab} (-i\partial^2) \chi_{2R}^b \\
& + (Z_2 - 1) \bar{\psi}_R^i (i\not{\partial} - m_R) \psi_R^i - Z_2 (Z_m - 1) m_R \bar{\psi}_R^i \psi_R^i \\
& - (Z_1 - 1) \frac{1}{2} g_R f^{abc} (\partial_\mu A_{\nu R}^a - \partial_\nu A_{\mu R}^a) A_R^{\mu b} A_R^{\nu c} \\
& - (Z_4 - 1) \frac{1}{4} g_R^2 f^{abe} f^{cde} A_{\mu R}^a A_{\nu R}^b A_R^{c\mu} A_R^{d\nu} \\
& - (\tilde{Z}_1 - 1) i g_R f^{abc} (\partial^\mu \chi_{1R}^a) \chi_{2R}^b A_{\mu R}^c \\
& + (Z_{1F} - 1) g_R \bar{\psi}^i T_{ij}^a \gamma^\mu \psi_R^j A_{\mu R}^a,
\end{aligned} \tag{5.14}$$

where Z_1, Z_4, \tilde{Z}_1 and Z_{1F} are defined as follows:

$$\begin{aligned}
Z_1 &\equiv Z_g Z_3^{\frac{3}{2}}, & Z_4 &\equiv Z_g^2 Z_3^2, \\
\tilde{Z}_1 &\equiv Z_g \tilde{Z}_3 Z_3^{\frac{1}{2}}, & Z_{1F} &\equiv Z_g Z_2 Z_3^{\frac{1}{2}}.
\end{aligned} \tag{5.15}$$

From this counterterm term we obtain the corresponding Feynman rules (see Appendix B).

The gauge nature of the theory implies the Slavnov-Taylor identity [62, 63],

$$\frac{Z_1}{Z_3} = \frac{\tilde{Z}_1}{\tilde{Z}_3} = \frac{Z_{1F}}{Z_2} = \frac{Z_4}{Z_1}. \tag{5.16}$$

The Slavnov-Taylor identity ensures the universality of the renormalized coupling constant g_R .


By the power counting analysis in the case of QCD, we have seven amplitudes which possess overall divergences. The Feynman diagrams with non-negative superficial degree of divergence in QCD are outlined below:

Note that the superficial degrees of divergence d for the self-energy part for the gluon, Faddeev-Popov ghost and quark and three-gluon vertex are 2, 1, 1 and 1, respectively, but the actual degrees of divergences of these amplitudes are all logarithmic due to the gauge invariance. Next, we present one-loop contributions to the seven superficially divergent amplitudes [64–66]:

(i) The gluon self-energy $\Pi_{\mu\nu}^{ab}(k)$ is

$$\Pi_{\mu\nu}^{ab}(k) = \delta_{ab}(k_\mu k_\nu - k^2 g_{\mu\nu})\Pi(k^2), \quad (5.17)$$

$$\Pi(k^2) = \frac{g_R^2}{(4\pi)^2} \left[\frac{4}{3} T_R N_f - \frac{1}{2} C_G \left(\frac{13}{3} - \alpha_R \right) \right] \frac{1}{\epsilon} + Z_3 - 1 + \text{finite terms}, \quad (5.18)$$



where $\epsilon = (4 - D)/2$. In Eq. (5.18) we have taken N_f flavors of quarks into account, and T_R and C_G are the constants defined by

$$\begin{aligned} \text{tr}[T_a T_b] &= \delta_{ab} T_R, \\ f_{acd} f_{bcd} &= \delta_{ab} C_G. \end{aligned} \quad (5.19)$$

Here we have $T_R = \frac{1}{2}$ and $C_G = 3$ for $SU(3)_C$. Note that the one-loop contribution to the gluon self-energy satisfies the Ward-Takahashi identities,


$$k^\mu \Pi_{\mu\nu}^{ab}(k) = 0, \quad (5.20)$$

which is a natural consequence of gauge invariance. Due to this constraint the amplitude $\Pi_{\mu\nu}^{ab}$ must have the factor $k_\mu k_\nu - k^2 g_{\mu\nu}$ and the degree of divergence for $\Pi_{\mu\nu}^{ab}(k)$ is lowered by 2 units. This structure of $\Pi_{\mu\nu}^{ab}(k)$ forbids a mass terms and so there is no mass renormalization. Therefore the gluon remains massless under the radiative corrections. In the MS scheme [67] the gauge field renormalization constant Z_3 is given by

$$Z_3 = 1 - \frac{g_R^2}{(4\pi)^2} \left[\frac{4}{3} T_R N_f - \frac{1}{2} C_G \left(\frac{13}{3} - \alpha_R \right) \right] \frac{1}{\epsilon} + O(g_R^4). \quad (5.21)$$

(ii) The Faddeev-Popov ghost self-energy $\tilde{\Pi}^{ab}(k)$ is

$$\tilde{\Pi}^{ab}(k) \delta_{ab} \left[- \frac{g_R^2}{(4\pi)^2} C_G \frac{3 - \alpha_R}{4} \frac{1}{\epsilon} + \tilde{Z}_3 - 1 \right] + \text{finite terms}. \quad (5.22)$$




Note that the divergent part above is proportional to k^2 and thus there is no mass renormalization. So the Faddeev-Popov ghost self-energy remain massless after radiative corrections, too. The ghost field renormalization constant \tilde{Z} in the MS scheme is given by

$$\tilde{Z}_3 = 1 + \frac{g_R^2}{(4\pi)^2} C_G \frac{3 - \alpha_R}{4} \frac{1}{\epsilon} + O(g_R^4). \quad (5.23)$$

(iii) The quark self-energy $\Sigma^{ij}(p)$ is

$$\begin{aligned}
\Sigma^{ij}(p) &= \delta_{ij}[(Am_R - B\not{p}) - (Z_2Z_m - 1)m_R + (Z_2 - 1)\not{p}] + \text{finite terms}, \\
A &= -\frac{g_R^2}{(4\pi)^2}C_F(3 + \alpha_R)\frac{1}{\epsilon} + O(g_R^4), \\
B &= -\frac{g_R^2}{(4\pi)^2}C_F\alpha_R\frac{1}{\epsilon} + O(g_R^4).
\end{aligned} \tag{5.24}$$




As we see, the divergence in the quark self-energy consists of two kinds, the mass type Am_R and the kinetic energy type $-B\not{p}$. Then the mass and quark-field renormalization constants in the $\overline{\text{MS}}$ scheme are determined by

$$\begin{aligned}
Z_m &= 1 + A - B + O(g_R^4) \\
&= 1 - \frac{g_R^2}{(4\pi)^2}C_F(3 + \alpha_R)\frac{1}{\epsilon} + \frac{g_R^2}{(4\pi)^2}C_F\alpha_R\frac{1}{\epsilon} + O(g_R^4), \\
Z_2 &= 1 + B + O(g_R^4) \\
&= 1 - \frac{g_R^2}{(4\pi)^2}C_F\alpha_R\frac{1}{\epsilon} + O(g_R^4).
\end{aligned} \tag{5.25}$$

(iv) The three-gluon vertex $\Lambda_{\mu\nu\lambda}^{abc}(K_1, k_2, k_3)$ is given by

$$\Lambda_{\mu\nu\lambda}^{abc}(K_1, k_2, k_3) = -ig_R f^{abc} V_{\mu\nu\lambda}(k_1, k_2, k_3) \left\{ \frac{g_R^2}{(4\pi)^2} \left[C_G \left(-\frac{17}{12} + \frac{3\alpha_R}{4} \right) + \frac{4}{3} T_R N_f \right] \frac{1}{\epsilon} + Z_1 - 1 \right\} + \text{finite terms}, \quad (5.26)$$



where


$$V_{\mu\nu\lambda}(k_1, k_2, k_3) = (k_1 - k_2)_\lambda g_{\mu\nu} + (k_2 - k_3)_\mu g_{\nu\lambda} + (k_3 - k_1)_\nu g_{\mu\lambda}. \quad (5.27)$$

The three-gluon vertex renormalization constant Z_1 in the MS scheme is

$$Z_1 = 1 - \frac{g_R^2}{(4\pi)^2} \left[C_G \left(-\frac{17}{12} + \frac{3\alpha_R}{4} \right) + \frac{4}{3} T_R N_f \right] \frac{1}{\epsilon} + O(g_R^4). \quad (5.28)$$

(v) The ghost-gluon vertex $\tilde{\Lambda}_\mu^{abc}(k, p, p')$ has the express

$$\tilde{\Lambda}_\mu^{abc}(k, p, p') = -ig_R f^{abc} p_\mu \left[\frac{g_R^2}{(4\pi)^2} C_G \frac{\alpha_R}{2} \frac{1}{\epsilon} + \tilde{Z}_1 - 1 \right] + \text{finite terms}, \quad (5.29)$$




where the momentum p_μ denotes the ghost-line which carries the ghost number flowing out of the vertex. The ghost-gluon vertex renormalization constant \tilde{Z}_1 reads in the MS scheme

$$\tilde{Z}_1 = 1 - \frac{g_R^2}{(4\pi)^2} C_G \frac{\alpha_R}{2} \frac{1}{\epsilon} + O(g_R^4). \quad (5.30)$$

(vi) The quark-gluon vertex $\Lambda_{F\mu}^{aij}(k, p, p')$ is

$$\Lambda_{F\mu}^{aij}(k, p, p') = g_R \gamma_\mu T_{ij}^a \left[\frac{g_R^2}{(4\pi)^2} \left(\frac{3 + \alpha_R}{4} C_G + \alpha_R C_F \right) \frac{1}{\epsilon} + Z_{1F} - 1 \right] + \text{finite terms}. \quad (5.31)$$




The quark-gluon vertex renormalization constant Z_{1F} is given by

$$Z_{1F} = 1 - \frac{g_R^2}{(4\pi)^2} \left(\frac{3 + \alpha_R}{4} C_G + \alpha_R C_F \right) \frac{1}{\epsilon} + O(g_R^4). \quad (5.32)$$

(vii) The four-gluon vertex $\Lambda_{\mu_1 \dots \mu_4}^{a_1 \dots a_4}(k_1, k_2, k_3, k_4)$ is

$$\begin{aligned} & \Lambda_{\mu_1 \dots \mu_4}^{a_1 \dots a_4}(k_1, k_2, k_3, k_4) \\ &= -g_R^2 W_{\mu_1 \dots \mu_4}^{a_1 \dots a_4} \left\{ \frac{g_R^2}{(4\pi)^2} \left[\left(-\frac{2}{3} + \alpha_R \right) C_G + \frac{4}{3} T_R N_f \right] \frac{1}{\epsilon} + Z_4 - 1 \right\} \\ &+ \text{finite terms,} \end{aligned} \quad (5.33)$$



where

$$\begin{aligned}
W_{\mu_1 \dots \mu_4}^{a_1 \dots a_4} &= (f^{13,24} - f^{14,32})g_{\mu_1 \mu_2} g_{\mu_3 \mu_4} + (f^{12,34} - f^{14,23})g_{\mu_1 \mu_3} g_{\mu_2 \mu_4} \\
&\quad + (f^{13,42} - f^{12,34})g_{\mu_1 \mu_4} g_{\mu_3 \mu_2}, \\
f^{ij,kl} &\equiv f^{a_i a_j a} f^{a_k a_l a}, \quad i, j, k = 1, 2, 3, 4.
\end{aligned} \tag{5.34}$$

The four-gluon vertex renormalization constant Z_4 in the MS scheme reads

$$Z_4 = 1 - \frac{g_R^2}{(4\pi)^2} \left[\left(-\frac{2}{3} + \alpha_R \right) C_G + \frac{4}{3} T_R N_f \right] \frac{1}{\epsilon} + O(g_R^4). \tag{5.35}$$

Now we find that all the one-loop divergences in the seven superficially divergent amplitudes can be cancelled by the contributions of the counter terms derived from \mathcal{L}^C . Therefore the renormalizability of QCD at the one-loop order is shown.

5.2 Renormalization Group Equation and Asymptotic Freedom

Among renormalizable theories in four spacetime dimensions, non-Abelian gauge theories are unique because of the exclusive possession of asymptotic freedom. It is the significant property that makes QCD such a prominent candidate for the theory of strong interactions in which it gives a substantial basis for incorporating and extending the successful parton model for describing deep inelastic phenomena. In this section, we are dedicated to introduce the renormalization group equations, the concept of running coupling constant, the definition and the physical significance of asymptotic freedom [6–9, 64, 68, 69].

5.2.1 Renormalization Group Equation

According to the renormalization procedure we subtract all the divergences from the Green functions systematically order by order in the perturbative theory. In the subtraction procedure there exists an arbitrariness of defining a divergence part

in a Green function, i.e., how much of the finite part will be subtracted together with the infinity. This arbitrariness is equivalent to that in splitting the Lagrangian into a renormalized Lagrangian and the counterterms and leads to various renormalization schemes.

The arbitrariness remains while defining the renormalized quantities. For example, in QCD, the renormalized coupling constant g_R may be defined in terms either of the three-gluon vertex or of the four-gluon vertex. In general different coupling constants g_R are determined by these different definitions. For QCD, with the help of Slavnov-Taylor identity, these two coupling constants coincide.

In subtracting the singularities we have to introduce an arbitrary mass scale μ which is called the renormalization scale. For instance, in the on-shell scheme, the renormalization scale μ is chosen as the physical mass of the relevant particle at which the renormalization condition is established. In the $\overline{\text{MS}}$ scheme, at first glance, the mass scale seems unnecessary because only the pole in the spacetime dimension is subtracted. However, in fact, the mass dimension of the coupling constant in arbitrary spacetime dimensions plays a role of the renormalization scale. The renormalization scale μ is arbitrary and persists in the finite part of the Green functions. Therefore the renormalized Green functions after subtracting divergences remains arbitrary.

In general, the renormalized coupling constant g_R and mass m_R depend on the renormalization scale μ for which the subtraction procedure is determined, and the explicit dependence can be expressed as

$$\begin{aligned} g_R(\mu) &= Z_g(\mu)^{-1} g, \\ m_R(\mu) &= Z_m(\mu)^{-\frac{1}{2}} m. \end{aligned} \tag{5.36}$$

The renormalized coupling constant $g_R(\mu)$ and $g_R(\mu')$ which are defined via two different subtraction procedures characterized by the renormalization scales μ and μ' respectively. They are related to each other by a finite renormalization $z_g(\mu', \mu)$,

$$g_R(\mu') = z_g(\mu', \mu)g_R(\mu), \quad (5.37)$$

where $z_g(\mu', \mu)$ is defined by

$$z_g(\mu', \mu) = \frac{Z_g(\mu)}{Z_g(\mu')}. \quad (5.38)$$

Similarly, we have

$$m_R(\mu') = z_m(\mu', \mu)m_R(\mu), \quad (5.39)$$

where $z_m(\mu', \mu)$ is defined by

$$z_m(\mu', \mu) = \left(\frac{Z_m(\mu)}{Z_m(\mu')} \right)^{\frac{1}{2}}. \quad (5.40)$$

Note that eq. (5.40) defines a set of finite renormalizations $\{z_g(\mu', \mu)\}$ for varying renormalization scales μ' and μ . We treat the finite renormalization (5.40) as a transformation. It can be shown that this set of transformations have group properties [70].

In fact we could define a product of two elements $z_g(\mu'', \mu')$ and $z_g(\mu', \mu)$

$$z_g(\mu'', \mu')z_g(\mu', \mu), \quad (5.41)$$

which stands for the change of $g_R(\mu)$ through the successive changes of the scales $\mu \rightarrow \mu' \rightarrow \mu''$. Since

$$z_g(\mu'', \mu')z_g(\mu', \mu) = \frac{Z_g(\mu)}{Z_g(\mu'')} = z_g(\mu'', \mu), \quad (5.42)$$

$z_g(\mu'', \mu)$ the finite renormalization of $g_R(\mu)$ caused by the scale change $\mu \rightarrow \mu''$.

Therefore the product $z_g(\mu'', \mu')z_g(\mu', \mu)$ belongs to the set $\{z_g(\mu', \mu)\}$. Furthermore,

the inverse of $z_g(\mu', \mu)$ can be defined by

$$z_g^{-1}(\mu', \mu) = z_g(\mu, \mu'), \quad (5.43)$$

and the identity

$$z_g(\mu, \mu) = 1 \quad (5.44)$$

belongs to the set $\{z_g(\mu', \mu)\}$. Therefore the set of finite renormalizations $\{z_g(\mu', \mu)\}$ is a Abelian group, called the renormalization group.

Furthermore, we define the renormalized one-particle irreducible (1PI) amplitudes by

$$\Gamma_R(p, g_R(\mu'), m_R(\mu'), \mu') = Z_\Gamma \Gamma(p, g_R(\mu), m_R(\mu), \mu) \quad (5.45)$$

where Z_Γ is the product of the necessary scaling factors for the set of operators, depending the number and types of the external lines. For example, in quantum electrodynamics, Γ might be an amputated Green function's with n_e external fermion lines and n_γ external photon lines, and then Z_Γ is given by

$$Z_\Gamma = Z_2^{\frac{n_e}{2}} Z_3^{\frac{n_\gamma}{2}}. \quad (5.46)$$

The finite renormalization for Γ_R is determined by

$$\Gamma_R(p, g_R(\mu'), m_R(\mu'), \mu') = z(\mu', \mu) \Gamma_R(p, g_R(\mu), m_R(\mu), \mu) \quad (5.47)$$

where the renormalization factor $z(\mu', \mu)$ is defined by

$$z(\mu', \mu) = \frac{Z_\Gamma(\mu')}{Z_\Gamma(\mu)}. \quad (5.48)$$

Due to the arbitrariness for choosing the renormalization condition and fixing the renormalization scale μ , we may have many possible expressions for one physical quantity which depends on the choice of the renormalization scheme and scale. These

different expressions are connected by a finite renormalization described above. A natural concern is whether these different expressions for one physical quantity are equivalent or not. Since they represent one physical quantity and are derived from the unique Lagrangian, they describe the same physical phenomenon and therefore must be equivalent. In other words, physical quantities such as renormalized 1PI amplitudes are invariant under finite renormalization.

Given that the choice of renormalization scale is arbitrary, according to the discussion above, we conclude that any change in the renormalization scale μ can be compensated by all the renormalized quantities such that the renormalized 1PI amplitudes remain unchanged. This fact is reflected by the renormalization group equation [67, 71–74].

We can derive the renormalization group equation for the renormalized Green's functions by differentiating eq. (5.45) with respect to μ . Considering g_R and m_R depend on μ , while the unrenormalized amplitude Γ does not, we have immediately that

$$\left[\mu \frac{\partial}{\partial \mu} + \beta(g_R) \frac{\partial}{\partial g_R} - \gamma_m(g_R) m_R \frac{\partial}{\partial m_R} - \gamma_\Gamma(g_R) \right] \Gamma_R = 0, \quad (5.49)$$

where β , γ_m and γ_Γ are defined by

$$\begin{aligned} \beta &= \mu \left. \frac{\partial g_R}{\partial \mu} \right|_{g,m}, \\ \gamma_m &= -\mu \left. \frac{\partial \log m_R}{\partial \mu} \right|_{g,m}, \\ \gamma_\Gamma &= \frac{1}{2} \frac{\partial \log Z_\Gamma}{\partial \mu}. \end{aligned} \quad (5.50)$$

We wish to use the renormalization group equation to study the momentum dependence of the Green function. Assume that all the momentum components vary

together with the fixed ratio, $p = \lambda p_0$, where p_0 is a set of fixed momenta and λ is a momentum scale variable. If Γ has the dimensions of mass to the power D_Γ , then

$$\left[\mu \frac{\partial}{\partial \mu} + m_R \frac{\partial}{\partial m_R} + \lambda \frac{\partial}{\partial \lambda} \right] \Gamma_R = D_\Gamma \Gamma_R, \quad (5.51)$$

so eq. (5.49) can be rewritten as

$$\left\{ \lambda \frac{\partial}{\partial \lambda} - \beta(g_R) \frac{\partial}{\partial g_R} - [1 + \gamma_m(g_R)] m_R \frac{\partial}{\partial m_R} - D_\Gamma + \gamma_\Gamma(g_R) \right\} \Gamma_R(\lambda p_0, g_R, m_R, \mu) = 0. \quad (5.52)$$

Let us define a λ -dependent effective coupling and mass through the differential equations

$$\lambda \frac{d}{d\lambda} g(\lambda) = \beta(g(\lambda)), \quad (5.53)$$

$$\lambda \frac{d}{d\lambda} m(\lambda) = -[1 + \gamma_m(g(\lambda))] m(\lambda) \quad (5.54)$$

and the initial conditions

$$g(1) = g_R, \quad m(1) = m_R. \quad (5.55)$$

Then the eq. (5.52) has the solution

$$\Gamma_R(\lambda p_0, g_R, m_R, \mu) = \lambda^{D_\Gamma} \Gamma_R(p_0, g(\lambda), m(\lambda), \mu) \exp \left[- \int_1^\lambda \gamma_\Gamma(g(\lambda')) \frac{d\lambda'}{\lambda'} \right], \quad (5.56)$$

where the exponential term is the "anomalous dimension". Thus, solution of the renormalization group equation can be expressed in terms of the running coupling constant $g(\lambda)$ and the running mass $m(\lambda)$. The asymptotic behavior of the Green's functions Γ_R is governed the asymptotic behavior of the $g(\lambda)$ and $m(\lambda)$.

According to eq. (5.53), the running coupling constant $g(\lambda)$ must tend to a "fixed point" as $k \rightarrow \infty$, which may be either the point at infinity, or any zeros of the β -function. Thus we need to distinguish three different cases qualitatively: (i) If β at g_R has the same sign as g_R , and if there are no zeros of β between g_R and $\pm\infty$ (for

$g_R > 0$ or $g_R < 0$), then $|g(\lambda)|$ must increase, approaching infinity for $\lambda \rightarrow \infty$. (ii) If β has zeros, and if the first zero encountered as its argument increases for $\beta(g_R) > 0$ or decreases for $\beta(g_R) < 0$ from g_R is as a finite point $g_\infty \neq 0$, then $g(\lambda)$ will increase or decrease to g_∞ as $k \rightarrow \infty$. (iii) If β at g_R has the opposite sign to g_R , and has no zeros between g_R and the origin, then $|g(\lambda)|$ must decrease from $|g_R|$ as λ increases, $|g(\lambda)| \rightarrow 0$ as $\lambda \rightarrow \infty$. Such theories are called "asymptotically free". In the usual case, the perturbation theory gives [6–9],

$$\beta(g_R) = -\beta_0 g_R^3 - \beta_1 g_R^5 - \beta_2 g_R^7 + O(g_R^9). \quad (5.57)$$

Asymptotically free field theories are of great theoretical interests. In such theories, the asymptotic behavior of amplitudes is calculable by the perturbation theory. In the next subsection, we will introduce the renormalization group equation for QCD and asymptotic freedom in QCD.

5.2.2 Asymptotic Freedom in QCD

First, let us derive the renormalization group equation for QCD in the MS scheme. Our basic Lagrangian is given by eq. (5.10) and we split it into two parts, the renormalized part and the counter terms (5.14). We refined the gluon field A_μ^a , ghost field χ^a and quark field ψ through eq. (5.11) and the renormalized parameter g_R , m_R and α_R are defined by eq. (5.12) in terms of renormalization constants Z_g , Z_m and Z_3 . Thus the renormalization group equation for QCD is straightforward and reads off

$$\left[\mu \frac{\partial}{\partial \mu} + \beta(g_R, \alpha_R) \frac{\partial}{\partial g_R} - \gamma_m(g_R, \alpha_R) m_R \frac{\partial}{\partial m_R} + \delta(g_R, \alpha_R) \frac{\partial}{\partial \alpha_R} - n_G \gamma_G(g_R, \alpha_R) - n_F \gamma_F(g_R, \alpha_R) \right] \Gamma_{n_G, n_F} = 0, \quad (5.58)$$

where Γ_{n_G, n_F} is the 1PI renormalized Green's function with n_G external gluon lines and n_F external fermion lines (we do not consider the Green functions with external ghost lines), g_r is the dimensionless renormalized gauge coupling constant defined by

$$g_r = \left(\frac{\mu_0}{\mu} \right)^\epsilon Z_g^{-1} g_0, \quad (5.59)$$

with $g_r = g_R \mu^{-\epsilon}$, $g_0 = g \mu_0^{-\epsilon}$, $\epsilon = \frac{4-D}{2}$, $m_r = m_R$ and $\alpha_r = \alpha_R$. Here the mass scale μ_0 for the bare coupling constant g is fixed scale while the mass scale μ for the renormalized coupling constant g_R is a variable. The renormalization group functions β , γ_m , δ , γ_G and γ_F are defined by

$$\beta(g_r, \alpha_r) = \mu \frac{\partial g_r}{\partial \mu} \Big|_{g, m, \alpha}, \quad (5.60)$$

$$\gamma_m(g_r, \alpha_r) = -\mu \frac{\partial \log m_r}{\partial \mu} \Big|_{g, m, \alpha}, \quad (5.61)$$

$$\delta(g_r, \alpha_r) = \mu \frac{\partial \alpha_r}{\partial \mu} \Big|_{g, m, \alpha}, \quad (5.62)$$

$$\gamma_G(g_r, \alpha_r) = \frac{\mu}{2} \frac{\partial \log Z_3}{\partial \mu} \Big|_{g, m, \alpha}, \quad (5.63)$$

$$\gamma_F(g_r, \alpha_r) = \frac{\mu}{2} \frac{\partial \log Z_2}{\partial \mu} \Big|_{g, m, \alpha}. \quad (5.64)$$

Here γ_G and γ_F are the anomalous dimensions of the gluon and quark fields, respectively. The bare parameters g and m are regarded as fixed constants and are free from the renormalization scale μ . Then we have

$$\frac{dg_R}{d\mu} = 0, \quad \frac{dm}{d\mu} = 0. \quad (5.65)$$

According to eqs. (5.58), (5.59), and (5.64), we have

$$\beta = -\epsilon g_R - \frac{\mu}{Z_g} \frac{dZ_g}{d\mu} g_R. \quad (5.66)$$

Therefore, in order to compute the β -function to one-loop order, we need to know the renormalized coupling constant g_R in one-loop order with the renormalization scale μ . There are four different ways of doing it because Z_g can be evaluated with four different definitions (5.15). These four approaches are equivalent due to the Slavnov-Taylor identity (5.16). We here introduce an easy way of calculating Z_g by using the definition

$$Z_g = \tilde{Z}_1 / (\tilde{Z}_3 Z_3^{\frac{1}{2}}). \quad (5.67)$$

With the help of eqs. (5.21), (5.23) and (5.30), we obtain

$$Z_g = 1 - \frac{g_R^2}{(4\pi)^2} \frac{1}{6} (11C_G - 4T_R N_f) \frac{1}{\epsilon} + O(g_R^4). \quad (5.68)$$

Then, we have, according to eqs. (5.59), (5.66) and (5.68),

$$\begin{aligned} \beta(g_R) &= -\epsilon g_R - \frac{\mu}{Z_g} \frac{dZ_g}{d\mu} g_R \\ &= -\epsilon g_R + \frac{11C_G - 4T_R N_f}{3} \frac{g_R^2}{(4\pi)^2} \frac{1}{\epsilon} \beta(g_R) + O(g_R^5) \\ &= -\frac{1}{(4\pi)^2} \frac{11C_G - 4T_R N_f}{3} g_R^3 + O(g_R^5, \epsilon). \end{aligned} \quad (5.69)$$

Therefore we find that the coefficient β_0 defined in eq. (5.57) is given by

$$\beta_0 = \frac{1}{(4\pi)^2} \frac{11C_G - 4T_R N_f}{3}. \quad (5.70)$$

Asymptotic freedom occurs if $\beta_0 > 0$, i.e., $11C_G - 4T_R N_f > 0$. For $SU(3)$ $C_G = 3$ and $T_R = \frac{1}{2}$, the condition for the asymptotic freedom is

$$N_f < \frac{33}{2}. \quad (5.71)$$

Thus QCD is asymptotically free as long as the number of quark flavors is less than 16. Note that for $N_f = 0$ the coefficient β_0 is positive definite. It is the presence of

quarks that can undermine asymptotic freedom. The fundamental origin of asymptotic freedom may be traced back to the existence of the three-gluon coupling terms in the Lagrangian. Since this term is peculiar to the Yang-Mills theory, we can conclude that the asymptotic freedom is an inherent nature of non-Abelian gauge theory.

So far we discussed the β -function up to one loop order. The β -function up to two loops [75, 76] is given by

$$\beta(g) = -\beta_0 g^3 - \beta_1 g^5 + O(g^7), \quad (5.72)$$

where β_0 is given by eq. (5.70) and

$$\beta_1 = \frac{1}{(4\pi)^4} \left[\frac{34}{3} C_G^2 - 4 \left(\frac{5}{3} C_G + C_F \right) T_R N_f \right]. \quad (5.73)$$

Next, let us turn to the running coupling constant. The running coupling constant $\bar{g}(t)$ at the momentum scale e^t is determined by eq. (5.53), where $t = -\log \lambda$. We choose the momentum scale to be

$$e^t = \frac{\sqrt{-q^2}}{\mu}, \quad (5.74)$$

where q is the space-like momentum and μ is the fixed momentum scale which is chosen to be the renormalization scale for $\bar{g}(0) = g$. Integrating eq. (5.53), we have

$$t = \int_g^{\bar{g}(t)} \frac{dg'}{\beta(g')}. \quad (5.75)$$

Then, we obtain by inserting eq. (5.72) into eq. (5.75)

$$t = -\frac{1}{2} \int_g^{\bar{g}(t)} \frac{d\lambda}{\lambda^2} \frac{dg'}{\beta_0 + \beta_1 \lambda + O(\lambda^2)}. \quad (5.76)$$

If we choose g and λ sufficient small, then we might safely truncate the perturbative series for the β -function to this approximation. Keeping only the one loop order we have

$$t = \frac{1}{2\beta_0} \left(\frac{1}{\bar{g}^2} - \frac{1}{g^2} \right). \quad (5.77)$$

Therefore the running coupling constant \bar{g} is given by

$$\bar{g}^2 = \frac{g^2}{1 + 2\beta_0 g^2 t} = \frac{1}{\beta_0 \log(-q^2/\Lambda^2)}, \quad (5.78)$$

where the new momentum scale Λ is defined by

$$\Lambda = \mu \exp\left[-\frac{1}{2\beta_0 g^2}\right]. \quad (5.79)$$

The momentum scale Λ is referred to as the QCD scale parameter and is the only adjustable parameter in QCD besides the quark mass. The expression for the running coupling constant can be improved by taking into account terms with the coefficient β_1 in eq. (5.76). Performing the integration we have

$$t = \frac{1}{2\beta_0} \left[\frac{1}{\bar{g}^2} - \frac{1}{g^2} + \frac{\beta_1}{\beta_0} \log \frac{\bar{g}^2(\beta_0 + \beta_1 g^2)}{g^2(\beta_0 + \beta_1 \bar{g}^2)} \right]. \quad (5.80)$$

Definin the scale parameter Λ by

$$\Lambda = \mu \exp\left[-\frac{1}{2\beta_0 g^2}\right] \left(\frac{1 + \beta_1 g^2/\beta_0}{\beta_0 g^2}\right)^{\frac{\beta_1}{2\beta_0^2}}, \quad (5.81)$$

we can rewrite eq. (5.80) as follows:

$$\frac{1}{\bar{g}^2} + \frac{\beta_1}{\beta_0} \log \frac{\beta_0 \bar{g}^2}{1 + \beta_1 \bar{g}^2/\beta_0} = \beta_0 \log \left(\frac{-q^2}{\Lambda^2}\right). \quad (5.82)$$

Note that eq. (5.81) reduces to eq. (5.79) for $\beta_1 = 0$. The eq. (5.82) can be solved for \bar{g}^2 iteratively if $-q^2 \gg \Lambda^2$,

$$\bar{g}^2 = \frac{1}{\beta_0 \log(-q^2/\Lambda^2)} \left[1 - \frac{\beta_1}{\beta_0} \frac{\log \log(-q^2/\Lambda^2)}{\log(-q^2/\Lambda^2)} + \dots \right]. \quad (5.83)$$

Note that the second term in the parentheses in the equation above represents the next-to-leading order which corresponds to the two loop correction.

In quantum electrodynamics the coupling constant defined on the mass shell is small enough to ensure the perturbative expansion. However, in quantum chromodynamics, there is no method independent of perturbation theory to determine experimentally the magnitude of the coupling constant. We know nothing about

the validity of perturbation theory in QCD until we perform practical perturbative calculations. Specifically, we first tentatively neglect the question of the validity of perturbation theory and evaluate the β -function in the lowest order of perturbation theory. Then we find that the renormalized coupling constant tends to be small as the relevant momentum scale grows. According to the property of asymptotic freedom, we realize that the perturbative calculation is legitimate for the large momentum scale. Therefore, the perturbation theory in QCD is valid in the large momentum region.

CHAPTER SIX

Yennie-Frautschi-Suura Resummation

The essential idea for understanding the infrared divergences was first proposed by Block and Nordsieck [77] before the invention of relativistic perturbation theory. The idea is that the infrared divergence arises from some soft photons which will escape detection. They showed that the probability that only a finite number of photons will escape detection is precisely zero because of the infrared divergence associated with the soft virtual photons. On the other hand a nonvanishing result would be obtained when the cross section is summed over all possible final states compatible with the detection arrangement. In fact, they proved the cancellation between the real and virtual infrared divergences. As an extension of the idea above, Yennie, Frautschi and Suura (YFS) [78] developed a modern field theoretical treatment of the infrared divergence phenomena. The main feature of this approach is the separation of the infrared divergences as multiplicative factors, which are treated to all orders of perturbation theory, and the conversion of the residual perturbation expansion into one which has no infrared divergence, and hence no need for an infrared cutoff. In the infrared factors, which are in exponential form, the infrared divergences arising from the real and virtual photons cancel out. This procedure depends on no specific details. The beauty of this formalism is that it could be systematically improved order by order in the electromagnetic coupling constant α . In this chapter, we will give a brief introduction to Yennie-Frautschi-Suura theory [78–80]

6.1 Resummation of Virtual Photon Radiative Corrections

Consider a process in which a certain number of photons are generated in the fermion scattering from an initial state of momentum \vec{p} to a final state of momentum \vec{p}' . Let $M_n(\vec{p}, \vec{p}')$ be the contribution to the amplitude corresponding to all n virtual photon loop diagrams. The complete amplitude is then

$$M(\vec{p}, \vec{p}') = \sum_{n=0}^{\infty} M_n(\vec{p}, \vec{p}'). \quad (6.1)$$

Because there are n photons, it is clear that M_n have an infrared divergence of n th order and will be a polynomial of degree of n in the logarithm of the infrared cutoff.

Thus, we could show that the M_n 's have the structure

$$\begin{aligned} M_0 &= m_0, \\ M_1 &= m_0 \alpha B + m_1, \\ M_2 &= m_0 \frac{(\alpha B)^2}{2} + m_1 \alpha B + m_2, \\ &\dots \\ M_n &= \sum_{r=0}^n m_{n-r} \frac{(\alpha B)^r}{r!}, \end{aligned} \quad (6.2)$$

where m_j 's are infrared divergenceless and of order α^j relative to M_0 . Summing over all numbers of virtual photon n , we formally arrive at

$$M(\vec{p}, \vec{p}') = \exp(\alpha B) \sum_{n=0}^{\infty} m_n. \quad (6.3)$$

This is the YFS exponentiation of virtual infrared divergences.

To construct the YFS exponentiation of virtual infrared divergences, we begin with defining that

$$M_n = \frac{1}{n!} \int \cdots \int \prod_{i=1}^n \frac{d^4 k_i}{k_i^2 - m_\gamma^2} \rho_n(k_1, \cdots, k_n), \quad (6.4)$$

where m_γ is the cutoff of the infrared divergence. The factor $\frac{1}{n!}$ arises from the symmetrization of the n virtual photons in ρ_n . Now consider ρ_n is a function of k_n .




Figure 6.1. Basic digrams with any set of real photons and $(n - 1)$ virtual photons

ρ_{n-1} is represented by the set of basic diagrams associated with the first $(n - 1)$ photons and an arbitrary number of potential interactions (see Figure 6.1). ρ_n is represented by the possible ways the n th photon can be inserted into various basic diagrams.

From the Lamb shift analysis, we know that Figure 6.2.(a),(b) and (c) give IR divergences in k_n . The diagrams in Figure 6.2 (d), (e) and (f) are finite as $k_n \rightarrow 0$ if the remaining photon momenta k_i 's are nonzero. As $k_n \rightarrow 0$ and $k_i \rightarrow 0$ simultaneously, overlapping divergences arise and they cancel in gauge invariant combination of terms. Thus, the only remaining divergences corresponds to Figure 6.2 (a), (b) and (c) with $k_n = 0$ in the basic diagram.




Figure 6.2. Basic diagrams with an additional virtual photon.

Thus, we have

$$\rho(k_1, \dots, k_n) = S(k_n)\rho_{n-1}(k_1, \dots, k_{n-1}) + \beta_n^{(1)}(k_1, \dots, k_{n-1}; k_n), \quad (6.5)$$

where $S(k_n)$ contains the k_n infrared contribution from Figure (6.2). The integral of β is infrared divergenceless in k_n .

Iteration of eq. (6.5) gives us

$$\begin{aligned}
\rho_n(k_1, \dots, k_n) &= S(k_n)S(k_{n-1})\rho_{n-2}(k_1 \cdots k_{n-2}) \\
&\quad + S(k_n)\beta_{n-1}^{(1)}(k_1 \cdots k_{n-2}; k_{n-1}) \\
&\quad + S(k_{n-1})\beta_{n-1}^{(1)}(k_1 \cdots k_{n-2}; k_n) \\
&\quad + \{-S(k_{n-1})\beta_{n-1}^{(1)}(k_1 \cdots k_{n-2}; k_n) + \beta_n^1(k_1 \cdots k_{n-1}; k_n)\}.
\end{aligned} \tag{6.6}$$

The symmetry of ρ_n in k_n and k_{n-1} indicates the invariance of bracketed quantity at the end of the above equation under the interchange of k_n and k_{n-1} . So we denote this property by

$$\begin{aligned}
&\{-S(k_{n-1})\beta_{n-1}^{(1)}(k_1 \cdots k_{n-2}; k_n) + \beta_n^1(k_1 \cdots k_{n-1}; k_n)\} \\
&\equiv \beta_n^{(2)}(k_1 \cdots k_{n-2}; k_{n-1}, k_n)
\end{aligned} \tag{6.7}$$

Repeated application of this IR separation procedure and exploitation of the symmetry of ρ_n yield

$$\begin{aligned}
\rho_n(k_1 \cdots k_n) &= S(k_1) \cdots S(k_n)\beta_0 \\
&\quad + \sum_{i=1}^n S(k_1) \cdots S(k_{i-1})S(k_{i+1}) \cdots S(k_n)\beta_1(k_i) \\
&\quad + \cdots \\
&\quad + \sum_{i=1}^n S(k_i)\beta_{n-1}(k_1, k_{i-1}, k_{i+1} \cdots k_n) \\
&\quad + \beta_n(k_1 \cdots k_n).
\end{aligned} \tag{6.8}$$

Rewriting eq. (6.8) in terms of all permutations of k_i and k_j yields

$$\rho_n(k_1 \cdots k_n) = \sum_{Perm} \sum_{r=0}^n \frac{1}{r!(n-r)!} \prod_{i=1}^r S(k_i)\beta_{n-r}(k_{r+1} \cdots k_n). \tag{6.9}$$

Thus, we have

$$M_n = \sum_{r=0}^n \frac{1}{r!(n-r)!} \left(\int \frac{d^4 k S(k)}{k^2 - m_\gamma^2} \right)^r \int \prod_{i=1}^{n-r} \beta_{n-r}(k_1 \cdots k_{n-r}). \tag{6.10}$$

Finally, we define

$$\alpha B(p, p') \equiv \int \frac{d^4 k_i}{k_i^2} \beta_r(k_1 \cdots k_r), \quad (6.11)$$

which yields the desired result, i.e., eq. (6.2).

6.2 Resummation of Real Photon Radiative Corrections

From eq (6.3), we see the cross section is proportional to $\exp(2\alpha\Re B)$. Except that, we need to compute the contribution from the emission of n undetect real photons with total energy ϵ , symmetrized in real photons. The cross section should have the form

$$\begin{aligned} \frac{d\sigma(m_\gamma)}{d\epsilon} &= \sum_{\infty}^{n=0} \frac{d\sigma_n(m_\gamma)}{d\epsilon}, \\ \frac{d\sigma_n(m_\gamma)}{d\epsilon} &= \exp(2\alpha\Re B) \frac{1}{n!} \int \prod_{m=1}^n \frac{d^3 k_m}{\sqrt{k_m^2 + m_\gamma^2}} \\ &\quad \times \delta\left(\epsilon - \sum_{i=1}^n k_i\right) \tilde{\rho}_n(p, p', k_1 \cdots k_n). \end{aligned} \quad (6.12)$$

so that $\tilde{\rho}_n$ plays a similar role to that of ρ_n for the treatment of virtual photons and is given by the absolute square of $\sum m_r$, where

$$E' = E - \sum_{i=1}^{n'} k_i^0 = E - \epsilon.$$

Thus, the sum over all possible undected photons provides the complete differential cross section

$$\frac{d\sigma}{d\epsilon} = \lim_{m_\gamma \rightarrow 0} \sum_{n=0}^{\infty} \frac{d\sigma_n}{d\epsilon}. \quad (6.13)$$

Since $\tilde{\rho}_n$ is symmetric in the real photons and overlapping infrared divergences cancel in the same manner for both real and virtual photons, infrared terms could be factored out of $\tilde{\rho}_n$ by the same treatments applied for ρ_n . Because of the cancellation of overlapping divergences, only the photons which terminate exclusively on external fermion lines (fig.6.3 (a) and (b)) contribute infrared divergence.




Figure 6.3. Basic diagrams with an additional real photon.

We obtain a similar relation to that obtained for the virtual photon case:

$$\begin{aligned}
\tilde{\rho}_n(k_1 \cdots k_n) &= \tilde{S}(k_1) \cdots \tilde{S}(k_n) \tilde{\beta}_0 \\
&+ \sum_{i=1}^n \tilde{S}(k_1) \cdots \tilde{S}(k_{i-1}) \tilde{S}(k_{i+1}) \cdots \tilde{S}(k_n) \tilde{\beta}_1(k_i) \\
&+ \cdots \\
&+ \sum_{i=1}^n \tilde{S}(k_i) \tilde{\beta}_{n-1}(k_1, k_{i-1}, k_{i+1} \cdots k_n) \\
&+ \tilde{\beta}_n(k_1 \cdots k_n). \tag{6.14}
\end{aligned}$$

Similar to the virtual photon case, \tilde{S} contains the infrared divergence and $\tilde{\beta}$ has none. \tilde{S} must be evaluated at $E' = E - \sum k_n^0$, and $\tilde{\beta}_0$ is defined only at $E' = E$.

The energy-conserving δ function eq. (6.12) is conveniently represented by [81,82]

$$\delta\left(\epsilon - \sum_{m=1}^n k_m\right) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \exp\left[iy\left(\epsilon - \sum_{m=1}^n k_m\right)\right]. \tag{6.15}$$

After some manipulations, we obtain

$$\begin{aligned}
\frac{d\sigma}{d\epsilon} &= \lim_{m_\gamma \rightarrow 0} \exp(2\Re B) \frac{1}{2\pi} \int_{-\infty}^{+\infty} dy e^{iy\epsilon} \\
&\times \exp\left[\int^{k \leq \epsilon} \frac{d^3 k}{\sqrt{k^2 + m_\gamma^2}} \tilde{S}(k, p, p') e^{-iyk}\right] \\
&\left\{ \tilde{\beta}_0 + \sum_{n=1}^{\infty} \frac{1}{n!} \int \prod_{m=1}^n \frac{d^3 k_m}{k_m} e^{-iyk} \tilde{\beta}_n(p, p', k_1 \cdots k_n) \right\}. \tag{6.16}
\end{aligned}$$

From the third exponential of eq. (6.16), we see the real infrared photons are still kinematically connected with other real photons by the factor e^{-iyk} which guarantees that $\sum k = \epsilon$. In order to make the infrared photons kinematically independent, we define

$$\int^{k \leq \epsilon} \frac{d^3 k}{(k^2 + m_\gamma^2)^{\frac{1}{2}}} \tilde{S}(k, p, p') e^{-iyk} \equiv 2\alpha \tilde{B} + D, \tag{6.17}$$

where

$$2\alpha \tilde{B}(p, p') \equiv \int^{k \leq \epsilon} \frac{d^3 k}{(k^2 + m_\gamma^2)^{\frac{1}{2}}} \tilde{S} \tag{6.18}$$

and

$$D \equiv \int^{k \leq \epsilon} \frac{d^3 k}{k} \tilde{S}(e^{-iyk} - 1). \quad (6.19)$$

With eq. (6.17) to eq. (6.19), we obtain the noninfrared part of eq. (6.17)

$$\frac{d\hat{\sigma}}{d\epsilon} \equiv \frac{1}{2\pi} \int_{-\infty}^{+\infty} dy e^{iy\epsilon + D} \left\{ \tilde{\beta}_0 + \sum_{n=1}^{\infty} \frac{1}{n!} \int \prod_{m=1}^n \frac{d^3 k_m}{k_m} e^{-iyk_m} \tilde{\beta}_n \right\}. \quad (6.20)$$

Then, eq (6.17) becomes

$$\frac{d\sigma}{d\epsilon} = \exp \left\{ \lim_{m_\gamma \rightarrow 0} 2\alpha(B + \tilde{B}) \right\} \frac{d\hat{\sigma}}{d\epsilon}. \quad (6.21)$$

So far, the problem is whether or not $\lim_{m_\gamma \rightarrow 0} 2\alpha(B + \tilde{B})$ is finite or not. Next, we will show the cancellation of infrared terms to all orders of the electromagnetic coupling α by exploiting details of infrared factors.

6.3 Details of Infrared Factors

B and \tilde{B} can be represented by the gauge invariant expressions [82]

$$B = \frac{i}{(2\pi)^3} \int \frac{d^4 k}{k^2 - m_\gamma^2} \left(\frac{2p'_\mu - k_\mu}{2p' \cdot k - k^2} - \frac{2p_\mu - k_\mu}{2p \cdot k - k^2} \right)^2 \quad (6.22)$$

and

$$B = \frac{-1}{8\pi^2} \int_0^\epsilon \frac{d^4 k}{(k^2 + m_\gamma^2)^{\frac{1}{2}}} \left(\frac{p'_\mu}{p' \cdot k} - \frac{p_\mu}{p \cdot k} \right)^2 \quad (6.23)$$

We can see that the infrared divergent part of $\Re B$ arises from the pole

$$\frac{1}{k^2 - m_\gamma^2 + i\epsilon} = P.V. \frac{1}{k^2 - m_\gamma^2} - i\pi\delta(k^2 - m_\gamma^2) \quad (6.24)$$

These poles contribute the amount

$$B = \frac{1}{8\pi^2} \int \frac{d^4 k}{(k^2 + m_\gamma^2)^{\frac{1}{2}}} \left(\frac{2p'_\mu - k_\mu}{2p' \cdot k - k^2} - \frac{2p_\mu - k_\mu}{2p \cdot k - k^2} \right)^2 + \text{finite terms}. \quad (6.25)$$

As $k \rightarrow 0$, the diverging integrands of eq. (6.25) and \tilde{B} cancel.

Thus, $\lim_{m_\gamma \rightarrow 0} (2\alpha\Re B(m_\gamma) + 2\alpha\tilde{B}(m_\gamma))$ is finite, i.e., we have cancelled the infrared divergence in the theory to all order in α .

After exact calculations, we have [79, 80]

$$\begin{aligned}
\lim_{m_\gamma \rightarrow 0} (2\alpha \Re B + 2\alpha \tilde{B}) = & \frac{\alpha}{\pi} \left\{ \left(\log \frac{pp'}{m^2} - 1 \right) \log \frac{k_m^2}{EE'} + \frac{1}{2} \log \frac{2pp'}{m^2} - \frac{1}{2} \log^2 \frac{p_0}{p'_0} \right. \\
& - \frac{1}{4} \log^2 \frac{(\Delta + \delta)^2}{4p^0 p'^0} - \frac{1}{4} \log^2 \frac{(\Delta - \delta)^2}{4p^0 p'^0} - \Re Li_2 \left(\frac{\Delta + \omega}{\Delta + \delta} \right) \\
& - \Re Li_2 \left(\frac{\Delta + \omega}{\Delta - \delta} \right) - \Re Li_2 \left(\frac{\Delta - \omega}{\Delta + \delta} \right) - \Re Li_2 \left(\frac{\Delta - \omega}{\Delta - \delta} \right) \\
& \left. + \frac{\pi^2}{3} - 1 \right\}, \tag{6.26}
\end{aligned}$$

where $k_m = \epsilon$,

$$\Delta = \sqrt{2pp' + (p^0 - p'^0)},$$

$$\omega = p^0 + p'^0,$$

$$\delta = p^0 - p'^0,$$

and we have introduced the Spence function

$$Li_2(x) = - \int_0^x \log(1-t) dt.$$

At high energies and small ϵ , B and \tilde{B} have the approximate forms

$$B = -\frac{1}{2\pi} \left[\log \frac{2p \cdot p'}{m^2} \left(\log \frac{m^2}{m_\gamma^2} + \frac{1}{2} \log \frac{2p \cdot p'}{m^2} - \frac{1}{2} \right) - \log \frac{m^2}{m_\gamma^2} \right], \tag{6.27}$$

$$\begin{aligned}
\tilde{B} = & \frac{1}{2\pi} \left[\log \frac{2p \cdot p'}{m^2} \left(\log \frac{m^2}{m_\gamma^2} + \frac{1}{2} \log \frac{2p \cdot p'}{m^2} - \log \frac{EE'}{\epsilon^2} \right) - \log \frac{m^2}{m_\gamma^2} + \log \frac{EE'}{\epsilon^2} \right]. \tag{6.28}
\end{aligned}$$

If we use a photon momentum k_{min} instead of the photon mass m_γ , eqs. (6.27) and

(6.28) become

$$B = -\frac{1}{2\pi} \left[\log \frac{2p \cdot p'}{m^2} \left(\log \frac{EE'}{k_{min}^2} - \frac{1}{2} \right) - \log \frac{EE'}{k_{min}^2} \right] \tag{6.29}$$

and

$$\tilde{B} = \frac{1}{2\pi} \left(\log \frac{2p \cdot p'}{m^2} - 1 \right) \log \frac{\epsilon^2}{k_{min}^2}. \tag{6.30}$$

However, the sum $\Re B(m_\gamma) + \tilde{B}(m_\gamma)$ is the same as $\Re B(k_{min}) + \tilde{B}(k_{min})$:

$$2\alpha(\Re B + \tilde{B}) = -\frac{\alpha A}{2} \log \frac{EE'}{\epsilon^2} + \frac{\alpha}{2\pi} \log \frac{2p \cdot p'}{m^2}, \quad (6.31)$$

where

$$\begin{aligned} \alpha A &\equiv -\frac{k^2 \alpha}{4\pi^2} \int d\Omega \left(\frac{p'_\mu}{p' \cdot k} - \frac{p_\mu}{p \cdot k} \right)^2 \\ &\cong \frac{2\alpha}{\pi} \left(\log \frac{2p \cdot p'}{m^2} - 1 \right). \end{aligned} \quad (6.32)$$

It shows the physical consequence is independent of the selection of regularization schemes.

6.4 Details of Noninfrared Virtual Photon Terms

We now discuss the virtual photon remainders (6.2)

$$\begin{aligned} m_1 &= M_1 - \alpha B M_0, \\ m_2 &= M_2 - \alpha B M_1 + \frac{(\alpha B)^2}{2!} M_0 \\ &\dots\dots \end{aligned} \quad (6.33)$$

Note that since αB (and $\alpha \tilde{B}$) are not unique, the separation of M_1 into $\alpha B M_0$ and m_1 is not unique. Recoil terms such as k^2 in $(k^2 - 2kp)^{-1}$ does not affect the infrared singularity, but are preserved in B to make the integral (6.22) converge naturally as $k \rightarrow \infty$. Recoil currents such as k_μ in $(2p_\mu - k_\mu)$ do not contribute to the infrared divergence, but they are remained in the integral (6.22) to make B gauge invariant. Thus, different representations of B with these terms having different coefficients would yield the same infrared singularities. In general, m_1 could be very complicated, however, we could still obtain some good results through discussing the lowest order for an fermion.

Applying the Feynman rules for the amplitude above, we could represent the incoming part as

$$\dots \frac{(p - k_i + m)\epsilon_i}{k_i^2 - 2p \cdot k_i} u(p) = \dots \frac{(2p - k_i) \cdot \epsilon_i - \frac{1}{2}[k_i, \epsilon_i]}{k_i^2 - 2K_i \cdot p} u(p) \quad (6.34)$$

where the first term on the right is the current we have used in the factor B , and the second term is the magnetic term.




Figure 6.4. Vacuum polarization of the potential.

Applying the Feynman rules for the amplitude above, we could represent the incoming part as

$$\dots \frac{(p - k_i + m)\epsilon_i}{k_i^2 - 2p \cdot k_i} u(p) = \dots \frac{(2p - k_i) \cdot \epsilon_i - \frac{1}{2}[k_i, \epsilon_i]}{k_i^2 - 2K_i \cdot p} u(p) \quad (6.35)$$

where the first term on the right is the current we have used in the factor B , and the second term is the magnetic term.

The calculation of the magnetic terms at the high energy limit gives the contribution

$$\frac{\alpha M_0}{2\pi} \log \frac{2p \cdot p'}{m^2} + O(\alpha M_0) \quad (6.36)$$

And the vacuum polarization contributes

$$\frac{\alpha M_0}{3\pi} \log \frac{2p \cdot p'}{m^2} + O(\alpha M_0). \quad (6.37)$$

6.5 Details of Noninfrared Real Photon Terms

We have discussed the infrared terms B and \tilde{B} and the virtual corrections in $\tilde{\beta}_n$, we are focusing on the noninfrared real photon corrections in eq. (6.20). The photons have a spectrum of dk rather than dk/k for $k \rightarrow 0$, and the expansion of the real photon correction in n is an expansion in the number of noninfrared real photons. Thus, the n th order correction is from noninfrared photons. We assume the energy loss for emitting one real photon is ϵ . The $n = 0$ term contains dk/k contribution from the emitted photon. And the $n = 1$ term contains dk/k and dk contributions from one photon. It means the $n = 0$ and $n = 1$ terms both start at the order of $O(\alpha)$.

We start with the $n = 0$ case in eq. (6.20):

$$\begin{aligned}\frac{d\hat{\sigma}_0}{d\epsilon} &= \tilde{\beta}_0 I, \\ \frac{d\sigma_0}{d\epsilon} &= \frac{d\hat{\sigma}_0}{d\epsilon} e^{2\alpha(\Re B + \tilde{B}(\epsilon))},\end{aligned}\tag{6.38}$$

where

$$\begin{aligned}I &\equiv \frac{1}{2\pi} \int_{-\infty}^{+\infty} dy e^{iy\epsilon + D}, \\ D &= \int^{k^0 \geq \epsilon} \frac{dk}{k} \tilde{S}(p, p', k) (e^{-iyk^0 - 1}), \\ \tilde{S}(p, p', k) &= -\frac{\alpha Q_f^2}{4\pi^2} \left(\frac{p_\mu}{k \cdot p} - \frac{p'_\mu}{k \cdot p'} \right).\end{aligned}\tag{6.39}$$

I could be computed in terms of tabulated functions [82]. The result is

$$I = \frac{\alpha A}{\epsilon} F_{YFS}(\alpha A),\tag{6.40}$$

where

$$\begin{aligned} F_{YFS}(\alpha A) &= \frac{e^{-\alpha AC}}{\Gamma(1 + \alpha A)} \\ &= 1 - \frac{\pi^2(\alpha A)^2}{12} + \dots, \end{aligned} \quad (6.41)$$

Therefore we have

$$\frac{d\sigma_0}{d\epsilon} = e^{2\alpha(\Re B + \tilde{B}(\epsilon))} \frac{\alpha A}{\epsilon} F_{YFS}(\alpha A) \tilde{\beta}_0, \quad (6.42)$$

where

$$2\alpha(\Re B + \tilde{B}(\epsilon)) = \alpha A \log \frac{\epsilon}{E} + \frac{\pi}{\alpha} \left(\frac{1}{2} \log \frac{pp'}{m^2} - 1 - \frac{\pi^2}{6} \right). \quad (6.43)$$

Next, for the $n = 1$ case, we have

$$\begin{aligned} \frac{d\sigma_1}{d\epsilon} &= e^{2\alpha(\Re B + \tilde{B}(\epsilon))} \frac{d\hat{\sigma}_1}{d\epsilon}, \\ \frac{d\hat{\sigma}_1}{d\epsilon} &= \int^{k_1 \leq \epsilon} \frac{d^3 k_1}{k_1} \tilde{\beta}_1(k_1) \frac{1}{\pi} \int_{-\infty}^{+\infty} dy e^{iy(\epsilon - k_1) + D}, \end{aligned} \quad (6.44)$$

where the $\tilde{\beta}_1$ contains dk and kdk terms from the emission of one real photon. And $\tilde{\beta}_1(k_1)$ is evaluated at $E' = E - k_1$

After the similar calculation for the $n = 0$ case, we arrive at

$$\frac{d\sigma_1}{d\epsilon} = \frac{\alpha A}{\epsilon} F(\alpha A) \int_0^\epsilon dk_1 G_1(k_1) \left(\frac{\epsilon}{\epsilon - k_1} \right)^{1 - \alpha A}. \quad (6.45)$$

We can show that the above integral is convergent for $\alpha A \geq 0$. Because of the peaking behavior at $k_1 = \epsilon$, we could expand $G_1(k_1)$ about $k_1 = \epsilon$:

$$G_1(k_1) = G_1(\epsilon) + (k_1 - \epsilon) \left. \frac{dG_1(k_1)}{dk_1} \right|_{k_1=\epsilon} + \dots, \quad (6.46)$$

so that we have

$$\frac{d\hat{\sigma}_1}{d\epsilon} = F_{YFS}(\alpha A) \left\{ G_1(\epsilon) - \frac{\alpha A \epsilon}{\alpha A + 1} \frac{dG_1(k_1 = \epsilon)}{dk_1} \right\}. \quad (6.47)$$

The $G_1(\epsilon)$ term is of order α from a hard photon. And the other terms in $\{\dots\}$ is from the infrared photons in addition to the "dk" photon in G_1

Therefore, we have [80]

$$\begin{aligned} \frac{d\sigma_1}{d\epsilon} = & \exp \left[\alpha A \log \frac{\epsilon}{E} + \frac{\pi}{\alpha} \left(\frac{1}{2} \log \frac{pp'}{m^2} - 1 - \frac{\pi^2}{6} \right) \right] \\ & \times F_{YFS} \left\{ G_1(\epsilon) - \frac{\alpha A \epsilon}{1 + \alpha A} G_1'(\epsilon) + \dots \right\}. \end{aligned} \quad (6.48)$$

In some literatures, we set $\epsilon = v \frac{\sqrt{s}}{2} = vE$ so that

$$\frac{d\sigma_1}{dv} = v^{\alpha A} F_{YFS}(\alpha A) e^{\frac{\alpha}{\pi} \left(\frac{1}{2} \log \frac{pp'}{m^2} - 1 - \frac{\pi^2}{6} \right)} G_1(v) + O(\alpha^2) \quad (6.49)$$

which is useful for many applications in precision EW, QCD and quantum gravity [83].

CHAPTER SEVEN

CEEX and KKMC-hh

In last chapter, we have given a brief introduction to Yenni-Frautschi-Suura(YFS) theory. The YFS theory provides an elegant treatment for infrared singularities to all orders of the electromagnetic coupling α . Rooted from YFS framework on QED exponentiation [78], many important Monte Carlo (MC) event generators were developed in pursuit of ultimate precision in theoretical particle theory, such as KORALB [84], KORALZ [85], BHLUMI [86, 87], YFS2 [88], KKMC [89], KKMC-hh [90–92] and etc.

KKMC-hh is an extension of the generator KKMC for the process $e^+e^- \rightarrow f\bar{f} + n\gamma$ in LEP energies, where $f\bar{f}$ represents a final state fermion pair. The MC structure is based on CEEX [89, 94–96], an amplitude based analog to the original YFS exponentiation, and includes residuals through the order of $\alpha^2 L$, where $L \equiv \log(\frac{s}{m_f^2})$. (big logarithm). Electroweak matrix element corrections are computed by DIZET 6.21 from the program ZFITTER [97]. ZFITTER calculates vacuum polarization for the photon and Z boson, and adds the form factor corrections to the vector coupling and angle-dependent form factor to includes the box diagram corrections. The decay is realized by TAUOLA [98–100].

The KKMC of version 4.22 supports quark initial states, and a modified version 4.22 is incorporated into KKMC-hh to select the quarks via PDF's with the help of an LHAPDF [101] interface. KKMC-hh utilizes an adaptive MC program FOAM [102] to generate the quark momentum fractions x_i , the total ISR energy, and the quark flavor using a crude distribution which is constructed during an initialization phase.

In this chapter, we will introduce the two types of QED matrix elements and exponentiations: the coherent exclusive exponentiation(CEEX) and exclusive exponentiation (EEX) at first. Then we would like to give a brief review the MC algorithms for KKMC.

7.1 Amplitudes for Exclusive Exponentiation

In this context, exclusivity means that the procedure of exponentiation(summing up the infrared real and virtual contributions within the scheme of perturbative quantum field theory) is done at the level of fully differential (multiphoton) cross section or at the level of the scattering matrix element (spin amplitudes) before integrating over photon momenta in the phase space. As opposite to exclusivity, inclusivity represents executing the procedure of exponentiation after phase space integration over photon momenta. EEX is formulated in terms of spin summed or averaged differential distributions, which results both advantages and disadvantages. The advantage of EEX formulation is that the differential distributions are given analytically in terms of Mandelstam variables and they are easily examined by checking certain important limit, such as leading-logarithmic and soft limits. However, the disadvantage is that the squaring of the sums of spin amplitudes from Feynman diagrams leads to many interference terms, which in the exponentiation are calculated analytically and individually. In spite of disadvantages, the EEX matrix element still play an important role to provide a testing environment for the new , more complicated matrix element of the CEEX class. In this section, we will give a concise introduction of amplitudes for exclusive exponentiation. We use the process $e^-e^+ \rightarrow f\bar{f} + n\gamma$ to illustrate the


EEX, which could be extended to partonic process $q\bar{q} \rightarrow f\bar{f} + n\gamma$ in KKMC-hh ($q \equiv \text{quark}$).

7.1.1 Master Formula

The kinematics of the process $e^-e^+ \rightarrow f\bar{f} + n\gamma$ is described in the figure below. In this case, we neglect the initial-final state interference. Therefore, we are allowed to distinguish between photons radiated from the initial-state fermions and those radiated from the final-state fermions. The four-momentum

$$X = p_1 + p_2 - \sum_{j=1}^n k_j = q_1 + q_2 + \sum_{l=1}^{n'} k'_l \quad (7.1)$$

of the s-channel virtual boson (Z/γ^*) is well defined. Let us denote the rest frame of X as XMS (the X zero momentum system).



The kinematics with multiple photon emissions in the process $e^+e^- \rightarrow f\bar{f} + n\gamma$

Denoting the Lorentz-invariant phase-space by

$$d^n \text{Lips}(P; p_1, p_2, \dots, p_n) = \prod_{j=1}^n \frac{d^3 p_j}{p_j^0} \delta^{(4)}\left(P - \sum_{j=1}^n p_j\right) \quad (7.2)$$

for the process $e^-(p_1) + e^+(p_2) \rightarrow f(q_1) + \bar{f}(q_2) + n\gamma(k_j) + n'\gamma(k'_l)$, the $O(\alpha^r)$ total cross section reads

$$\begin{aligned} \sigma_{EEEX}^{(r)} &= \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \frac{1}{n!} \frac{1}{n'!} \iint d^{n+n'+2} \text{Lips}(p_1 + p_2; q_1, q_2, k_1 \dots, k_n, k'_1 \dots, k'_{n'}) \\ &\rho_{EEEX}^{(r)}, r = 0, 1, 2, 3, \end{aligned} \quad (7.3)$$

in terms of the fully differential multiphoton distribution

$$\begin{aligned} &\rho_{EEEX}^{(r)}(p_1, p_2; q_1, q_2, k_1 \dots, k_n, k'_1 \dots, k'_{n'}) \\ &= \exp[Y_e(\Sigma_I; p_1, p_2) + Y_f(\Sigma_F; q_1, q_2)] \prod_{j=1}^n \tilde{S}_I(k_j) \Theta(\Sigma_I^-; k_j) \tilde{S}_F(k'_l) \Theta(\Sigma_F^-; k'_l) \\ &\left\{ \bar{\beta}_0^{(r)}(X, p_1, p_2, q_1, q_2) + \sum_{j=1}^n \frac{\bar{\beta}_1^{(r)} I(X, p_1, p_2, q_1, q_2, k_j)}{\tilde{S}_I(k_j)} + \right. \\ &\sum_{l=1}^{n'} \frac{\bar{\beta}_1^{(r)} F(X, p_1, p_2, q_1, q_2, k'_l)}{\tilde{S}_F(k'_l)} + \sum_{n \geq j > k \geq 1} \frac{\bar{\beta}^{(r)2II}(X, p_1, p_2, q_1, q_2, k_j, k_k)}{\tilde{S}_I(k_j) \tilde{S}_I(k_k)} \\ &+ \sum_{n' \geq l > m \geq 1} \frac{\bar{\beta}^{(r)2FF}(X, p_1, p_2, q_1, q_2, k'_l, k'_m)}{\tilde{S}_F(k'_l) \tilde{S}_F(k'_m)} \\ &+ \sum_{j=1}^n \sum_{l=1}^{n'} \frac{\bar{\beta}^{(r)2IF}(X, p_1, p_2, q_1, q_2, k_j, k'_l)}{\tilde{S}_I(k_j) \tilde{S}_F(k'_l)} \\ &\left. + \sum_{n \geq j > k > l \geq 1} \frac{\bar{\beta}_{3III}^{(r)}(X, p_1, p_2, q_1, q_2, k_j, k_k, k_l)}{\tilde{S}_I(k_j) \tilde{S}_I(k_k) \tilde{S}_I(k_l)} \right\}. \end{aligned} \quad (7.4)$$

As we noted in last chapter, the YFS soft factors for real photons emitted from the initial and final state fermions are

$$\begin{aligned} \tilde{S}_I(k_j) &= -Q_e^2 \frac{\alpha}{4\pi^2} \left(\frac{p_1}{k_j p_1} - \frac{p_2}{k_j p_2} \right)^2, \\ \tilde{S}_F(k'_l) &= -Q_f^2 \frac{\alpha}{4\pi^2} \left(\frac{q_1}{k'_l q_1} - \frac{q_2}{k'_l q_2} \right)^2, \end{aligned} \quad (7.5)$$

where Q_e and Q_f are the electric charges of the electron e and fermion f . The Y function in the exponential YFS from factor is defined as in last chapter:

$$\begin{aligned}
Y_f(\Omega, p, \bar{p}) &\equiv 2Q_f^2\alpha\tilde{B}(\Omega, p, \bar{p}) + 2Q_f^2\alpha\Re B(\Omega, p, \bar{p}) \\
&\equiv -2Q_f^2\frac{\alpha}{8\pi^2}\int\frac{d^k}{k^0}\Theta(\Omega; k)\left(\frac{p}{kp}-\frac{\bar{p}}{k\bar{p}}\right)^2 \\
&\quad + 2Q_f^2\alpha\Re\int\frac{d^4k}{k^2}\frac{i}{2\pi^3}\left(\frac{2p-k}{kp-k^2}-\frac{2\bar{p}-k}{2k\bar{p}-k^2}\right)^2. \tag{7.6}
\end{aligned}$$

The form factor above is IR convergent and depends explicitly on the soft-photon domains $\Omega = \Omega_I, \Omega_F$, which includes the IR divergence point $k = 0$. We define $\Theta(\Omega; k) = 1$ for $k \in \Omega$ and $\Theta(\Omega; k) = 0$ for $k \in \bar{\Omega}$. The sum over contributions from the real photons inside the domain Ω to infinite order, together with the analogous contributions from virtual photons, forms the exponential YFS form factor. In the Monte Carlo calculation we generate photons $k \in \bar{\Omega}$ via function $\Theta(\bar{\Omega}, k) = 1 - \Theta(\Omega, k)$. Usually it is required that Ω_I and Ω_F are small enough in the total cross section. And physical observables are independent of the choice of Ω_I and Ω_F . so mathematically speaking, $\Omega_{I/F}$ is required to be dummy parameters in the calculation. If we neglect the initial final state inference, we could choose Ω_I and Ω_F differently. For example, let us define Ω_I with $k^0 < E_{min}$ in the center of mass system of the incoming e^+e^- beams and Ω with $k'^0 < E'_{min}$ in the center of mass system of the outgoing fermions $f\bar{f}$. This is the easiest definition for the Monte Carlo Generation, but in the latter discussion, we would should how we deal with the situation $\Omega_I = \Omega_F$. The YFS form factor for the above choices of $\Omega_{I/F}$ are

$$Y_e(\Omega_I; p_1, p_2) = \gamma_e \log \frac{2E_{min}}{\sqrt{2p_1p_2}} + \frac{1}{4}\gamma_e + Q_e^2\frac{\alpha}{\pi}\left(-\frac{1}{2} + \frac{\pi^2}{3}\right), \tag{7.7}$$

$$Y_f(\Omega_F; p_1, p_2) = \gamma_f \log \frac{2E_{min}}{\sqrt{2p_1p_2}} + \frac{1}{4}\gamma_f + Q_e^2\frac{\alpha}{\pi}\left(-\frac{1}{2} + \frac{\pi^2}{3}\right), \tag{7.8}$$

where

$$\gamma_e = 2Q_e^2 \frac{\alpha}{\pi} \left(\log \frac{2p_1 p_2}{m_e^2} - 1 \right), \quad (7.9)$$

$$\gamma_f = 2Q_f^2 \frac{\alpha}{\pi} \left(\log \frac{2q_1 q_2}{m_f^2} - 1 \right), \quad (7.10)$$

7.1.2 Pure Virtual Corrections

As we discussed in the last chapter, the perturbative QED matrix element is located in the $\bar{\beta}$ functions. The $\bar{\beta}_0$ function is proportional to the Born differential cross section $d\sigma^{Born}(s, \theta)/d\Omega$ for the process $e^+e^- \rightarrow f\bar{f}$ and it contains calculable infrared convergent corrections order by order. We shall calculate $\bar{\beta}_0$ and other $\bar{\beta}$'s in the $O(\alpha^j)_{prag}$, $i = 0, 1, 2$.

The $O(\alpha^j)_{prag}$ expressions for $\bar{\beta}_0^{(i)}$, $i = 0, 1, 2$ are

$$\bar{\beta}_0^{(r)}(X, p_1, p_2, q_1, q_2) = \frac{1}{4} \sum_{k,l=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{kl})(1 + \delta_I^{(r)})(1 + \delta_F^{(r)}), \quad (7.11)$$

$$\delta_I^{(0)} = 0, \quad \delta_I^{(1)} = \frac{1}{2}\gamma, \quad \delta_I^{(2)} = \delta_I^{(1)} + \frac{1}{8}\gamma^2, \quad \delta_I^{(3)} = \delta_I^{(2)} + \frac{1}{48}\gamma^3, \quad (7.12)$$

$$\delta_F^{(0)} = 0, \quad \delta_F^{(1)} = \frac{1}{2}\gamma_f, \quad \delta_F^{(2)} = \delta_F^{(1)} + \frac{1}{8}\gamma_f^2, \quad \delta_F^{(3)} = \delta_F^{(2)} + \frac{1}{48}\gamma_f^3, \quad (7.13)$$

where

$$\begin{aligned} \cos \theta_{11} &= \frac{\vec{p}_1 \cdot \vec{q}_1}{|\vec{p}_1| |\vec{q}_1|}, & \cos \theta_{12} &= \frac{-\vec{p}_1 \cdot \vec{q}_2}{|\vec{p}_1| |\vec{q}_2|}, \\ \cos \theta_{21} &= \frac{-\vec{p}_2 \cdot \vec{q}_1}{|\vec{p}_2| |\vec{q}_1|}, & \cos \theta_{22} &= \frac{\vec{p}_2 \cdot \vec{q}_2}{|\vec{p}_2| |\vec{q}_2|}, \end{aligned} \quad (7.14)$$

with all three-vectors in the rest frame of the four momentum X , namely, in the frame XMS .

Notice that we take an average over four θ_{kl} instead of having a single $\frac{d\sigma^{\text{Born}}}{d\Omega(\theta)}$. The reason for this more complex choice is due to the treatment of the first- and higher-order real photon contributions in the next subsections. According to the Refs. [103, 104], the exact single-photon ISR/FSR matrix element can be expressed as a linear combination of the two $d\sigma^{\text{Born}}/d\Omega(\theta_k), k = 1, 2$, distributions. Our implementation of the LL matrix element of the two or three real photons could also apply this kind of linear combinations. Thus, it is practical and reasonable to adopt a similar approach ahead for $\bar{\beta}_0$. Note that in the soft limit, all four angles θ_{kl} are identical.

Some reader might question the authenticity of the freedom of defining θ in $d\sigma^{\text{Born}}/d\Omega(\theta)$ in the first place. This question is answered in Ref. [87, 88]. Briefly speaking, the differential cross section $d\sigma^{\text{Born}}/d\Omega(\theta)$ and $\bar{\beta}_0^{(i)}$ are defined in the two-body phase-space. However, they will be used all over the phase-space with additional photons(either soft or hard). This needs some extrapolations of $d\sigma^{\text{Born}}/d\Omega(\theta)$ and $\bar{\beta}_0^{(i)}$ beyond the two-body phase-space. The extrapolation is realized by manipulating the four-momenta in Ref. [88] and it is done as an extrapolation for the Mandelstam variables(s, t and u). However, we could solve this problem from another perspective. The effect due to changing from one specific extrapolation to another is a kind of "higher order" effect for the entire calculation. For example, at $O(\alpha^1)$, changing the type of extrapolation brings in an $O(\alpha^2)$ effect. Therefore, it is logical to use a certain extrapolation to minimize the higher-order effects.

7.1.3 One Real Photon with Virtual Corrections

The contributions $\bar{\beta}_1^{(2)}$ are built from the QED distributions with a single real-photon emission and up to one virtual-photon contribution. They are defined as follows:

$$\bar{\beta}_{1I}^{(i)}(X, p_1, p_2, q_1, q_2, k_j) = D_{1I}^{(i)}(X, p_1, p_2, q_1, q_2, k_j) - \tilde{S}_I(k_j) \bar{\beta}_0^{(i-1)}(X, p_1, p_2, q_1, q_2), \quad (7.15)$$

$$\bar{\beta}_{1F}^{(i)}(X, p_1, p_2, q_1, q_2, k'_i) = D_{1F}^{(i)}(X, p_1, p_2, q_1, q_2, k'_i) - \tilde{S}_F(k'_i) \bar{\beta}_0^{(i-1)}(X, p_1, p_2, q_1, q_2), \quad (7.16)$$

where $i = 1, 2$. We define all the ingredients for the initial-state contribution. The single initial-state photon emission differential distribution at the $O(\alpha^r)$, $r = 1, 2, 3$ with up to two-loop virtual correction from the initial- and/or final-state photon is

$$\begin{aligned} D_{1I}^{(r)}(X, p_1, p_2, q_1, q_2, k_j) &= Q_e^2 \frac{\alpha}{4\pi^2} \frac{2p_1 p_2}{(k_j p_1)(k_j p_2)} W_e(\hat{\alpha}_j, \hat{\beta}_j) \\ &\times \left\{ \frac{(1 - \hat{\alpha}_j)^2}{2} \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) + \frac{(1 - \hat{\beta}_j)^2}{2} \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{2r}) \right\} \\ &\times \left[1 + \Delta_I^{(r-1)}(z_j) \right] (1 + \delta_F^{(r-1)}), \end{aligned} \quad (7.17)$$

where

$$\begin{aligned} \hat{\alpha}_j &= \frac{k_j p_2}{p_1 p_2}, \quad \hat{\beta}_j = \frac{k_j p_1}{p_1 p_2}, \quad z_j = (1 - \hat{\alpha}_j)(1 - \hat{\beta}_j), \\ \Delta_I^{(0)}(z) &\equiv 0, \quad \Delta_I^{(1)}(z) \equiv \frac{1}{2}\gamma - \frac{1}{4}\gamma \log(z), \\ \Delta_I^{(2)}(z) &\equiv \Delta_I^{(1)}(z) + \frac{1}{8}\gamma^2 - \frac{1}{8}\gamma^2 \log(z) + \frac{1}{24}\gamma^2 \log^2(z), \\ W_e(a, b) &\equiv 1 - \frac{m_e^2}{2p_1 p_2} \frac{(1-a)(1-b)}{(1-a)^2 + (1-b)^2} \left(\frac{a}{b} + \frac{b}{a} \right). \end{aligned} \quad (7.18)$$

Let us check the soft limit at first. In the case of more than one photon, if we take the soft limit $k_j \rightarrow 0$, keeping the momenta of the other photons constant, then θ_{kr} are generally all different. However, the sums over $d\sigma^{Born}/d\Omega$ in eq. (7.17)

combine into a simple average over all four angles, as in eq. (7.11). So the single photon distribution reduces to

$$D_{1I}^{(2,1)}(X, p_1, p_2, q_1, q_2, k_j) \sim \tilde{S}_I(k_j) \bar{\beta}_0^{(1,0)}(X, p_1, p_2, q_1, q_2) \quad (7.19)$$

and thus $\bar{\beta}_{1I}^{(2,1)}(X, p_1, p_2, q_1, q_2, k_j)$ is IR finite. The above discussion implies that extrapolations for $\bar{\beta}_0$ and $\bar{\beta}_1$ have to be of the same type.

The collinear limit is our another concern. If all of the photons are collinear to the initial or final fermions, then all of the angles $\theta_{ij}, i, j = 1, 2$, are identical and equal to the LL effective scattering angle for the hard process in the frame XMS. This will facilitate the introduction of the higher order LL corrections in the following.

Note that there are many equivalent ways, modulo a term of $O(m^2/s)$, to express the single-bremsstrahlung spin-summed differential distribution [105]. Our choice results from minimizing the machine rounding errors which implementing Monte Carlo programs [85, 88, 89]. And eq. (7.17) is explicitly expressed in terms of the Born differential cross sections, which helps the introduction of electroweak corrections.

The virtual correction term $[1 + \Delta_I^{(1)}(z_j)]$ is taken in the LL approximation and it agrees with the corresponding contribution in the Ref. [106]. In the limit $k_j \rightarrow 0$, we have $\Delta_I^{(1)}(z) \rightarrow \delta_I^{(1)}$ and $\bar{\beta}_{1F}^{(2)}$ is infrared finite. The factor $(1 + \delta_F^{(1)})$ represents the contribution from the simultaneous emission of the real initial and the virtual final photons.

The key ingredients for the $O(\alpha^r)$ final state $\bar{\beta}_{1F}^{(r)}$, $r = 1, 2$, is the single final state photon emission matrix element with up to one-loop virtual initial-/final-state

photon corrections:

$$\begin{aligned}
D_{1F}^{(r)}(X, p_1, p_2, q_1, q_2, k'_l) &= Q_f^2 \frac{\alpha}{4\pi^2} \frac{2q_1 q_2}{(k'_l q_1)(k'_l q_2)} W_f(\hat{\eta}_l, \hat{\zeta}_l) \\
&\times \left\{ \frac{(1 - \hat{\eta}_l)^2}{2} \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{r1}) + \frac{(1 - \hat{\zeta}_l)^2}{2} \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{r2}) \right\} \\
&\times \left[1 + \Delta_F^{(r-1)}(z_l) \right] (1 + \delta_I^{(r-1)}), \tag{7.20}
\end{aligned}$$

where

$$\begin{aligned}
\eta_l &= \frac{k'_l q_2}{q_1 q_2}, \quad \zeta_l = \frac{k'_l q_1}{q_1 q_2}, \quad \hat{\eta}_l = \frac{\eta_l}{1 + \eta_l + \zeta_l}, \\
\Delta_F^{(0)}(z) &\equiv 0, \quad \Delta_F^{(1)}(z) \equiv \frac{1}{2} \gamma_f + \log(z) \\
W_e(a, b) &\equiv 1 - \frac{m_f^2}{2q_1 q_2} \frac{(1-a)(1-b)}{(1-a)^2 + (1-b)^2} \left(\frac{a}{b} + \frac{b}{a} \right). \tag{7.21}
\end{aligned}$$

The discussions on the ISR distribution of eq. (7.17) also works for the FSR distribution above.

7.1.4 Two Real Photons with Virtual Corrections

The contributions $\bar{\beta}_{2II}^{(2)}$, $\bar{\beta}_{2FF}^{(2)}$, and $\bar{\beta}_{2IF}^{(2)}$ are related to the emission of two real photons, two initial, two final and one initial and on final, respectively. The contribution $\bar{\beta}_{2II}^{(2)}$, related to the emission of two real photons, is defined formally:

$$\begin{aligned}
&\bar{\beta}_{2II}^{(r)}(X, p_1, p_2, q_1, q_2, k_j, k_k) \\
&= D_{2II}^{(r)}(X, p_1, p_2, q_1, q_2, k_j, k_k) - \tilde{S}_I(k_j) \bar{\beta}_{1I}^{(r-1)}(X, p_1, p_2, q_1, q_2, k_k) \\
&\quad - \tilde{S}_I(k_k) \bar{\beta}_{1I}^{(r-1)}(X, p_1, p_2, q_1, q_2, k_j) - \tilde{S}_I(k_j) \tilde{S}_I(k_k) \bar{\beta}_0^{(r-2)}(X, p_1, p_2, q_1, q_2), \\
&r = 2, 3, \tag{7.22}
\end{aligned}$$

And the contributions $\bar{\beta}_{2FF}^{(2)}$ and $\bar{\beta}_{2IF}^{(2)}$, related to the emission of two final photons and one initial photon and one final photon, are defined respectively:

$$\begin{aligned}
& \bar{\beta}_{2FF}^{(r)}(X, p_1, p_2, q_1, q_2, k'_l, k'_m) \\
&= D_{2FF}^{(r)}(X, p_1, p_2, q_1, q_2, k'_l, k'_m) - \tilde{S}_F(k'_l) \bar{\beta}_{1F}^{(r-1)}(X, p_1, p_2, q_1, q_2, k'_m) \\
&\quad - \tilde{S}_F(k'_m) \bar{\beta}_{1F}^{(r-1)}(X, p_1, p_2, q_1, q_2, k'_l) - \tilde{S}_F(k'_l) \tilde{S}_F(k'_m) \bar{\beta}_0^{(r-2)}(X, p_1, p_2, q_1, q_2), \\
& r = 2, 3, \tag{7.23}
\end{aligned}$$

$$\begin{aligned}
& \bar{\beta}_{2IF}^{(r)}(X, p_1, p_2, q_1, q_2, k_j, k'_l) \\
&= D_{2IF}^{(r)}(X, p_1, p_2, q_1, q_2, k_j, k'_l) - \tilde{S}_I(k_j) \bar{\beta}_{1F}^{(r-1)}(X, p_1, p_2, q_1, q_2, k'_l) \\
&\quad - \tilde{S}_I(k'_l) \bar{\beta}_{1I}^{(r-1)}(X, p_1, p_2, q_1, q_2, k_j) - \tilde{S}_I(k_j) \tilde{S}_F(k'_l) \bar{\beta}_0^{(r-2)}(X, p_1, p_2, q_1, q_2), \\
& r = 2, 3. \tag{7.24}
\end{aligned}$$

The new terms $D_{2II}^{(2)}$, $D_{2FF}^{(2)}$, and $D_{2IF}^{(2)}$ in the above expressions are the differential distributions for the double bremsstrahlung. They are not calculated directly from Feynman diagrams but they are set up in the following way: if one photon is soft and the other is hard, then the single-bremsstrahlung expressions of (7.17) and (7.20) are recovered; if both photons are hard and collinear, then the proper LL limit is also recovered.

The definition of the double real ISR distribution is

$$\begin{aligned}
& D_{2II}^{(2)}(X, p_1, p_2, q_1, q_2, k_1, k_2) \\
& \equiv Q_e^4 \frac{\alpha}{4\pi^2} \frac{2p_1 p_2}{(k_1 p_1)(k_1 P_2)} \frac{\alpha}{4\pi^2} \frac{2p_1 p_2}{(k_2 p_1)(k_2 p_2)} W_e(\hat{\alpha}_1, \hat{\beta}_1) W_e(\hat{\alpha}_2, \hat{\beta}_2) \\
& \left\{ \Theta(v_1 - v_2) \left[1 + \Delta_{II}^{(r-1)}(z_1, z_{12}) \right] (1 + \delta_F^{(r-1)}) \left[\chi_2(\hat{\alpha}_1; \hat{\alpha}'_2, \hat{\beta}'_2) \right. \right. \\
& \quad \left. \left. \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) + \chi_2(\hat{\alpha}_1; \hat{\alpha}'_2, \hat{\beta}'_2) \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) + \right] + \right. \\
& \quad \left. \left\{ \Theta(v_2 - v_1) \left[1 + \Delta_{II}^{(r-1)}(z_2, z_{21}) \right] (1 + \delta_F^{(r-1)}) \left[\chi_2(\hat{\beta}_1; \hat{\alpha}'_2, \hat{\beta}'_2) \right. \right. \right. \\
& \quad \left. \left. \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{2r}) + \chi_2(\hat{\beta}_1; \hat{\alpha}'_2, \hat{\beta}'_2) \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) + \right] \right\}, \tag{7.25}
\end{aligned}$$

where

$$\begin{aligned}
\hat{\alpha}'_1 &= \frac{\hat{\alpha}_1}{1 - \hat{\alpha}_2}, \quad \hat{\alpha}'_2 = \frac{\hat{\alpha}_2}{1 - \hat{\alpha}_1}, \quad \hat{\beta}'_1 = \frac{\hat{\beta}_1}{1 - \hat{\beta}_2}, \quad \hat{\beta}'_2 = \frac{\hat{\beta}_2}{1 - \hat{\beta}_1}, \\
v_i &= \hat{\alpha}_i + \hat{\beta}_i, \quad z_i = (1 - \hat{\alpha}_i)(1 - \hat{\beta}_i), \quad z_{ij} = (1 - \hat{\alpha}_i - \hat{\alpha}_j)(1 - \hat{\beta}_i - \hat{\beta}_j), \\
\chi_2(u; a, b) &\equiv \frac{1}{4}(1 - u)^2[(1 - a)^2 + (1 - b)^2], \\
\Delta_{II}^{(0)} &= 0, \quad \Delta_{II}^{(1)}(z_i - \frac{1}{6}\gamma \log(z_i), z_{ij}) = \frac{1}{2}\gamma - \frac{1}{6}\gamma \log(z_{ij}). \tag{7.26}
\end{aligned}$$

The variables $\hat{\alpha}_i, \hat{\beta}_i$ for the i th photon are defined as in eq. (7.18).

In order to understand the construction, we examine the realization of the LL collinear limit in the exact single-bremsstrahlung matrix element of eq. (7.17). Suppose a photon carrying the fraction x_1 of the beam energy is collinear with p_1 , then $\hat{\alpha}_1 \sim x, \hat{\beta}_1 \sim 0$, all four angles are the same $\theta_{sr} \rightarrow \theta^*$ and we at once recover the

correct LL formula

$$\begin{aligned} & \frac{1}{2}(1 - \hat{\alpha}_1)^2 \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(\theta_{1r}) + \frac{1}{2}(1 - \hat{\beta}_1)^2 \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(\theta_{2r}) \\ & \rightarrow \frac{1}{2}[1 + (1 - x)^2] \frac{d\sigma^{Born}}{d\Omega}(\theta^*). \end{aligned}$$

So it is natural to use the the angular-dependent Altarelli-Parisi (AP) factors of the type

$$\frac{1}{2}[(1 - \hat{\alpha}_2)^2 + (1 - \hat{\beta}_2)^2] \frac{1}{2}[(1 - \hat{\alpha}_1)^2 + (1 - \hat{\beta}_1)^2]$$

for the double emission. But the above expression is too simple to reproduce correctly the result of the double convolution of the AP kernels in the case that both photons are collinear with the same fermion

$$\frac{1}{2}[1 + (1 - x_1)^2] \frac{1}{2} \left[1 + \left(1 - \frac{x_2}{1 - x_1} \right)^2 \right] \frac{d\sigma^{Born}}{d\Omega}(\theta^*),$$

where $x_2/(1 - x_1)$ reflects the energy loss in the emission cascade because of the emission of k_1 . In order to deal with the above situation, we need to reconstruct angular dependent AP factor as

$$\frac{1}{2}[(1 - \hat{\alpha}_1)^2 + (1 - \hat{\beta}_1)^2] \frac{1}{2}[(1 - \hat{\alpha}'_2)^2 + (1 - \hat{\beta}'_2)^2].$$

The above formula fits both kinds of the LL collinear limit, when two photons are collinear with a single beam or each of them follows a different beam. Finally, we reproduce the limit in which one photon is hard and the other is soft, $v_2 = \hat{\alpha}_2 + \hat{\beta}_2 \rightarrow 0$. In this case, we split the above double-bremsstrahlung angular dependent AP factor into two parts

$$\begin{aligned} \chi_2(\hat{\alpha}_1; \hat{\alpha}'_2, \hat{\beta}'_2) &= \frac{1}{2}(1 - \hat{\alpha}_1)^2 \frac{1}{2}[(1 - \hat{\alpha}'_2)^2 + (1 - \hat{\beta}'_2)^2], \\ \chi_2(\hat{\beta}_1; \hat{\alpha}'_2, \hat{\beta}'_2) &= \frac{1}{2}(1 - \hat{\beta}_1)^2 \frac{1}{2}[(1 - \hat{\alpha}'_2)^2 + (1 - \hat{\beta}'_2)^2], \end{aligned} \quad (7.27)$$

and relate each one with the corresponding $\frac{d\sigma^{Born}}{d\Omega}$ as we did in eq. (7.17). The order in the cascade does not affect the result. So we just symmetrize over the two orderings in the cascade (Bose-Einstein symmetrization).

The construction above gives the correct limit $D_{2II}^{(2)}(k_1, k_2) \rightarrow \tilde{S}(k_2)D_{1I}^{(1)}(k_2)$ for $v_1 = \text{const}$ and $v_2 \rightarrow 0$. Consequently, $\bar{\beta}_{II}^{(2)}(X, p_1, p_2, q_1, q_2, k_1, k_2)$ is finite in the limit of one or both photon momenta approaching to zero.

The double final-state bremsstrahlung distribution is constructed in an analogous way:

$$\begin{aligned}
& D_{2FF}^{(2)}(X, p_1, p_2, q_1, q_2, k_1, k_2) \\
& \equiv Q_f^4 \frac{\alpha}{4\pi^2} \frac{2q_1 p_2}{(k'_1 q_1)(k'_1 p_2)} \frac{\alpha}{4\pi^2} \frac{2q_1 p_2}{(k'_2 q_1)(k'_2 p_2)} W_f(\hat{\eta}_1, \hat{\zeta}_1) W_e(\hat{\eta}_2, \hat{\zeta}_2) \\
& \left\{ \Theta(v'_1 - v'_2) \left[\chi_2(\hat{\eta}_1; \hat{\eta}'_2, \hat{\zeta}'_2) \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) \right. \right. \\
& \quad \left. \left. + \chi_2(\hat{\eta}_1; \hat{\eta}'_2, \hat{\zeta}'_2) \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) \right] + \Theta(v'_2 - v'_1) \left[\chi_2(\hat{\zeta}_1; \hat{\eta}'_2, \hat{\zeta}'_2) \right. \right. \\
& \quad \left. \left. \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{2r}) + \chi_2(\hat{\zeta}_1; \hat{\eta}'_1, \hat{\zeta}'_1) \sum_{r=1,2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{1r}) + \right] \right\} \\
& \times [1 + \Delta_I^{(r-1)}(z_j)], \tag{7.28}
\end{aligned}$$

where

$$\eta'_1 = \frac{\eta_1}{1 + \eta_2}, \quad \eta'_2 = \frac{\eta_2}{1 + \eta_1}, \quad \zeta'_1 = \frac{\zeta_1}{1 + \zeta_2}, \quad \zeta'_2 = \frac{\zeta_2}{1 + \zeta_1}. \tag{7.29}$$

Note that the definition of the "primed" Sudakov variables is different from that in the ISR case, because the fermion momenta $q_{1,2}$ are affected by photon emission. The virtual corrections are absent because we restrict the FSR to $O(\alpha^2)_{LL}$. The above expression is tagged with $r = 2, 3$ for $O(\alpha^r)$, but the FSR is implemented only in $O(\alpha^2)$ and the only correction in $O(\alpha^3)$ is the ISR one loop correction.

The distribution for one photon from the initial-state and the other from the final-state at $O(\alpha^r)$, $r = 1, 2$, is given by

$$\begin{aligned}
& D_{2IF}^{(r)}(X, p_1, p_2, q_1, q_2, k_j, k'_l) \\
&= Q_e^2 \frac{\alpha}{4\pi^2} \frac{2p_1 p_2}{(k_j p_1)(k_j p_2)} W_e(\hat{\alpha}_j, \hat{\beta}_j) Q_f^2 \frac{\alpha}{4\pi^2} \frac{2q_1 q_2}{(k'_l q_1)(k'_l q_2)} W_f(\hat{\eta}_l, \hat{\zeta}_l) \times \\
& \left\{ \frac{(1 - \hat{\alpha}_j)^2 (1 - \hat{\eta}_l)^2}{2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{11}) + \frac{(1 - \hat{\alpha}_j)^2 (1 - \hat{\zeta}_l)^2}{2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{12}) \right. \\
& \left. + \frac{(1 - \hat{\beta}_j)^2 (1 - \hat{\eta}_l)^2}{2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{21}) + \frac{(1 - \hat{\beta}_j)^2 (1 - \hat{\zeta}_l)^2}{2} \frac{d\sigma^{Born}}{d\Omega}(X^2, \theta_{22}) \right\} \\
& \times [1 + \Delta_I^{(r-1)}(z_1)][1 + \Delta_F^{(r-1)}(z'_2)], \tag{7.30}
\end{aligned}$$

where $\hat{\alpha}_j, \hat{\beta}_j, \hat{\eta}_l, \hat{\zeta}_l$ and other componets are defined in eqs. (7.18) and (7.21).

7.1.5 Three Real Photons

The differential distribution for three real ISR photons is obtained by the triple convolution of the AP kernel for each beam. In spite of the primary importance of the collinear limit, preserving all soft limit is also our concern while constructing the fully differential triple-photon distribution.

In these limit the triple-photon differential distribution must reproduce the previously define Born, single-, and double-bremmsstrahlung distributions times the

the appropriate soft factors. If not, we may encounter an issue of the IR finiteness of

$$\begin{aligned}
& \bar{\beta}_{3III}^{(3)}(X, p_i, q_j, k_1, k_2, k_3) \\
&= D_{3III}^{(3)}(X, p_i, q_j, k_1, k_2, k_3) - \tilde{S}_I(k_1)\bar{\beta}_{2II}^{(2)}(X, p_i, q_j, k_2, k_3) \\
&\quad - \tilde{S}_I(k_2)\bar{\beta}_{2II}^{(2)}(X, p_i, q_j, k_1, k_3) - \tilde{S}_I(k_3)\bar{\beta}_{2II}^{(2)}(X, p_i, q_j, k_1, k_2) \\
&\quad - \tilde{S}_I(k_1)\tilde{S}_I(k_2)\bar{\beta}_{1I}^{(1)}(X, p_i, q_j, k_3) - \tilde{S}_I(k_3)\tilde{S}_I(k_1)\bar{\beta}_{1I}^{(1)}(X, p_i, q_j, k_2) \\
&\quad - \tilde{S}_I(k_2)\tilde{S}_I(k_3)\bar{\beta}_{1I}^{(1)}(X, p_i, q_j, k_1) - \tilde{S}_I(k_1)\tilde{S}_I(k_2)\tilde{S}_I(k_3)\bar{\beta}_0^{(0)}(X, p_i, q_j).
\end{aligned} \tag{7.31}$$

As in the case of the double real ISR photons, the guideline for constructing the differential distributions includes (i) the hardest photon decides which of the angles is used in $\frac{d\sigma^{\text{Born}}}{d\Omega}(X^2, \theta_{1r})$ and (ii) we have to sum over all orderings in a cascade emission of several photons from one beam (Bose-Einstein Symmetrization). For the case of three real photons there are no virtual corrections.

The construction for three real ISR is

$$\begin{aligned}
& D_{3III}^{(3)}(X, p_1, p_2, q_1, q_2, k_1, k_2, k_3) \\
&\equiv \prod_{l=1,3} Q_e^2 \frac{\alpha}{4\pi^2} \frac{2p_1 p_2}{(k_l p_1)(k_l p_2)} W_e(\hat{\alpha}_l, \hat{\beta}_l) \left\{ \Theta(v_1 - v_2)\Theta(v_2 - v_3) \right. \\
&\quad \left[\chi_3(\hat{\alpha}_1; \hat{\alpha}'_2, \hat{\beta}'_2, \hat{\alpha}''_3, \hat{\beta}''_3) \sum_{r=1,2} \frac{d\sigma^{\text{Born}}}{d\Omega}(X^2, \theta_{1r}) + \chi_3(\hat{\beta}_1; \hat{\alpha}'_2, \hat{\beta}'_2, \hat{\alpha}''_3, \hat{\beta}''_3) \right. \\
&\quad \left. \left. \sum_{r=1,2} \frac{d\sigma^{\text{Born}}}{d\Omega}(X^2, \theta_{2r}) \right] + \text{remaining five permutations of } (1,2,3) \right\} \tag{7.32}
\end{aligned}$$

where

$$\begin{aligned}
\chi_3(u_1; a_2, b_2, a_3, b_3) &\equiv \frac{1}{8}(1 - u_1)^2 [(1 - a_2)^2 + (1 - b_2)^2] [(1 - a_2)^2 + (1 - b_2)^2], \\
\hat{\alpha}''_3 &= \frac{\bar{\alpha}_3}{1 - \bar{\alpha}_1 - \bar{\alpha}_2}, \quad \hat{\beta}''_3 = \frac{\bar{\beta}_3}{1 - \bar{\beta}_1 - \bar{\beta}_2}.
\end{aligned} \tag{7.33}$$

7.2 Amplitudes for Coherent Exclusive Exponentiation

The coherent exclusive exponentiation was first introduced in Ref. [107], which is rooted in the YFS exponentiation [78]. The exponentiation procedure which is a reorganization of the QED perturbative series such that the IR divergences are summed to up infinite order, is realized at the spin-amplitude level for both real and virtual IR divergences. This is contrast with the EEX which is based on the traditional YFS theory, in which the isolation for the real IR-singularities is achieved for the squared spin-summed spin amplitudes. The computation of the spin amplitudes is finished with the help of the Kleiss and Stirling Spinor technique [52] (please read Chapter Three for details). It is very interesting that the IR cancellation of the CEEX occur for the integrated cross sections as usual even though the CEEX is formulated completely in terms of the spin amplitudes. In this section, we shall introduce the construction of the CEEX matrix element, the IR cancellation in the CEEX scheme and the virtual and photonic correction for CEEX.

7.2.1 Master Formula

Let us define the Lorentz-invariant phase-space as

$$\int d\text{Lips}_n(P; p_1, p_2, \dots, p_n) = \int (2\pi)^4 \delta\left(P - \sum_{i=1}^n p_i\right) \prod_{i=1}^n \frac{d^3p}{(2\pi)^3 2p_i^0}, \quad (7.34)$$

then we write the CEEX total cross section for the process

$$e^-(p_a) + e^+(p_b) \rightarrow f(p_c) + \bar{f}(p_d) + \gamma(k_1) + \gamma(k_2) + \dots + \gamma(k_n), \quad n = 0, 1, 2, \dots \quad (7.35)$$

with polarized beams and decays of unstable final fermions which are sensitive to fermion spin polarizations as follows:

$$\begin{aligned} \sigma^{(r)} &= \frac{1}{\text{flux}(s)} \sum_{n=0}^{\infty} \int d\text{Lip} s_{n+2}(p_a + p_b; p_c, p_d, k_1, \dots, k_n) \\ &\quad \times \rho_{\text{CEEX}}^{(r)}(p_a, p_b, p_c, p_d, k_1, \dots, k_n), \end{aligned} \quad (7.36)$$

where, in the CMS (center of mass) $\text{flux}(s) = 2s + O(m_e^2)$,

$$\begin{aligned} &\rho_{\text{CEEX}}^{(r)}(p_a, p_b, p_c, p_d, k_1, \dots, k_n) \\ &= \frac{1}{n!} \exp[Y(\Omega; p_a, \dots, p_d)] \bar{\Theta}(\Omega) \sum_{\sigma_i=\pm 1} \sum_{\lambda_i, \lambda_j=\pm 1} \sum_{i,j,l,m=0}^3 \hat{\epsilon}_a^i \hat{\epsilon}_b^j \sigma_{\lambda_a \bar{\lambda}_a}^i \sigma_{\lambda_b \bar{\lambda}_b}^j \\ &\quad \times \mathfrak{M}_n^{(r)} \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \left[\mathfrak{M}_n^{(r)} \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \bar{\lambda} \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \right]^* \sigma_{\lambda_c \lambda_d}^l \sigma_{\bar{\lambda}_c \bar{\lambda}_d}^j \hat{h}_c^l \hat{h}_d^m. \end{aligned} \quad (7.37)$$

Assume that the s -channel exchanges dominate and resonances are included, then we can define the complete set of spin amplitudes for the n photon emission, in $O(\alpha^r)_{\text{CEEX}}$, $r = 0, 1, 2$, as follows,

$$\begin{aligned} &\mathfrak{M}_n^{(1)} \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \\ &\equiv \sum_{\varphi \in \{I, F\}^n} \prod_{i=1}^n \mathfrak{s}_{[i]}^{\{\varphi_i\}} \left\{ \hat{\beta}_0^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X_\varphi \right) + \sum_{j=1}^n \frac{\hat{\beta}_{1\{\varphi_j\}}^{(1)} \left(\begin{array}{c} pk_j \\ \lambda \sigma_j \end{array} ; X_\varphi \right)}{\mathfrak{s}_{[j]}^{\{\varphi_i\}}} \right\}, \end{aligned} \quad (7.38)$$

$$\begin{aligned}
& \mathfrak{M}_n^{(2)} \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \\
& \equiv \sum_{\varphi \in \{I, F\}^n} \prod_{i=1}^n \mathfrak{s}_{[i]}^{\{\varphi_i\}} \left\{ \hat{\beta}_0^{(2)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X_\varphi \right) + \sum_{j=1}^n \frac{\hat{\beta}_{1\{\varphi_j\}}^{(2)} \left(\begin{array}{c} pk_j \\ \lambda \sigma_j \end{array} ; X_\varphi \right)}{\mathfrak{s}_{[j]}^{\{\varphi_i\}}} \right. \\
& \quad \left. + \sum_{1 \leq j < n \leq n} \frac{\hat{\beta}_{2\{\varphi_j \varphi_l\}}^{(2)} \left(\begin{array}{c} pk_j k_l \\ \lambda \sigma_j \sigma_l \end{array} ; X_\varphi \right)}{\mathfrak{s}_{[j]}^{\{\varphi_j\}} \mathfrak{s}_{[l]}^{\{\varphi_l\}}} \right\}.
\end{aligned} \tag{7.39}$$

In order to simplify our expressions, we introduce a compact collective notation:

$$\left(\begin{array}{c} p \\ \lambda \end{array} \right) = \left(\begin{array}{c} p_a p_b p_c p_d \\ \lambda_a \lambda_b \lambda_c \lambda_d \end{array} \right) \tag{7.40}$$

for the fermion four-momenta p_A , $A = a, b, c, d$ (i.e., $p_1 = p_a$, $p_2 = p_b$, $q_1 = p_c$, $q_2 = p_d$) and helicities λ_A , $A = a, b, c, d$. For $k = 1, 2, 3$, σ^k are the Pauli matrices and $\sigma_{\lambda, \mu}^0 = \delta_{\lambda, \mu}$ is the unit matrix. The components $\hat{\epsilon}_1^j$, $\hat{\epsilon}_2^k$, where $j, k = 1, 2, 3$, are the components of the conventional spin-polarization vectors of the incoming fermions, defined in the GPS fermion rest frames (Plase read Appendix D for details). We define $\hat{\epsilon}_A^0 = 1$ in a nonstandard way (i.e., $p_A \cdot \hat{\epsilon}_A = m_e$, $A = a, b$). The polarimeter vector \hat{h}_C are similarly defined in th proper GPS rest frames of the final unstable fermions ($p_C \cdot \hat{h}_C = m_f$, $C = c, d$).

Next, we introduce and explain the notation for the IR integration limits for the real photons in eqs. (7.36) and (7.37). The factor $\bar{\Theta}(\Omega)$ in eq. (7.36) defines

the IR integration limits for all real photons. For a single photon, Ω is the domain surrounding the IR divergences point $k = 0$, which is excluded from the MC phase-space. In CEEX, Ω is the same for all photons since there is no actual difference between ISR and FSR photons. We define a characteristic function $\Theta(\Omega, k)$ of the IR domain Ω as

$$\Theta(\Omega, k) = \begin{cases} 1 & \text{for } k \in \Omega, \\ 0 & \text{for } k \notin \Omega. \end{cases} \quad (7.41)$$

The characteristic functions for the part of the phase-space include in the MC integration for a single real photon is $\bar{\Theta}(\Omega, k) = 1 - \Theta(\Omega, k)$. Similarly, the characteristic function for all real photons is as follows:

$$\bar{\Theta}(\Omega) = \prod_{i=1}^n \bar{\Theta}(\Omega, k_i). \quad (7.42)$$

In the computation corresponding to the KKMC program we define Ω in a traditional way with the photon-energy cut condition $k^0 < E_{\min}$.

The YFS form factor [78] for Ω defined with the condition $k^0 < E_{\min}$ is

$$\begin{aligned} Y(\Omega; p_a, \dots, p_d) = & Q_e^2 Y_\Omega(p_a, p_b) + Q_f^2 Y_\Omega(p_c, p_d) + Q_e Q_f Y_\Omega(p_a, p_c) \\ & + Q_e Q_f Y_\Omega(p_a, p_c) - Q_e Q_f Y_\Omega(p_a, p_c) - Q_e Q_f Y_\Omega(p_a, p_c), \end{aligned} \quad (7.43)$$

where

$$\begin{aligned} Y_\Omega(p, q) \equiv & 2\alpha \widehat{B}(\Omega, p, q) + 2\alpha \Re B(\Omega, p, q) \\ \equiv & -2\alpha \frac{1}{8\pi^2} \int \frac{d^3 k}{k^0} \Theta(\Omega; k) \left(\frac{p}{kp} - \frac{q}{kq} \right)^2 \\ & + 2\alpha \Re \int \frac{d^4 k}{k^2} \frac{i}{(2\pi)^3} \left(\frac{2p-k}{2kp-k^2} - \frac{2q-k}{2kq-k^2} \right) \end{aligned} \quad (7.44)$$

is given analytically in terms of dilogarithm functions.

The coherent sum is taken over the set $\{\varphi\} = \{I, F\}$ for all 2^n partitions, the single partition φ is defined as a vector $(\varphi_1, \varphi_2, \dots, \varphi_n)$ where $\varphi_i = I$ for an ISR photon and $\varphi_F = F$ for an FSR photon. The set of all partitions is explicitly written as follows:

$$\{\varphi\} = \{(I, I, I, \dots, I), (F, I, I, \dots, I), (I, F, I, \dots, I), \dots, (F, F, F, \dots, F)\}.$$

The s -channel four-momentum in the resonant s -channel propagator is $X_\varphi = p_a + p_b - \sum_{\varphi_i=I} k_i$.

The soft amplitude factors $\mathfrak{s}_{[i]}^{\{\omega\}}$, $\omega = I, F$, are defined as follows:

$$\begin{aligned} \mathfrak{s}_{[i]}^{\{I\}} &\equiv \mathfrak{s}_{[i]}^{\{I\}}(k) = -eQ_e \frac{b_\sigma(k, p_a)}{2k_i p_a} + eQ_e \frac{b_\sigma(k, p_b)}{2k_i p_b}, \\ \left| \mathfrak{s}_{[i]}^{\{I\}} \right|^2 &= -\frac{e^2 Q_e^2}{2} \left(\frac{p_a}{k_i p_a} - \frac{p_b}{k_i p_b} \right)^2, \end{aligned} \quad (7.45)$$

$$\begin{aligned} \mathfrak{s}_{[i]}^{\{F\}} &\equiv \mathfrak{s}_{[i]}^{\{I\}}(k) = +eQ_f \frac{b_\sigma(k, p_c)}{2k_i p_c} + eQ_e \frac{b_\sigma(k, p_d)}{2k_i p_d}, \\ \left| \mathfrak{s}_{[i]}^{\{F\}} \right|^2 &= -\frac{e^2 Q_f^2}{2} \left(\frac{p_c}{k_i p_c} - \frac{p_d}{k_i p_d} \right)^2, \end{aligned} \quad (7.46)$$

$$b_\sigma(k, p) = \sqrt{2} \frac{\bar{u}_\sigma(k) \not{p} u_\sigma(\zeta)}{\bar{u}_{-\sigma} u_\sigma(\zeta)}. \quad (7.47)$$

The simplest IR-finite $\hat{\beta}$ function $\hat{\beta}_0^{(0)}$ is the Born spin amplitude times a kinematical factor

$$\hat{\beta}_0^{(0)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) = \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) \frac{X^2}{(p_c + p_d)^2}. \quad (7.48)$$

Note that the Born spin amplitude $\mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right)$ is an essential block for building all of the spin amplitudes. Applying the Feynman rules and the basic massive spinors with the definite GPS helicities, the Born spin amplitudes for $e^-(p_a) + e^+(p_b) \rightarrow$

$f(p_c) + f(p_d)$ are given by

$$\begin{aligned}
\mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) &= \mathfrak{B} \left(\begin{array}{c} p_a p_b p_c p_d \\ \lambda_a \lambda_b \lambda_c \lambda_d \end{array} ; X \right) = \mathfrak{B} \left[\begin{array}{c} p_b p_a \\ \lambda_b \lambda_a \end{array} \right] \left[\begin{array}{c} p_c p_d \\ \lambda_c \lambda_d \end{array} \right] (X) \\
&= \mathfrak{B}_{[ba][cd]}(X) = ie^2 \sum_{B=\gamma, Z} \prod_B^{\mu\nu} (X) (G_{e,\mu}^B)_{[ba]} (G_{f,\mu}^B)_{[cd]} H_B \\
&= \sum_{B=\gamma, Z} \mathfrak{B}_{[ba][cd]}^B(X), \\
(G_{e,\mu}^B)_{[ba]} &\equiv \bar{v}(p_b \lambda_b) G_{e,\mu}^B u(p_a, \lambda_a) \\
(G_{f,\mu}^B)_{[ba]} &\equiv \bar{v}(p_c \lambda_c) G_{f,\mu}^B u(p_d, \lambda_d) \\
G_{e,\mu}^B &= \gamma_\mu \sum_{\lambda=\pm} \omega_\lambda g_\lambda^{B,e} \\
G_{f,\mu}^B &= \gamma_\mu \sum_{\lambda=\pm} \omega_\lambda g_\lambda^{B,f}, \quad \omega_\lambda = \frac{1}{2}(1 + \lambda\gamma_5), \\
\Pi_B^{\mu\nu}(X) &= \frac{g^{\mu\nu}}{X^2 - M_B^2 + i\Gamma_B X^2 / M_B^2}, \tag{7.49}
\end{aligned}$$

where $g_\lambda^{B,f}$ are the chiral coupling constants ($\lambda = \pm = R, L$) of the vector boson $B = \gamma, Z$ to the fermion f in units of the electric charge e . Usually, the "hook function" H_B is trivial: $H_\gamma = H_Z = 1$. And spinor products can be reorganized with the help of Chisholm identities:

$$\mathfrak{B}_{[ba][cd]}^B(X) = 2ie^2 \frac{\delta_{\lambda_a, -\lambda_b} [g_{\lambda_a}^{B,e} g_{-\lambda_a}^{B,f} T_{\lambda_c \lambda_a} T'_{\lambda_b \lambda_d} + g_{\lambda_a}^{B,e} g_{-\lambda_a}^{B,f} U'_{\lambda_c \lambda_b} U_{\lambda_a \lambda_d}]}{X^2 - M_B^2 + i\Gamma_B X^2 / M_B} \tag{7.50}$$

where

$$\begin{aligned}
T_{\lambda_c \lambda_a} &= \bar{u}(p_c, \lambda_c) u(p_a, \lambda_a) = S(p_c, m_c, \lambda_c, p_a, 0, \lambda_a), \\
T'_{\lambda_b \lambda_d} &= \bar{v}(p_b, \lambda_b) v(p_d, \lambda_d) = S(p_b, 0, -\lambda_b, p_d, -m_d, -\lambda_d), \\
U'_{\lambda_c \lambda_b} &= \bar{u}(p_c, \lambda_c) v(p_b, -\lambda_b) = S(p_c, m_c, \lambda_c, p_b, 0, \lambda_b), \\
U_{\lambda_a \lambda_d} &= \bar{u}(p_a, -\lambda_a) v(p_d, \lambda_d) = S(p_b, 0, -\lambda_a, p_d, -m_d, -\lambda_d). \tag{7.51}
\end{aligned}$$

7.2.2 IR Structure in CEEX

In this subsection, we discuss the mechanism of the IR-cancellation in the CEEX scheme. Let us begin with the infinite-order perturbative expression for the total cross section given by the standard quantum-mechanical expression of the type “matrix element squared modulus times phase-space”:

$$\begin{aligned} \sigma^{(\infty)} &= \sum_{n=0}^{\infty} \frac{1}{n!} \int d\tau_n(p_a + p_b; p_c, p_d, k_1, \dots, k_n) \\ &\quad \times \frac{1}{4} \sum_{\lambda, \sigma_1, \dots, \sigma_n = \pm} \left| \mathcal{M}_n \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \right|^2, \end{aligned} \quad (7.52)$$

where $d\tau_n$ is the respective $(n\gamma + 2f)$ -Lorentz-invariant phase-space, and \mathcal{M}_n are the corresponding spin amplitudes. In order to simplify the discussion, we take the unpolarized case without narrow resonances here.

According to the YFS theory [78], all virtual IR corrections can be relocated into an exponential form factor order by order and in infinite order

$$\mathcal{M}_n^{(\infty)} = \exp[\alpha B_4(p_a, p_b, p_c, p_d)] \mathfrak{m}_n^{(\infty)}. \quad (7.53)$$

Since the convergence of the perturbative series is questionable, the equation above is practically treated as a symbolic representation of the order-by-order relation, which reads at $O(\alpha^r)$,

$$\mathcal{M}_n^{(r)} = \sum_{l=0}^{r-n} \frac{(\alpha B_4)^{r-l}}{(r-l)!} \mathfrak{m}_n^{[l+n]} \quad (n \leq r), \quad (7.54)$$

where the index l is the number of loops in $\mathfrak{M}_n^{[l+n]}$. The $\mathfrak{M}_n^{[l+n]}$'s are not only free of the virtual IR divergences, they are also universal: they are the same in every perturbative order r . The formula above can be reformulated as follows:

$$\mathfrak{m}_n^{(r)} = \sum_{l=0}^{r-n} \mathfrak{m}_n^{[l+n]} = \{\exp[-\alpha B_4(p_a, p_b, p_c, p_d)] \mathcal{M}_n^{(r)}\}_{|O(\alpha^r)}, \quad (7.55)$$

where $\mathcal{M}_n^{(r)}$ has to be evaluated from the Feynman diagrams in at least $O(\alpha^r)$. The above treatments are exactly the same as in Chapter Six.

The YFS form factor B_4 for $e^-(p_a) + e^+(p_b) \rightarrow f(p_c) + \bar{f}(p_d) + n\gamma$ is

$$\alpha B_4(p_a, p_b, p_c, p_d) = \int \frac{d^4k}{k^2 - m_\gamma^2 + i\epsilon} \frac{i}{(2\pi)^3} |J_I(k) - J_F(k)|^2, \quad (7.56)$$

where

$$\begin{aligned} J_I &= eQ_e[\hat{J}_a(k) - \hat{J}_b(k)], \\ J_F &= eQ_f[\hat{J}_c(k) - \hat{J}_d(k)], \\ \hat{J}_f^\mu(k) &= \frac{2p_f^\mu + k^\mu}{k^2 + 2k \cdot p_f + i\epsilon}. \end{aligned} \quad (7.57)$$

Using the identity $(\sum_k Z_k J_k)^2 = -\sum_{i>k} Z_i Z_k (J_i - J - k)^2$ for $\sum Z_k = 0$, where Z_k is the charge of the particle with minus charge in the initial or final state respectively, we can rewrite B_4 as the sum of the simpler dipole components. Note that the IR singularities are regularized with a fictitious photon mass m_γ .

$$\begin{aligned} B_4(p_a, p_b, p_c, p_d) &= Q_e^2 B_2(p_a, p_b) + Q_f^2 B_2(p_c, p_d) + Q_e Q_f B_2(p_a, p_c) \\ &\quad + Q_e Q_f B_2(p_b, p_d) - Q_e Q_f B_2(p_a, p_d) - Q_e Q_f B_2(p_b, p_c), \end{aligned} \quad (7.58)$$

$$B_2(p_i, p_j) \equiv \int \frac{d^4k}{k^2 - m_\gamma^2 + i\epsilon} \frac{i}{(2\pi)^3} [\hat{J}(p_i, k) - \hat{J}(p_j, k)]. \quad (7.59)$$

Next, we elaborate the isolation of the real IR divergences in the CEEX scheme, which differs in essential details from the original YFS method [78] (please read Chapter Six for details). The essential difference is that we do not square the amplitudes immediately, and it is done numerically at a later stage. We use the results of the basic analysis of the real IR divergences of Ref. [78]. The basic analysis of IR cancellations

in Ref. [78] is done in terms of the currents

$$j_f^\mu(k) = \frac{2p_f^\mu}{2p_f \cdot k}, \quad f = a, b, c, d. \quad (7.60)$$

The above currents are simply related to the \mathfrak{s} factors:

$$\begin{aligned} \mathfrak{s}_\sigma^I(k) &= \text{const} \times Q_e(j_a - j_b) \cdot \epsilon_\sigma(\beta) \\ \mathfrak{s}_\sigma^F(k) &= \text{const} \times Q_f(j_c - j_d) \cdot \epsilon_\sigma(\beta). \end{aligned} \quad (7.61)$$

Note that the whole structure of the real IR singularities is completely controlled by the squares of the currents $|j(k)|^2$, for $j = j_a - j_b$ or $j = j_c - j_d$ because only the squares $|j(k)|^2$ are IR divergent and other contractions do not matter. Similarly, if we express spin amplitudes in terms of \mathfrak{s} factors, only the squares $|\mathfrak{s}|^2$ are IR divergent and not the interference terms.

The IR-divergent part of \mathcal{M} is proportional to the products of n \mathfrak{s} factors

$$\mathfrak{M}_n \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \sim \hat{\beta}_0 \left(\begin{array}{c} p \\ \lambda \end{array}; X \right) \mathfrak{s}_{\sigma_1}(k_1) \mathfrak{s}_{\sigma_2}(k_2) \dots \mathfrak{s}_{\sigma_n}(k_n), \quad (7.62)$$

$$\mathfrak{s}_\sigma(k) \equiv \mathfrak{s}_\sigma^{\{F\}} + \mathfrak{s}_\sigma^{\{I\}}. \quad (7.63)$$

Considering there are also non-leading IR singularities, the whole real-IR structure is revealed in the following decomposition:

$$\begin{aligned} &\mathfrak{M}_n^{(\infty)}(k_1, k_2, k_3, \dots, k_n) \\ &= \hat{\beta}_0 \prod_{s=1}^n \mathfrak{s}(k_s) + \sum_{j=1}^n \hat{\beta}_1(k_j) \prod_{s \neq j} \mathfrak{s}(k_s) + \sum_{j_1 > j_2} \hat{\beta}_2(k_{j_1}, k_{j_2}) \prod_{s \neq j_1, j_2} \mathfrak{s}(k_s) \\ &\quad + \sum_{j_1 > j_2 > j_3} \hat{\beta}_2(k_{j_1}, k_{j_2}, k_{j_3}) \prod_{s \neq j_1, j_2, j_3} \mathfrak{s}(k_s) + \dots \\ &\quad + \sum_{j=1} \hat{\beta}_{n-1}(k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_n) \mathfrak{s}(k_j) + \hat{\beta}_n(k_1, k_2, k_3, \dots, k_n) \end{aligned} \quad (7.64)$$

where the functions $\hat{\beta}_i$ are IR free and include finite loop corrections to infinite order. The decomposition of eq. (7.64) also has the order-by-order representation and it is written as follows:

$$\begin{aligned}
& \mathfrak{M}_n^{(n+l)}(k_1, k_2, k_3, \dots, k_n) \\
&= \hat{\beta}_0^{(l)} \prod_{s=1}^n \mathfrak{s}(k_s) + \sum_{j=1}^n \hat{\beta}_1^{(1+l)}(k_j) \prod_{s \neq j} \mathfrak{s}(k_s) + \sum_{j_1 > j_2} \hat{\beta}_2^{(2+l)}(k_{j_1}, k_{j_2}) \prod_{s \neq j_1, j_2} \mathfrak{s}(k_s) \\
&+ \sum_{j_1 > j_2 > j_3} \hat{\beta}_3^{(3+l)}(k_{j_1}, k_{j_2}, k_{j_3}) \prod_{s \neq j_1, j_2, j_3} \mathfrak{s}(k_s) + \dots \\
&+ \sum_{j=1}^n \hat{\beta}_{n-1}^{(n-1+l)}(k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_n) \mathfrak{s}(k_j) + \hat{\beta}_n^{(n+l)}(k_1, k_2, k_3, \dots, k_n) \\
&= \prod_{s=1}^n \mathfrak{s}(k_s) \left\{ \hat{\beta}_0^{(l)} + \sum_{j=1}^n \frac{\hat{\beta}_1^{(1+l)}(k_j)}{\mathfrak{s}(k_j)} + \sum_{j_1 < j_2} \frac{\hat{\beta}_2^{(2+l)}(k_{j_1}, k_{j_2})}{\mathfrak{s}(k_{j_1}) \mathfrak{s}(k_{j_2})} \right. \\
&+ \sum_{j_1 < j_2 < j_3} \frac{\hat{\beta}_3^{(3+l)}(k_{j_1}, k_{j_2}, k_{j_3})}{\mathfrak{s}(k_{j_1}) \mathfrak{s}(k_{j_2}) \mathfrak{s}(k_{j_3})} + \sum_{j=1}^n \frac{\hat{\beta}_{n-1}^{(n-1+l)}(k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_n)}{\prod_{s \neq j} \mathfrak{s}(k_s)} \\
&\left. + \frac{\hat{\beta}_n^{(n+l)}(k_1, k_2, k_3, \dots, k_n)}{\prod_s \mathfrak{s}(k_s)} \right\} \tag{7.65}
\end{aligned}$$

at $O(\alpha^r)$, $r = n + l$. The functions $\hat{\beta}_n^{(n+l)}(k_1, k_2, k_3, \dots, k_n)$ include up to l -loop corrections. The $\hat{\beta}_n^{(n+l)}(k_1, k_2, k_3, \dots, k_n)$ functions are not only completely IR finite, but are universal as well. This feature is essential for reversing the relations of eq. (7.65). From this feature, we could calculate $\hat{\beta}_n^{(n+l)}$ from $\mathfrak{M}_n^{(r)}$ directly from the

Feynman rules order-by-order:

$$\begin{aligned}
\hat{\beta}_0^{(l)} &= \mathfrak{M}_0^{(l)}, \\
\hat{\beta}_1^{(l+1)}(k_1) &= \mathfrak{M}_1^{(1+l)}(k_1) - \hat{\beta}_0^{(l)} \mathfrak{s}(k_1), \\
\hat{\beta}_2^{(2+l)}(k_1, k_2) &= \mathfrak{M}_2^{(2+l)}(k_1, k_2) - \hat{\beta}_1^{(l+1)}(k_1) \mathfrak{s}(k_2) - \hat{\beta}_1^{(l+1)}(k_2) \mathfrak{s}(k_1) \\
&\quad - \hat{\beta}_0^{(l)} \mathfrak{s}(k_1) \mathfrak{s}(k_2), \\
\hat{\beta}_3^{(3+l)}(k_1, k_2, k_3) &= \mathfrak{M}_3^{(3+l)}(k_1, k_2, k_3) - \hat{\beta}_2^{(2+l)}(k_1, k_2) \mathfrak{s}(k_3) - \hat{\beta}_2^{(2+l)}(k_1, k_3) \mathfrak{s}(k_2) \\
&\quad - \hat{\beta}_2^{(2+l)}(k_2, k_3) \mathfrak{s}(k_1) - \hat{\beta}_1^{(1+l)}(k_1) \mathfrak{s}(k_2) \mathfrak{s}(k_3) - \hat{\beta}_1^{(1+l)}(k_2) \mathfrak{s}(k_1) \mathfrak{s}(k_3) \\
&\quad - \hat{\beta}_1^{(1+l)}(k_3) \mathfrak{s}(k_1) \mathfrak{s}(k_2) - \hat{\beta}_0^{(l)} \mathfrak{s}(k_1) \mathfrak{s}(k_2) \mathfrak{s}(k_3), \dots, \\
\hat{\beta}_n^{(n+l)}(k_1, \dots, k_n) \\
&= \mathfrak{M}_n^{(n+l)}(k_1, \dots, k_n) - \sum_{j=1}^n \hat{\beta}_{n-1}^{(n-1+l)}(k_1, \dots, k_{j-1}, k_{j+1}, \dots, k_n) \mathfrak{s}(k_j) \\
&\quad - \sum_{j_1 < j_2}^{n-2} \hat{\beta}_{n-2}^{(n-2+l)}(k_1, \dots, k_{j_1-1}, k_{j_1+1}, \dots, k_{j_2-1}, k_{j_2+1}, \dots, k_n) \mathfrak{s}(k_{j_1}) \mathfrak{s}(k_{j_2}) - \dots \\
&\quad - \sum_{j_1 < j_2} \hat{\beta}_2^{(2+l)}(k_{j_1}, k_{j_2}) \prod_{s \neq j_1, j_2} \mathfrak{s}(k_s) - \sum_{j=1}^n \hat{\beta}_1^{(1+l)}(k_j) \prod_{s \neq j} \mathfrak{s}(k_s) - \bar{\beta}_0^{(l)} \prod_{s=1}^n \mathfrak{s}(k_s).
\end{aligned} \tag{7.66}$$

The above set of equations is a recursive rule, i.e., the higher-order $\hat{\beta}$'s are built in terms of the lower-order ones. In practical calculations one does not go to the infinite order but stops at some $O(\alpha^r)$ and the above set of equations is truncated for $\hat{\beta}_n^{(n+l)}$ by the requirement $n + l \leq r$. The above truncation is valid since we omit higher order $\hat{\beta}$'s which are IR finite. As a result of the fixed-order truncation, eq. (7.64)

reads as follows

$$\begin{aligned}
& \mathfrak{M}_n^{(r)}(k_1, k_2, k_3, \dots, k_n) \\
&= \prod_{s=1}^n \mathfrak{s}(k_s) \left\{ \bar{\beta}_0^{(r)} + \sum_{j=1}^n \frac{\bar{\beta}_1^{(r)}(k_j)}{\mathfrak{s}(k_j)} + \sum_{j_1 < j_2} \frac{\bar{\beta}_2^{(r)}(k_{j_1}, k_{j_2})}{\mathfrak{s}(k_{j_1})\mathfrak{s}(k_{j_2})} \right. \\
&\quad \left. + \sum_{j_1 < j_2 < j_3} \frac{\bar{\beta}_3^{(r)}(k_{j_1}, k_{j_2}, k_{j_3})}{\mathfrak{s}(k_{j_1})\mathfrak{s}(k_{j_2})\mathfrak{s}(k_{j_3})} + \sum_{j_1 < j_2 < \dots < j_r} \frac{\bar{\beta}_3^{(r)}(k_{j_1}, k_{j_2}, \dots, k_{j_r})}{\mathfrak{s}(k_{j_1})\mathfrak{s}(k_{j_2}) \dots \mathfrak{s}(k_{j_r})} \right\}.
\end{aligned} \tag{7.67}$$

The formula above represents the general finite-order $O(\alpha^r)_{exp}$ case. For $r = 0$ case only the first term survives, and in the $O(\alpha^2)$ case there are three terms. The CEEX spin amplitudes in eq. (7.36) represent the case of $r = 0, 1, 2$.

Let us give an explicit example: in the recursive calculations of $\hat{\beta}$ in $O(\alpha^3)$, one needs to calculate $\hat{\beta}_0^{(l)}$, $l = 0, 1, 2, 3$; $\hat{\beta}_1^{(1+l)}$, $l = 0, 1, 2$; $\hat{\beta}_2^{(2+l)}$, $l = 0, 1$; and $\hat{\beta}_3^{(3)}$.

Therefore, according to eq. (7.66), we have

$$\begin{aligned}
\hat{\beta}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix} &= \mathfrak{M}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix}, \quad l = 0, 1, 2, \\
\hat{\beta}_1^{(1+l)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= \mathfrak{M}_1^{(1+l)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} - \hat{\beta}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix} \mathfrak{s}_{\sigma_1}(k_1), \quad l = 0, 1, \\
\hat{\beta}_2^{(2)} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} &= \mathfrak{M}_1^{(2)} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} - \hat{\beta}_1^{(1)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} \mathfrak{s}_{\sigma_2}(k_2) \\
&\quad - \hat{\beta}_1^{(1)} \begin{pmatrix} pk_2 \\ \lambda\sigma_2 \end{pmatrix} \mathfrak{s}_{\sigma_1}(k_1) - \hat{\beta}_0^{(0)} \begin{pmatrix} p \\ \lambda \end{pmatrix} \mathfrak{s}_{\sigma_1}(k_1)\mathfrak{s}_{\sigma_2}(k_2),
\end{aligned} \tag{7.68}$$

where the amplitude \mathfrak{M} is given by eq. (7.55).

At fixed-order $O(\alpha^r)_{\text{CEEX}}$, we have

$$\begin{aligned} \sigma^{(r)} &= \sum_{n=0}^{\infty} \frac{1}{n!} \int d\tau_n(p_1 + p_2; p_3, p_4, k_1, \dots, k_n) \\ &\quad \times \exp[2\alpha \Re B_4(p_a, \dots, p_d)] \frac{1}{4} \sum_{spin} |\mathfrak{M}_n^{(r)}(k_1, k_2, \dots, k_n)|^2, \end{aligned} \quad (7.69)$$

where $\mathfrak{M}_n^{(r)}$ is given by eq. (7.67) and we factorize out the \mathfrak{s} factors

$$\frac{1}{4} \sum_{spin} |\mathfrak{M}_n^{(r)}(k_1, k_2, k_3, \dots, k_n)|^2 = d_n(k_1, k_2, k_3, \dots, k_n) \prod_{s=1}^n |\mathfrak{s}(k_s)|^2, \quad (7.70)$$

where

$$\begin{aligned} &d_n(k_1, k_2, k_3, \dots, k_n) \\ &= \left| \bar{\beta}_0^{(r)} + \sum_{j=1}^n \frac{\hat{\beta}_1^{(r)}(k_j)}{\mathfrak{s}(k_j)} + \sum_{j_1 < j_2}^n \frac{\hat{\beta}_2^{(r)}(k_{j_1}, k_{j_2})}{\mathfrak{s}(k_{j_1}) \mathfrak{s}(k_{j_2})} + \sum_{j_1 < j_2 < j_3}^n \frac{\hat{\beta}_3^{(r)}(k_{j_1}, k_{j_2}, k_{j_3})}{\mathfrak{s}(k_{j_1}) \mathfrak{s}(k_{j_2}) \mathfrak{s}(k_{j_3})} \right. \\ &\quad \left. + \dots + \sum_{j_1 < j_2 < \dots < j_r}^n \frac{\hat{\beta}_r^{(r)}(k_{j_1}, k_{j_2}, \dots, k_{j_r})}{\mathfrak{s}(k_{j_1}) \mathfrak{s}(k_{j_2}) \dots \mathfrak{s}(k_{j_r})} \right|^2. \end{aligned} \quad (7.71)$$

Apparently the function $d_n(k_1, k_2, k_3, \dots, k_n)$ is IR finite and we can set $m_\gamma \rightarrow 0$ in it. Besides $2\Re B_4$ the IR regulator m_γ remains in all $\mathfrak{s}(k_i)$ factors and in the lower phase-space boundary of all real photons in $\int \frac{d^3 k}{2k^0}$.

The IF finiteness of above total cross section can be checked by partial differentiation with respect to the photon mass

$$\begin{aligned} \frac{\partial}{\partial m_\gamma} \sigma^{(r)} &= \sum_{n=0}^{\infty} \int d\tau_n(P; p_3, p_4, k_1, \dots, k_n) \exp(2\alpha \Re B_4) \frac{\partial}{\partial m_\gamma} \{2\alpha \Re B_4\} \\ &\quad \times \frac{1}{4} \sum_{spin} |\mathfrak{M}_n^{(r)}(k_1, k_2, k_3, \dots, k_n)|^2 + \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{s=1}^n \\ &\quad \times \int d\tau_{n-1}(P; p_3, p_4, k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_n) \\ &\quad \times \exp(2\alpha \Re B_4) \frac{\partial}{\partial m_\gamma} \left\{ \int \frac{d^3 k_s}{(2\pi)^3 2k_s^0} |\mathfrak{s}(k_s)|^2 \right\} \prod_{j \neq s} |\mathfrak{s}(k_j)|^2 \\ &\quad \times d_n(k_1, k_2, \dots, k_s, \dots, k_n). \end{aligned} \quad (7.72)$$

Note that

$$\frac{\partial}{\partial m_\gamma} \left\{ \int \frac{d^3 k_s}{2k_s^0} |\mathfrak{s}(k_s)|^2 \right\}$$

is a δ -like measure concentrated at $k_s = 0$ and therefore we may use the limit

$$\begin{aligned} d_n(k_1, \dots, k_s, \dots, k_n) &\rightarrow d_n(k_1, k_2, \dots, k_{s-1}, 0, k_{s+1}, \dots, k_n) \\ &\equiv d_{n-1}(k_1, k_2, \dots, k_{s-1}, k_{s+1}, \dots, k_n). \end{aligned}$$

With the help of the limit above, we notice that all of the terms in the $\sum_{s=1}^n$ are identical so that we could sum them up after formally renaming the photon integration variables in the second in integral and rewrite eq. (7.72) in the following way:

$$\begin{aligned} \frac{\partial}{\partial m_\gamma} \sigma^{(r)} &= \sum_{n=0}^{\infty} \int d\tau_n(P; p_3, p_4, k_1, \dots, k_n) \\ &\quad \times \exp(2\alpha \Re B_4) \frac{1}{4} \sum_{spin} |\mathfrak{M}_n^{(r)}(k_1, k_2, k_3, \dots, k_n)|^2 \\ &\quad \times \frac{\partial}{\partial \mu} \left\{ 2\alpha \Re B_4 + \int \frac{d^3 k_s}{(2\pi)^3 2k_s^0} |\mathfrak{s}(k_s)|^2 \right\} \\ &= 0, \end{aligned} \tag{7.73}$$

where the independence of m_γ of the sum of the one-photon real and virtual integrals is because of the cancellation of the IR singularities in the YFS theory.

We have introduced the general notation for the IR domain Ω in eq. (7.42). Now it is time to exclude the Ω domain from the real photon phase space. Splitting the real photon integration phase space, the total cross section (7.69) is rewritten as

$$\begin{aligned} \sigma^{(r)} &= \sum_{n=0}^{\infty} \frac{1}{n!} \left\{ \int \frac{d^3 k_j}{(2\pi)^3 2K_j^0} |\mathfrak{s}(k_j)|^2 \Theta(\Omega, k_j) + \int \frac{d^3 k_j}{(2\pi)^3 2k_j^0} |\mathfrak{s}(k_j)|^2 \Theta(\Omega, k_j) \right\} \\ &\quad \times \int d\tau_0 \left(P - \sum_{j=1}^n k_j; p_3, p_4 \right) \exp(2\alpha \Re B_4) d_n(k_1, k_2, \dots, k_n). \end{aligned} \tag{7.74}$$

After expanding the binomial product into 2^n terms we consider the sum of all $\binom{n}{1} = n$ terms in which one photon is in Ω and the other ones are not:

$$\begin{aligned}
& \frac{1}{n!} \sum_{s=1}^n \int \frac{d^3 k_s}{(2\pi)^3 2k_s^0} |\mathfrak{s}(k_s)|^2 \Theta(\Omega, k_s) \prod_{j \neq s}^n \int \frac{d^3 k_j}{(2\pi)^3 2k_j^0} |\mathfrak{s}(k_j)|^2 \bar{\Theta}(\Omega, k_j) \\
& \times \int d\tau_0 \left(P - \sum_{j=1}^n k_j; p_3, p_4 \right) \exp(2\alpha \Re B_4) d_n(k_1, k_2, \dots, k_{s-1}, 0, k_{s+1}, \dots, k_n) \\
& = \frac{1}{n!} \binom{n}{1} \int \frac{d^3 k}{(2\pi)^3 2k^0} |\mathfrak{s}(k)|^2 \Theta(\Omega, k) \int d\tau_{n-1}(P; p_3, p_4, k_1, k_2, \dots, k_{n-1}) \\
& \times \prod_{j=1}^{n-1} \bar{\Theta}(\Omega, k_j) |\mathfrak{s}(k_j)|^2 d_{n-1}(k_1, k_2, \dots, k_{n-1}). \tag{7.75}
\end{aligned}$$

A similar summation is taken for the $\binom{n}{s}$ terms where s photons are in the IR domain Ω , leading to

$$\begin{aligned}
\sigma^{(r)} & = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{s=0}^n \binom{n}{s} \left(\int \frac{d^3 k}{(2\pi)^3 2k^0} |\mathfrak{s}(k)|^2 \Theta(\Omega, k) \right)^s \\
& \times \int d\tau_{n-s}(P; p_3, p_4, k_1, k_2, \dots, k_{n-s}) \prod_{j=1}^{n-s} \{ |\mathfrak{s}(k_j)|^2 \bar{\Theta}(\Omega) \} \\
& \times \exp(2\alpha \Re B_4) d_{n-s}(k_1, k_2, \dots, k_{n-s}) \\
& = \sum_{n=0}^{\infty} \frac{1}{n!} \int d\tau_n(P; p_3, p_4, k_1, k_2, \dots, k_n) \exp \left(\int \frac{d^3 k_j}{(2\pi)^3 2k_j^0} |\mathfrak{s}(k_j)|^2 \Theta(\Omega, k_j) \right) \\
& \times \exp[2\alpha \Re B_4(p_1, \dots, p_4)] \prod_{j=1}^n \{ |\mathfrak{s}(k_j)|^2 \Theta(\bar{\Omega}, k_j) \} d_n(k_1, k_2, \dots, k_n). \tag{7.76}
\end{aligned}$$

The additional overall exponential factor contains

$$\begin{aligned}
2\alpha \tilde{B}_4(p_1, \dots, p_4) & = \int \frac{d^3 k_j}{(2\pi)^3 2k_j^0} |\mathfrak{s}(k_j)|^2 \Theta(\Omega, k_j) \\
& = 2\alpha [Q_e^2 \tilde{B}_2(p_1, p_2) + Q_f^2 \tilde{B}_2(p_3, p_4) + Q_e Q_f \tilde{B}_2(p_1, p_3) \\
& \quad + Q_e Q_f \tilde{B}_2(p_2, p_4) - Q_e Q_f \tilde{B}_2(p_1, p_4) - Q_e Q_f \tilde{B}_2(p_2, p_3)], \tag{7.77}
\end{aligned}$$

and

$$\begin{aligned}\tilde{B}_2(p, q) &\equiv - \int \frac{d^3k}{(2\pi)^3 2k^0} \Theta(\Omega, k) [j_p(k) - j_q(k)]^2 \\ &\equiv \int \frac{d^3k}{k^0} \Theta(\Omega, k) \frac{-1}{8\pi^2} \left(\frac{p}{kp} - \frac{q}{kq} \right)^2.\end{aligned}\quad (7.78)$$

Furthermore, the YFS form factor is

$$Y(\Omega; p_1, \dots, p_4) = 2\alpha \tilde{B}_4(p_1, \dots, p_4) + 2\alpha \Re \tilde{B}_4(p_1, \dots, p_4). \quad (7.79)$$

In pursue of the completeness of the discussion, let us check the IR cancellations in the total cross section with Ω as the new regulator:

$$\begin{aligned}\sigma^{(r)} &= \sum_{n=0}^{\infty} \frac{1}{n!} \int d\tau_n(P; p_3, p_4, k_1, k_2, \dots, k_n) \prod_{j=1}^n \{ |\mathfrak{s}(k_j)|^2 \bar{\Theta}(\Omega, k_j) \} \\ &\quad \times \exp \left[2\alpha \tilde{B}_4(\Omega; p_1, \dots, p_4) + 2\alpha \Re B_4(p_1, \dots, p_4) \right] d_n(k_1, k_2, \dots, k_n).\end{aligned}\quad (7.80)$$

Now IR finiteness of the total cross section is converted into the independence of the domain Ω

$$\frac{\partial}{\partial \Omega} \sigma^{(r)} = 0. \quad (7.81)$$

This can be proved by the same argument for the photon mass m_γ . Considering $\Omega \rightarrow \Omega' = \Omega + \delta\Omega$, that is $\bar{\Omega}' = \bar{\Omega} - \delta\Omega$ and Ω' can be either larger or smaller than Ω ,

the only requirement is that both are very small. Consequently we have

$$\begin{aligned}
\sigma^{(r)} &= \sum_{n=0}^{\infty} \frac{1}{n!} \prod_{j=1}^n \left\{ \int \frac{d^3 k_j}{(2\pi)^3 2k_j^0} |\mathfrak{s}(k_j)|^2 \bar{\Theta}(\Omega', k_j) + \int \frac{d^3 k_j}{(2\pi)^3 2k_j^0} |\mathfrak{s}(k_j)|^2 \bar{\Theta}(\delta\Omega', k_j) \right\} \\
&\quad \times \int d\tau_0 \left(P - \sum k_j; p_3, p_4 \right) \exp[2\alpha \tilde{B}_4(\Omega; p_1, \dots, p_4) + 2\alpha \Re B_4(p_1, \dots, p_4)] \\
&\quad \times d_n(k_1, k_2, \dots, k_n) \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{s=0}^n \binom{n}{s} \left\{ \int \frac{d^3 k}{(2\pi)^3 2k_0} |\mathfrak{s}(k)|^2 \Theta(\delta\Omega, k) \right\}^s \int d\tau_{n-s}(P; p_3, p_4, k_1, \dots, k_{n-s}) \\
&\quad \times \prod_{j=1}^{n-s} |\mathfrak{s}(k_j)|^2 \Theta(\Omega', k_j) \exp[2\alpha \tilde{B}_4(\Omega; p_1, \dots, p_4) + 2\alpha \Re B_4(p_1, \dots, p_4)] \\
&\quad \times d_{n-s}(k_1, k_2, \dots, k_{n-s}) \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} \int d\tau_n(P; p_3, p_4, k_1, \dots, k_n) \exp \left[\int \frac{d^3 k}{(2\pi)^3 2k_0} |\mathfrak{s}(k)|^2 \Theta(\delta\Omega, k) \right. \\
&\quad \left. + 2\alpha \tilde{B}_4(\Omega; p_1, \dots, p_4) + 2\alpha \Re B_4(p_1, \dots, p_4) \right] \prod_{j=1}^n \{ |\mathfrak{s}(k_j)|^2 \bar{\Omega}(\Omega', k_j) \} \\
&\quad \times d_n(k_1, k_2, \dots, k_n), \tag{7.82}
\end{aligned}$$

7.2.3 Narrow Neutral Resonance in CEEX

We have introduced the general mechanism of the IR-cancellation in the CEEX scheme in the last subsection. In this subsection, we will introduce the possible formulation of CEEX. There are three possible versions of CEEX so far. We will mainly describe the the version of the resonant Born. These three possible verions are as follows.

(A) The version the non-resonant Born without partitions:

$$\mathfrak{M}_n^{(0)} \left(\begin{array}{c} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) = \prod_{i=1}^n [\mathfrak{s}_{\sigma_i}^I(k_i) + \mathfrak{s}_{\sigma_i}^F(k_i)] \mathfrak{B}_{[ba][cd]}. \tag{7.83}$$

(B) The version for the non-resonant Born with partitions:

$$\mathfrak{M}_n^{(0)} \begin{pmatrix} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{pmatrix} = \sum_{\varphi \in \{I, F\}^n} \prod_{i=1}^n \mathfrak{s}_{\sigma_i}^{\{\varphi_i\}}(k_i) \mathfrak{B}_{[ba][cd]}(X_\varphi). \quad (7.84)$$

(C) The version for the resonant Born:

$$\mathfrak{M}_n^{(0)} \begin{pmatrix} pk_1 k_2 \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{pmatrix} = \sum_{\varphi \in \{I, F\}^n} \prod_{i=1}^n \mathfrak{s}_{\sigma_i}^{\{\varphi_i\}}(k_i) \frac{X_\varphi^2}{(p_3 + p_4)^2} \sum_{R=\gamma, Z} \mathfrak{B}_{[ba][cd]}^B(X_\varphi) \exp[\alpha \Delta B_4^R(X_\varphi)]. \quad (7.85)$$

We define the additional form factor for the Z resonance for case (C):

$$\alpha \Delta B_4^Z(X) = \int \frac{d^4 k}{k^2 - m_\gamma^2 + i\epsilon} \frac{i}{(2\pi)^3} J_{I\mu}(k) [J_F^\mu(k)]^* \left(\frac{X^2 - \bar{M}^2}{(X - k)^2 - \bar{M}^2} - 1 \right), \quad (7.86)$$

where $\bar{M}^2 = M_Z^2 - iM_Z\Gamma_Z$. The currents J^μ are given by eq. (7.57), while for the nonresonant part $\Delta B_4^\gamma(X) = 0$.

Let us make a brief comparison among these three versions. The case (B) will become case (A) if we neglect the partition dependence of the four momentum in the Born amplitude: $\mathfrak{B}_{[ba][cd]}(X_\varphi) \rightarrow \mathfrak{B}_{[ba][cd]}(P)$, where $P = p_a + p_b$ or $P = p_c + p_d$ or any other which is independent of the momentum of the individual photon. This feature is due to

$$\prod_{i=1}^n [\mathfrak{s}_{\sigma_i}^{\{F\}}(k_i) + \mathfrak{s}_{\sigma_i}^{\{I\}}(k_i)] \equiv \sum_{\varphi \in \{I, F\}^n} \prod_{i=1}^n \mathfrak{s}_{\sigma_i}^{\{\varphi_i\}}(k_i). \quad (7.87)$$

Obviously case (C) is efficient for the resonant process while cases (A) and (B) are only suitable for nonresonant process. If (A) does not sum the higher orders, it has a clear advantage over (B), which is simpler computer code and less consumption of CPU time because of no summation over partition. However, (B) sums up the LL higher orders more efficiently than (A). Considering our aim is to cover the resonant process, it is natural to utilize (B) for the nonresonant background of the spin amplitudes.

Once the summation over partitions happens, it is also easy to apply the case (B) for the nonresonant background. In other words, if (C) is carried out, then (B) comes automatically.

After comparing these three versions, we focus on the case (C) now because it becomes (B) for nonresonant background component. For the narrow neutral resonance, the photons emitted during the production and decay processes are separated by a long time interval; they are therefore totally independent. In the perturbative QED, this fact is reflected in a certain class of cancellations between ISR and FSR photon on the one hand and the virtual and real corrections on the other hand. Because of the presence of narrow resonances, it is not sufficient to sum up the real emissions coherently, taking the energy shift in the resonance propagator into account [108,109]. It is also necessary to sum the virtual emission up to infinite order—this is why the resonance form factor $\exp(B_4^Z)$ is included in eq. (7.85). Next we will derive eq. (7.86) and show that the IFI cancellations do work to infinite order.

Let us rewrite the YFS function in a modified notation

$$\alpha B_4(p_a, \dots, p_d) = \int \frac{d^4k}{k^2 - m_\gamma^2 + i\epsilon} \frac{i}{(2\pi)^3} S(k), \quad (7.88)$$

where

$$S_I(k) = |J_I(k)|^2, \quad S_F(k) = |J_F(k)|^2, \quad S_{Int}(k) = -2\Re[J_I(k) \cdot J_F^*(k)] \quad (7.89)$$

Due to the presence of the narrow resonance, the YFS factorization of the virtual IR contribution must take into account the dependence of the scalar part of the resonance propagator on photon energies of order Γ . The relevant integrals with n

virtual photons is given by

$$I = (P^2 - \bar{M}^2) \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{\wp \in \mathcal{P}_n} \prod_{i=0}^n \int \frac{i}{(2\pi)^3} \frac{d^4 k_i}{k_i^2 - m_\gamma^2} S_{\wp_i}(k_i) \frac{1}{P_\wp^2 - \bar{M}^2}, \quad (7.90)$$

where $\bar{M}^2 = M^2 - iM\Gamma$, and \mathcal{P}_n is a set of all 3^n partitions $(\wp_1, \wp_2, \dots, \wp_n)$ with $\wp_i = \text{I, F, Int}$, and $P_\wp \equiv P - \sum_{\wp_i} k_i$ includes only the momenta of the photons in S_{Int} . The $(P^2 - M^2)$ factor is conventional, making the integral dimensionless. We will show that the integral above factorizes into the conventional YFS form factor and the additional non-IR factor due to the resonance $R = Z$:

$$I = \exp[\alpha B_4^R(m_\gamma, s, \bar{M})] = \exp[\alpha B_4(m_\gamma, s) + \alpha \Delta B_4^R(s, \bar{M})]. \quad (7.91)$$

We aim to find the analytical expression of the additional function ΔB_4^R . In the present computation, we adopt the following approximate formula [108, 109],

$$\alpha \Delta B_4^R(s') = -2Q_e Q_f \frac{\alpha}{\pi} \log\left(\frac{t}{u}\right) \log\left(\frac{\bar{M}^2 - s}{\bar{M}^2}\right) = -\frac{1}{2} \gamma_{\text{Int}} \log\left(\frac{\bar{M}^2 - s}{\bar{M}^2}\right). \quad (7.92)$$

In the following, we will derive the equation above and show explicitly that the above virtual interference part of the form factor cancels exactly with the corresponding real interference contributions.

Because the soft virtual photons entering into S_I and S_F in eq. (7.90) do not enter the resonance propagator, we factorize and sum up the contributions with S_I and S_F :

$$\begin{aligned} I &= \sum_{n_1=0}^{\infty} \frac{1}{n_1!} \prod_{i_1=0}^{n_1} \int \frac{i}{(2\pi)^3} \frac{d^4 k_{i_1}}{k_{i_1}^2 - m_\gamma^2} S_I(k_{i_1}) \sum_{n_2=0}^{\infty} \frac{1}{n_2!} \prod_{i_2=0}^{n_2} \int \frac{i}{(2\pi)^3} \frac{d^4 k_{i_2}}{k_{i_2}^2 - m_\gamma^2} S_F(k_{i_2}) \\ &\quad \times \sum_{n_3=0}^{\infty} \frac{1}{n_3!} \prod_{i_3=0}^{n_3} \int \frac{i}{(2\pi)^3} \frac{d^4 k_{i_3}}{k_{i_3}^2 - m_\gamma^2} S_{\text{Int}}(k_{i_3}) \frac{1}{\left(P - \sum_{j=1}^{n_3} k_j\right) - \bar{M}^2} \\ &\equiv \exp(\alpha B_i + \alpha B_F) \sum_{n=0}^{\infty} \frac{1}{n!} \prod_{i=0}^n \int \frac{i}{(2\pi)^3} \frac{d^4 k_i}{k_i^2 - m_\gamma^2} S_{\text{Int}}(k_i) \frac{1}{\left(P - \sum_{j=1}^n k_j\right) - \bar{M}^2}. \end{aligned} \quad (7.93)$$

Here we neglect the quadratic terms in the photon energies $O(k_i k_j)$

$$\begin{aligned}
\frac{1}{\left(P - \sum_{j=1}^n k_j\right) - \bar{M}^2} &\simeq \frac{1}{P^2 - 2P \sum_{j=1}^n k_j - \bar{M}^2} \\
&= \frac{1}{P^2 - \bar{M}^2} \frac{1}{1 - \sum_{j=1}^n \frac{2Pk_j}{P^2 - \bar{M}^2}} \\
&\simeq \frac{1}{P^2 - \bar{M}^2} \prod_{j=1}^n \frac{1}{1 - \frac{2Pk_j}{P^2 - \bar{M}^2}} \\
&\simeq \frac{1}{P^2 - \bar{M}^2} \prod_{j=1}^n \frac{P^2 - \bar{M}^2}{(P - k_j)^2 - \bar{M}^2}, \tag{7.94}
\end{aligned}$$

and this gives us

$$\begin{aligned}
I &= \exp(\alpha B_I + \alpha B_F) \exp\left(\int \frac{i}{(2\pi)^3} \frac{d^4 k}{k^2 - m_\gamma^2 + i\epsilon} S_{In}(k) \frac{P^2 - \bar{M}^2}{(P - k)^2 - \bar{M}^2}\right) \\
&= \exp[\alpha B_4(m_\gamma) + \alpha \Delta B_4^R(\Gamma)], \tag{7.95}
\end{aligned}$$

where

$$\alpha \Delta B_4^R(\Gamma) = \int \frac{i}{(2\pi)^3} \frac{d^4 k}{k} S_{In}(k) \left(\frac{P^2 - \bar{M}^2}{(P - k)^2 - \bar{M}^2} - 1\right). \tag{7.96}$$

As $k \rightarrow 0$ the emission amplitude can be expressed as

$$\mathcal{M} \rightarrow \frac{1}{k} \left\{ \epsilon_1 + O\left(\frac{k}{\bar{M}}\right) + \frac{k}{\Gamma_Z} \left[\epsilon_2 + O\left(\frac{k}{\bar{M}}\right) \right] \right\},$$

where $\epsilon_{1,2}$ are constants independent of k , so that

$$\left| \frac{2Pk_j}{P^2 - \bar{M}^2} \right| \ll 1,$$

namely, photon energy is below the resonance width. This constraint is completely analogous to the usual YFS expansion into an IR-singular part and the rest [78]. The approach we choose here is based on the fact that the virtual and real contributions from the IFI for photons with $E_\gamma > \Gamma$ do cancel as a result of the time separation between the production and decay. We shall show the cancellation mechanism is valid next.

Let us check analytically the real multiphoton emission contribution for the IFI. We began with the integral in which the total photon energy $K = \sum_{j=1}^n k_j$ is kept below $E_{\max} = v_{\max}\sqrt{s}$, where $\Gamma < E_{\max} \ll \sqrt{s}$:

$$\begin{aligned}
\sigma &= \sum_{n=0}^{\infty} \frac{1}{n!} \int \prod_{i=1}^n \frac{d^3 k_i}{(2\pi)^3 2k_i^0} \sum_{\sigma_1 \dots \sigma_n} \left| \sum_{\varphi \in \{I, F\}^n} \prod_{j=1}^n \mathfrak{s}_{[j]}^{\{\varphi_j\}} \frac{1}{X_{\varphi}^2 - \bar{M}^2} \exp[\alpha B_4^R(X_{\varphi})] \right|^2 \\
&\times \Theta \left(E_{\max} - \sum_{j=1}^n k_j \right) \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} \int_{K^0 < v\sqrt{s}} \prod_{i=1}^n \frac{d^3 k_i}{(2\pi)^3 2k_i^0} \sum_{\sigma_1 \dots \sigma_n} \sum_{\varphi, \varphi' \in \{I, F\}^n} \prod_{j=1}^n \mathfrak{s}_{[j]}^{\{\varphi_j\}} \mathfrak{s}_{[j]}^{*\{\varphi'_j\}} \\
&\times \frac{\exp[\alpha B_4^R(X_{\varphi})]}{X_{\varphi}^2 - \bar{M}^2} \left(\frac{\exp[\alpha B_4^R(X_{\varphi'})]}{X_{\varphi'}^2 - \bar{M}^2} \right)^* \\
&= \sum_{n=0}^{\infty} \frac{1}{n!} \int_{K^0 < v\sqrt{s}} \prod_{i=1}^n \frac{d^3 k_i}{2k_i^0} \sum_{\varphi, \varphi' \in \{I^2, F^2, IF, FI\}^n} \prod_{\varphi_j = I^2} 2\tilde{S}_I(k_j) \prod_{\varphi_j = F^2} 2\tilde{S}_F(k_j) \\
&\times \prod_{\varphi_j = IF} 2\tilde{S}_{Int}(k_j) \prod_{\varphi_j = FI} 2\tilde{S}_{Int}(k_j) \frac{\exp[\alpha B_4^R(P - K_I - K_{IF})]}{(P - K_I - K_{IF})^2 - \bar{M}^2} \\
&\times \left(\frac{\exp[\alpha B_4^R(P - K_I - K_{FI})]}{(P - K_I - K_{FI})^2 - \bar{M}^2} \right)^*, \tag{7.97}
\end{aligned}$$

where

$$\begin{aligned}
2(2\pi)^2 \tilde{S}_I(k_j) &= \sum_{\sigma_j} \left| \mathfrak{s}_{[j]}^{\{I\}} \right|^2, \quad 2(2\pi)^2 \tilde{S}_F(k_j) = \sum_{\sigma_j} \left| \mathfrak{s}_{[j]}^{\{F\}} \right|^2, \\
2(2\pi)^2 \tilde{S}_{Int}(k_j) &= \sum_{\sigma_j} \mathfrak{s}_{[j]}^{\{I\}} \left(\mathfrak{s}_{[j]}^{\{F\}} \right)^* = \sum_{\sigma_j} \mathfrak{s}_{[j]}^{\{F\}} \left(\mathfrak{s}_{[j]}^{\{I\}} \right)^*, \\
K_{I^2} &= \sum_{\varphi_j = I^2} k_j, \quad K_{F^2} = \sum_{\varphi_j = F^2} k_j, \quad K_{IF} = \sum_{\varphi_j = IF} k_j, \quad K_{FI} = \sum_{\varphi_j = FI} k_j, \\
K &= K_{I^2} + K_{F^2} + K_{IF} + K_{FI}. \tag{7.98}
\end{aligned}$$

The product of two sums, each over 2^n partitions $\varphi, \varphi' \in \{I, F\}^n$, is now replaced by the single sum over 4^n partitions $\varphi \in \{I^2, F^2, IF, FI\}^n$, where the IF, FI represent the interference terms. And the summation over the number of the photons can be

reorganized as follows:

$$\begin{aligned}
\sigma(v_{max}) &= \sum_{n_1=0}^{\infty} \frac{1}{n_1!} \int \prod_{i_1=1}^{n_1} \frac{d^3 k_{i_1}}{2k_{i_1}^0} 2\tilde{S}_I(k_{i_1}) \prod_{i_2=1}^{n_2} \frac{d^3 k_{i_2}}{2k_{i_2}^0} 2\tilde{S}_F(k_{i_2}) \\
&\times \prod_{i_3=1}^{n_3} \frac{d^3 k_{i_3}}{2k_{i_3}^0} 2\tilde{S}_{Int}(k_{i_3}) \prod_{i_4=1}^{n_4} \frac{d^3 k_{i_4}}{2k_{i_4}^0} 2\tilde{S}_{Int}(k_{i_4}) \\
&\times \frac{\exp[\alpha B_4^R(P - K_{II} - K_{IF})]}{(P - K_{II} - K_{IF})^2 - \bar{M}^2} \left(\frac{\exp[\alpha B_4^R(P - K_{II} - K_{IF})]}{(P - K_{II} - K_{IF})^2 - \bar{M}^2} \right)^* \\
&\times \Theta(E_{max} - K_{II}^0 - K_{FF}^0 - K_{IF}^0 - K_{FI}^0)
\end{aligned} \tag{7.99}$$

where $K_{I^2} = \sum_{i_1} k_{i_1}$, $K_{F^2} = \sum_{i_2} k_{i_2}$, $K_{IF} = \sum_{i_3} k_{i_3}$ and $K_{FI} = \sum_{i_4} k_{i_4}$. The sums over the pure initial- and final-state contributions, and over the interference contributions are well factorized and ready to be performed analytically. First, we integrate and sum up contributions from the very soft photons below $\epsilon\sqrt{s}$,

$$\begin{aligned}
\sigma(v_{max}) &= \int_0^{E_{max}} dE' \int_0^{E_{max}} \delta(E' - E_I - E_F - E_{Int}) dE_I dE_F dE_{IF} dE_{FI} \\
&\times \sum_{n_1=0}^{\infty} \frac{1}{n_1!} \prod_{i_1=1}^{n_1} \int_{k_{i_1}^0 > \epsilon E} \frac{d^3 k_{i_1}}{2k_{i_1}^0} 2\tilde{S}_I(k_{i_1}) \exp[2\alpha\tilde{B}_I(\epsilon E) + 2\alpha\Re B_I] \delta\left(E_I - \sum_{i_1} k_{i_1}^0\right) \\
&\times \sum_{n_2=0}^{\infty} \frac{1}{n_2!} \prod_{i_2=1}^{n_2} \int_{k_{i_2}^0 > \epsilon E} \frac{d^3 k_{i_2}}{2k_{i_2}^0} 2\tilde{S}_F(k_{i_2}) \exp[2\alpha\tilde{B}_F(\epsilon E) + 2\alpha\Re B_F] \delta\left(E_F - \sum_{i_2} k_{i_2}^0\right) \\
&\times \sum_{n_3=0}^{\infty} \frac{1}{n_3!} \prod_{i_3=1}^{n_3} \int_{k_{i_3}^0 > \epsilon E} \frac{d^3 k_{i_3}}{2k_{i_3}^0} 2\tilde{S}_I(k_{i_3}) \frac{\exp[\alpha\Delta B_4^R(P - K_{II} - K_{IF})]}{(P - K_{II} - K_{IF})^2 - \bar{M}^2} \\
&\times \exp[2\alpha\tilde{B}_{Int}(\epsilon E) + 2\alpha\Re B_{Int}] \delta\left(E_{Int} - \sum_{i_3} k_{i_3}^0\right) \\
&\times \sum_{n_4=0}^{\infty} \frac{1}{n_4!} \prod_{i_4=1}^{n_4} \int_{k_{i_4}^0 > \epsilon E} \frac{d^3 k_{i_4}}{2k_{i_4}^0} 2\tilde{S}_I(k_{i_4}) \left(\frac{\exp[\alpha\Delta B_4^R(P - K_{II} - K_{IF})]}{(P - K_{II} - K_{IF})^2 - \bar{M}^2} \right)^* \\
&\times \exp[2\alpha\tilde{B}_{Int}(\epsilon E) + 2\alpha\Re B_{Int}] \exp(2\alpha\Re\Delta B_4^R) \delta\left(E_{Int} - \sum_{i_3} k_{i_3}^0\right),
\end{aligned} \tag{7.100}$$

where $E = \frac{\sqrt{s}}{2}$. The integration over photon momenta gives

$$\begin{aligned}
\sigma(v_{max}) &= \int_0^{v_{max}} dv \delta(v - v_I - v_F - v_{IF} - v_{FI}) \\
&\times \int dv_I F(\gamma_I) v_I^{\gamma_I - 1} \exp[2\alpha \tilde{B}_I(E) + 2\alpha \Re B_I] \\
&\times \int dv_F F(\gamma_F) v_F^{\gamma_F - 1} \exp[2\alpha \tilde{B}_F(E) + 2\alpha \Re B_F] \\
&\times \int dv_{IF} F\left(\frac{\gamma_{Int}}{2}\right) \frac{1}{2} \gamma_{Int} v_{IF}^{\frac{1}{2}\gamma_{IF} - 1} \\
&\times \left(\frac{\exp\{\alpha \Delta B_4^R[s(1 - v_I)(1 - v_{FI})]\}}{s(1 - v_I)(1 - v_{FI}) - \bar{M}^2} \right) \exp[\alpha \tilde{B}_{Int}(E) + \alpha \Re B_{Int}] \\
&\times \int dv_{FI} F\left(\frac{\gamma_{Int}}{2}\right) \frac{1}{2} \gamma_{Int} v_{FI}^{\frac{1}{2}\gamma_{FI} - 1} \left(\frac{\exp\{\alpha \Delta B_4^R[s(1 - v_I)(1 - v_{FI})]\}}{s(1 - v_I)(1 - v_{FI}) - \bar{M}^2} \right)^* \\
&\times \exp[\alpha \tilde{B}_{Int}(E) + \alpha \Re B_{Int}], \tag{7.101}
\end{aligned}$$

which is explicitly free of IR singularities.

The main problem is whether the $\log\left(\frac{\Gamma}{M_Z}\right)$ terms in the interference subintegral

$$\begin{aligned}
I_{Int} &= \Re \int_0^{v_{max} - v_I - v_F - v_{FI}} dv_{IF} F\left(\frac{\gamma_{IF}}{2}\right) \frac{1}{2} \gamma_{IF} v_{IF}^{\frac{1}{2}\gamma_{Int} - 1} \\
&\times \frac{\exp\{\alpha \Delta B_4^R[s'(1 - v_{IF})]\}}{s'(1 - v_{IF}) - \bar{M}^2}. \tag{7.102}
\end{aligned}$$

can be cancelled perfectly. We ignore the term $\exp[\alpha \tilde{B}_{Int}(E) + \alpha \Re B_{Int}]$, because it does not depend on the resonance parameters. The bulk of the integral comes from the neighborhood of $v_{IF} = 0$ and the integrand is $\sim \frac{1}{v}$ at large v because of the resonance. Thus we can extend the integration limit to $\int_0^\infty dv_{Int}$. We could apply the standard techniques of the complex functions to evaluate the integral. We first

reformulate the integral as an integral over the discontinuity C_1 along the real axis

$$I_{Int} = F\left(\frac{\gamma_{IF}}{2}\right) \exp[\alpha \Delta B_4^R(s')] \frac{1}{i \sin\left(\frac{\pi}{2}\gamma_{Int}\right)} \times \int_{C_1} dz \frac{1}{2} \gamma_{Int} (-z)^{\frac{1}{2}\gamma_{Int}-1} \frac{1}{s' - \bar{M}^2 - s'z}. \quad (7.103)$$

because the contour can be closed with big circle, the integral is given by the residue at $z = 1 - \frac{\bar{M}^2}{s'}$:

$$\begin{aligned} I_{Int} &= F\left(\frac{\gamma_{IF}}{2}\right) \exp[\alpha \Delta B_4^R(s')] \frac{\frac{\pi}{2}\gamma_{Int}}{\sin\left(\frac{\pi}{2}\gamma_{Int}\right)} \left(\frac{\bar{M}^2 - s'}{s'}\right)^{\gamma_{Int}-1} \frac{1}{s'} \\ &= \frac{1}{\bar{M}^2 - s'} F\left(\frac{\gamma_{Int}}{2}\right) \frac{\frac{\pi}{2}\gamma_{Int}}{\sin\left(\frac{\pi}{2}\gamma_{Int}\right)} \exp[\alpha \Delta B_4^R(s')] \left(\frac{\bar{M}^2 - s'}{s'}\right)^{\frac{1}{2}\gamma_{Int}} \\ &= \frac{1}{\bar{M}^2 - s'} [1 + O(\gamma_{Int})], \end{aligned} \quad (7.104)$$

where we use the result below

$$\begin{aligned} \alpha \Delta B_4^R(s') &= -2Q_e Q_f \frac{\alpha}{\pi} \log\left(\frac{t}{u}\right) \log\left(\frac{\bar{M}^2 - s'}{\bar{M}^2}\right) \\ &= -\frac{1}{2}\gamma_{Int} \log\left(\frac{\bar{M}^2 - s'}{\bar{M}^2}\right). \end{aligned} \quad (7.105)$$

Therefore we have proved the full cancellation of the dependence on the resonance parameters for the integrated cross section.

As we have shown before, the $\hat{\beta}$ -functions can be derived with the recursive relation of eq. (7.66). The only additional work here is we must keep track of the type of the external real photon (ISR or FSR) and of the total photon momentum

after emission of the ISR photons:

$$\begin{aligned}
\hat{\beta}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P &= \mathfrak{M}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P, \quad l = 0, 1, 2, \\
\hat{\beta}_{1\{I\}}^{(1+l)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} ; P - k_1 &= \mathfrak{M}_{1\{I\}}^{(1+l)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} \\
&\quad - \hat{\beta}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P - k_1 \mathfrak{s}_{\sigma_1}^{\{I\}}(k_1), \quad l = 0, 1 \\
\hat{\beta}_{1\{F\}}^{(1+l)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} ; P &= \mathfrak{M}_{1\{F\}}^{(1+l)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} \\
&\quad - \hat{\beta}_0^{(l)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P \mathfrak{s}_{\sigma_1}^{\{F\}}(k_1), \quad l = 0, 1 \\
\hat{\beta}_{2\{\omega_1, \omega_2\}}^{(2)} \begin{pmatrix} pk_1 k_2 \\ \lambda\sigma_1 \sigma_2 \end{pmatrix} ; X_\omega &= \mathfrak{M}_{2\{\omega_1, \omega_2\}}^{(2)} \begin{pmatrix} pk_1 k_2 \\ \lambda\sigma_1 \sigma_2 \end{pmatrix} \\
&\quad - \hat{\beta}_{1\{\omega_1\}}^{(1)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} ; X_\omega \mathfrak{s}_{\sigma_2}^{\{\omega_2\}}(k_2) \\
&\quad - \hat{\beta}_{1\{\omega_2\}}^{(1)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} ; X_\omega \mathfrak{s}_{\sigma_1}^{\{\omega_1\}}(k_1) \\
&\quad - \hat{\beta}_0^{(0)} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} ; X_\omega \mathfrak{s}_{\sigma_1}^{\{\omega_1\}}(k_1) \mathfrak{s}_{\sigma_2}^{\{\omega_2\}}(k_2), \quad (7.106)
\end{aligned}$$

where $X_\omega = P - \sum_{\omega_i=I} k_i$, $P = p_a + p_b$.

The amplitude \mathfrak{M} in eq. (7.106) is given actually by eq. (7.55) with the form factor including the resonance part:

$$\begin{aligned} & \mathfrak{M}_{n\{\omega\}}^{(r)R} \left(\begin{array}{c} pk_1 \cdots k_n \\ \lambda\sigma_1 \cdots \sigma_n \end{array} ; X_\omega \right) \\ & = \left\{ \exp[-\alpha B_4 - \alpha \Delta B_4^R(X_\omega)] \mathcal{M}_{n\{\omega\}}^{(r)R} \left(\begin{array}{c} pk_1 \cdots k_n \\ \lambda\sigma_1 \cdots \sigma_n \end{array} ; X_\omega \right) \right\} \Big|_O (\alpha^r). \end{aligned} \quad (7.107)$$

As we see the type $R = \gamma, Z$ of the resonance form factor B_4^R must be adjusted to the type of the component in $\mathcal{M}^{(r)R}$.

7.2.4 Virtual Corrections, No Photons

So far we have only obtained the formal expressions of $\hat{\beta}$ -functions by recursive relations. We will accumulate the actual formulas for the $\hat{\beta}$ -functions contributing to the CEEX amplitudes with the case of no real photons and up to two virtual photons.

Let us begin with the case of the $O(\alpha^1)$ spin amplitudes with one virtual and zero real photons coming from the Feynman diagrams, which contribute the first order $\hat{\beta}_0^{(1)}$. The spin amplitudes are given by

$$\begin{aligned} \mathcal{M}_0^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) & = \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) [1 + Q_e^2 F_1(s, m_e, m_\gamma)] [1 + Q_f^2 F_1(s, m_f, m_\gamma)] \\ & \quad + \mathcal{M}_{Box}^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right), \end{aligned} \quad (7.108)$$

where F_1 is the standard electric form factor regularized with a photon mass. We neglect the magnetic form factor F_2 temporarily, It will be restored in the future. In F_1 we keep the exact final fermion mass.

In the present work we adopt the spin amplitudes for γ - γ and γ - Z boxes in the small mass approximation $\frac{m_e^2}{s} \rightarrow 0$, $\frac{m_f^2}{s} \rightarrow 0$, according to Refs. [110, 111],

$$\begin{aligned} \mathcal{M}_{\text{Box}}^{(1)} = & 2ie^2 \sum_{B=\gamma, Z} \frac{g_{\lambda_a}^{B,e} g_{-\lambda_a}^{B,f} T_{\lambda_c \lambda_a} T'_{\lambda_b \lambda_d} + g_{\lambda_a}^{B,e} g_{\lambda_a}^{B,f} U'_{\lambda_c \lambda_b} U_{\lambda_a \lambda_d}}{X^2 - M_B^2 + i\Gamma \frac{X^2}{M_B}} \\ & \times \delta_{\lambda_a, -\lambda_b} \delta_{\lambda_c, -\lambda_d} \frac{\alpha}{\pi} Q_e Q_f [\delta_{\lambda_a, \lambda_c} f_{BDP}(\bar{M}_B^2, m_\gamma, s, t, u) \\ & - \delta_{\lambda_a, -\lambda_c} f_{BDP}(\bar{M}_B^2, m_\gamma, s, u, t)], \end{aligned} \quad (7.109)$$

where

$$\begin{aligned} f_{BDP}(\bar{M}, m_\gamma, s, u, t) = & \log\left(\frac{t}{u}\right) \log\left(\frac{m_\gamma^2}{\sqrt{tu}}\right) - 2 \log\left(\frac{t}{u}\right) \log\left(\frac{\bar{M}_B^2 - s}{\bar{M}_B^2}\right) \\ & + Li_2\left(\frac{\bar{M}^2 + u}{\bar{M}_B^2}\right) - Li_2\left(\frac{\bar{M}^2 + t}{\bar{M}_B^2}\right) \\ & + \frac{(M_B^2 - s)(u - t - \bar{M}_B^2)}{u^2} \left\{ \log\left(\frac{-t}{s}\right) \log\left(\frac{\bar{M}_B^2 - s}{\bar{M}_B^2}\right) \right. \\ & \left. + Li_2\left(\frac{\bar{M}_B^2 + t}{\bar{M}_B^2}\right) - Li_2\left(\frac{\bar{M}_B^2 - s}{\bar{M}_B^2}\right) \right\} \\ & + \frac{(\bar{M}_B^2 - s)^2}{us} \log\left(\frac{\bar{M}_B^2 - s}{\bar{M}_B^2}\right) + \frac{\bar{M}^2 - s}{u} \log\left(\frac{-t}{\bar{M}_B^2}\right), \end{aligned} \quad (7.110)$$

$\bar{M}_Z^2 = M_Z^2 - iM_Z\Gamma_Z$, $\bar{M}_\gamma^2 = m_\gamma^2$, and the function f_{BDP} is from Ref. [111]. The standard Mandelstam variables s , t and u are defined as usual: $s = (p_a + p_b)^2$, $t = (p_a - p_c)^2$ and $u = (p_a - p_d)^2$. Since in the rest of the calculation we do not use $\frac{m_f^2}{s} \rightarrow 0$, we intend to replace the above box spin amplitudes with the finite mass result according to Ref [112].

Using eq. (7.107) we have

$$\hat{\beta}_0^{(1)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; X = \mathfrak{B} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; X [1 + \delta_{\text{virt}}^{(1)e}(s)][1 + \delta_{\text{virt}}^{(1)f}(s)] + \mathcal{R}_{\text{Box}}^{(1)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; X, \quad (7.111)$$

where

$$\begin{aligned}
\delta_{Virt}^{(1)e}(s) &= Q_e^2 F_1(s, m_e, m_\gamma) - Q_e^2 \alpha B_2(p_a, p_b, m_\gamma) = Q_e^2 \frac{\alpha}{\pi} \frac{1}{2} \bar{L}_e \\
\delta_{Virt}^{(1)f}(s) &= Q_f^2 F_1(s, m_f, m_\gamma) - Q_e^2 \alpha B_2(p_c, p_d, m_\gamma) = Q_f^2 \frac{\alpha}{\pi} \frac{1}{2} \bar{L}_f \\
\bar{L}_e &= \log\left(\frac{s}{m_e^2}\right) + i\pi - 1, \quad \bar{L}_f = \log\left(\frac{s}{m_f^2}\right) + i\pi - 1.
\end{aligned} \tag{7.112}$$

The IR subtraction in $\mathcal{M}_{\text{Box}}^{(1)}$ using eq. (7.107) results in the IR-finite \mathcal{R}_{Box} .

The above subtraction is equivalent to the following substitution:

$$f_{BDP}(\bar{M}_B^2, m_\gamma, s, t, u) \rightarrow f_{BDP}(\bar{M}^2, m_\gamma, s, t, u) - f_{IR}(m_\gamma, t, u), \tag{7.113}$$

where

$$\begin{aligned}
f_{IR}(m_\gamma, t, u) &= \frac{2}{\pi} B_2(p_a, p_c, m_\gamma) - \frac{2}{\pi} B_2(p_a, p_d, m_\gamma) \\
&= \log\left(\frac{t}{u}\right) \log\left(\frac{m_\gamma^2}{\sqrt{tu}}\right) + \frac{1}{2} \log\left(\frac{t}{u}\right),
\end{aligned} \tag{7.114}$$

and the additional resonance factor $\exp[-\alpha \Delta B_4^Z(s)]$ in eq. (7.107) includes the additional subtraction in the γ -Z box part:

$$f_{BDP}(s, t, u) \rightarrow f_{BDP}(s, t, u) - \alpha \Delta B_4^Z(s). \tag{7.115}$$

Our $O(\alpha^2)$ expressions for $\hat{\beta}_0^{(2)}$ are still incomplete because we neglected some trivial transposition of the diagrams among the second-order vertex diagrams. By eq. (7.107) we have

$$\hat{\beta}_0^{(2)} \begin{pmatrix} p \\ \lambda \\ X \end{pmatrix} = \mathfrak{B} \begin{pmatrix} p \\ \lambda \\ X \end{pmatrix} [1 + \delta_{Virt}^{(2)e}(s)][1 + \delta_{Virt}^{(2)f}(s)] + \mathcal{R}_{\text{Box}}^{(2)} \begin{pmatrix} p \\ \lambda \\ X \end{pmatrix}, \tag{7.116}$$

In the present calculation we omit the two-loop double box contribution in $\mathcal{R}_{\text{Box}}^{(2)}$. In fact we keep only the first-order box contribution $\mathcal{R}_{\text{Box}}^{(1)}$ in our incomplete $O(\alpha^2)$ -type matrix element. Note that the lack of the above contribution will not undermine the

validity of our approach because what we omit is IR finite. And since the contribution we neglect is expected to be numerically small, of $O(\alpha^2 L^1)$, our overall physical precision is still reliable.

According to Refs. [106, 113, 114], we have the $O(\alpha^2)$ corrections to the electric form factor as follows:

$$\begin{aligned}\delta_{Virt}^{(2)e}(s, m_e) &= \delta_{Virt}^{(1)e}(s) + \left(\frac{\alpha}{\pi}\right)^2 \left[\frac{\bar{L}_e^2}{8} + \bar{L}_e \left(\frac{3}{32} - \frac{3}{4}\zeta_2 + \frac{3}{2}\zeta_3 \right) \right], \\ \delta_{Virt}^{(2)f}(s, m_f) &= \delta_{Virt}^{(1)f}(s) + \left(\frac{\alpha}{\pi}\right)^2 \left[\frac{\bar{L}_f^2}{8} + \bar{L}_e \left(\frac{3}{32} - \frac{3}{4}\zeta_2 + \frac{3}{2}\zeta_3 \right) \right].\end{aligned}\tag{7.117}$$

Next, let us discuss the electroweak corrections in CEEX. In the absence of Electroweak corrections, the coupling constants of two neutral boson γ and Z are defined conventionally as

$$\begin{aligned}G_\lambda^{Z,f} &= g_V^{Z,f} - \lambda g_A^{Z,f}, \quad G_\lambda^{\gamma,f} = g_V^{\gamma,f}, \quad \lambda = \pm = R, L, \\ g_V^{\gamma,e} &= Q_e = -1, \quad g_V^{\gamma,f} = Q_f, \quad g_A^{\gamma,e} = 0, \quad g_A^{\gamma,f} = 0, \\ g_V^{Z,e} &= \frac{2T_e^3 - 4Q_e \sin^2 \theta_W}{16 \sin^2 \theta_W \cos^2 \theta_W}, \quad g_V^{Z,f} = \frac{2T_f^3 - 4Q_f \sin^2 \theta_W}{16 \sin^2 \theta_W \cos^2 \theta_W}, \\ g_A^{Z,e} &= \frac{2T_e^3}{16 \sin^2 \theta_W \cos^2 \theta_W}, \quad g_A^{Z,f} = \frac{2T_f^3}{16 \sin^2 \theta_W \cos^2 \theta_W},\end{aligned}\tag{7.118}$$

where T_f^3 is the isospin of the left-handed component of the fermion.

Electroweak corrections in CEEX are implemented using DIZET package, which is a part of the ZFITTER semi-analytical program. The actual execution of the electroweak corrections goes as follows: the γ and Z propagators are multiplied

by the corresponding two scalar form factors due to vacuum polarizations:

$$\begin{aligned}
H_\gamma &\rightarrow H_\gamma \times \frac{1}{2 - \Pi_\gamma}, \\
H_Z &\rightarrow H_Z \times 16 \sin^2 \theta_W \cos^2 \theta_W \frac{G_\mu M_Z^2}{\alpha_{QED} 8\sqrt{2}\pi} \rho_{EW}.
\end{aligned} \tag{7.119}$$

Additionally the vector coupling constants of Z boson are multiplied by extra form factors

$$\begin{aligned}
g_V^{Z,e} &= \frac{2T_e^3 - 4Q_e \sin^2 \theta_W}{16 \sin^2 \theta_W \cos^2 \theta_W} \rightarrow \frac{2T_e^3 - 4Q_e \sin^2 \theta_W F_{EW}^e(s)}{16 \sin^2 \theta_W \cos^2 \theta_W}, \\
g_V^{Z,f} &= \frac{2T_f^3 - 4Q_f \sin^2 \theta_W}{16 \sin^2 \theta_W \cos^2 \theta_W} \rightarrow \frac{2T_f^3 - 4Q_f \sin^2 \theta_W F_{EW}^f(s)}{16 \sin^2 \theta_W \cos^2 \theta_W},
\end{aligned} \tag{7.120}$$

where the electroweak form factors $F_{EW}^3(s)$ and $F_{EW}^f(s)$ are given by DIZET library and they correspond to electroweak vertex corrections.

The electroweak box diagrams need a more complicated treatment. In the Born spin amplitudes two products of the coupling constants are given by

$$\begin{aligned}
g_\lambda^{Z,e} g_{-\lambda}^{Z,f} &= (g_V^{Z,e} - \lambda g_A^{Z,e})(g_V^{Z,f} + \lambda g_A^{Z,f}), \\
g_\lambda^{Z,e} g_\lambda^{Z,f} &= (g_V^{Z,e} - \lambda g_A^{Z,e})(g_V^{Z,f} - \lambda g_A^{Z,f}).
\end{aligned} \tag{7.121}$$

Therefore the doubly-vector component arrives at

$$g_V^{Z,e} g_V^{Z,f} = \frac{4T_e^3 T_f^3 - 8T_e^3 Q_f F_{EW}^f(s) - 8T_f^3 Q_e F_{EW}^e(s) + 16Q_f^2 F_{EW}^{ef}(s, t)}{(16 \sin^2 \theta_W \cos^2 \theta_W)^2}, \tag{7.122}$$

where the new form factor $F_{EW}^{e,f}(s, t)$ corresponds to electroweak box diagrams and is angle-dependent.

7.2.5 One Real Photon

Next let us discuss of the $\hat{\beta}_1$ tensors corresponding to the emission of a single real photon with the tree-level case (zero virtual photons). We start with $O(\alpha^1)$ split amplitude from the single bremsstrahlung (including ISR and FSR).

The first-order, one-photon, ISR matrix element from the Feynman diagrams


is

$$\begin{aligned}
& \mathcal{M}_{1\{I\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} \\
&= eQ_e \bar{v}(p_b, \lambda_b) \mathbf{M}_{\{I\}} \frac{\not{p}_a + m - \not{k}_1}{-2k_1 p_a} \not{\epsilon}_{\sigma_1}^*(k_1) u(p_a, \lambda_a) \\
&+ eQ_e \bar{v}(p_b, \lambda_b) \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-\not{p}_b + m + \not{k}_1}{-2k_1 p_a} \mathbf{M}_{\{I\}} u(p_a, \lambda_a),
\end{aligned} \tag{7.123}$$

where

$$\mathbf{M}_{\{I\}} = ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) G_{e\mu}^B (G_{f,\nu}^B)_{[cd]} \tag{7.124}$$

is the annihilation scattering spinor matrix, including the final-state spinors. We split the above formula into the soft IR parts proportional to $(\not{p} \pm m)$ and the non-IR parts proportional to \not{k}_1 .



ISR diagrams.


Then we have, using completeness relation (D.5) in the Appendix (D),

$$\begin{aligned}
\mathcal{M}_{1\{I\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= -\frac{eQ_e}{2k_1p_a} \sum_{\rho} \mathfrak{B} \begin{bmatrix} p_b p_a \\ \lambda_b \rho_a \end{bmatrix}_{[cd]} U \begin{bmatrix} p_a k_1 p_a \\ \rho_a \sigma_1 \lambda_a \end{bmatrix} \\
&+ \frac{eQ_e}{2k_1p_a} \sum_{\rho} V \begin{bmatrix} p_b k_1 p_b \\ \lambda_b \sigma_1 \rho_b \end{bmatrix} \mathfrak{B} \begin{bmatrix} p_b p_a \\ \rho_b \lambda_a \end{bmatrix}_{[cd]} \\
&+ \frac{eQ_e}{2k_1p_b} \sum_{\rho} \mathfrak{B} \begin{bmatrix} p_b k_1 \\ \lambda_b \rho \end{bmatrix}_{[cd]} U \begin{bmatrix} k_1 k_1 p_a \\ \rho \sigma_1 \lambda_a \end{bmatrix} \\
&- \frac{eQ_e}{2k_1p_b} \sum_{\rho} V \begin{bmatrix} p_b k_1 k_1 \\ \lambda_b \sigma_1 \rho \end{bmatrix} \mathfrak{B} \begin{bmatrix} k_1 p_a \\ \rho \lambda_a \end{bmatrix}_{[cd]}. \tag{7.125}
\end{aligned}$$

The summation in the first two terms gets canceled by the diagonality property of U and V and leads to

$$\begin{aligned}
\mathcal{M}_{1\{I\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= \mathfrak{s}_{\sigma_1}^{\{I\}}(k_1) \mathfrak{B} \begin{bmatrix} p \\ \lambda \end{bmatrix} + r_{\{I\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix}, \\
r_{\{I\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= +\frac{eQ_e}{2k_1p_b} \sum_{\rho} \mathfrak{B} \begin{bmatrix} p_b k_1 \\ \lambda_b \rho \end{bmatrix}_{[cd]} U \begin{bmatrix} k_1 k_1 p_a \\ \rho \sigma_1 \lambda_a \end{bmatrix} \\
&- \frac{eQ_e}{2k_1p_b} \sum_{\rho} V \begin{bmatrix} p_b k_1 k_1 \\ \lambda_b \sigma_1 \rho \end{bmatrix} \mathfrak{B} \begin{bmatrix} k_1 p_a \\ \rho \lambda_a \end{bmatrix}_{[cd]}, \\
\mathfrak{s}_{\sigma_1}^{\{I\}}(k_1) &= -eQ_e \frac{b_{\sigma_1}(k_1, p_a)}{2k_1p_a} + eQ_e \frac{b_{\sigma_1}(k_1, p_b)}{2k_1p_b}. \tag{7.126}
\end{aligned}$$

The soft part is now separated and the remaining non-IR part for CEEX is obtained.



FSR diagrams.

The case of the final-state, one-real-photon emission can be analyzed in an analogous way. The first-order FSR, one-photon, matrix element reads

$$\begin{aligned} \mathcal{M}_{1\{F\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= eQ_f \bar{u}(p_c, \lambda_c) \not{\epsilon}_{\sigma_1}^*(k_1) \frac{\not{p}_c + m + \not{k}_1}{2k_1 p_c} \mathbf{M}_{\{F\}} v(p_d, \lambda_d) \\ &+ eQ_f \bar{u}(p_c, \lambda_c) \mathbf{M}_{\{F\}} \frac{-\not{p}_d + m - \not{k}_1}{2k_1 p_d} \not{\epsilon}_{\sigma_1}^*(k_1) v(p_d, \lambda_d), \end{aligned} \quad (7.127)$$

where

$$\mathbf{M}_{\{F\}} = ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) (G_{e,\mu}^B)_{[ba]} G_{f\nu}^B \quad (7.128)$$

is the annihilation scattering spinor matrix, including the initial spinors. Analogously, the expansion into soft and non-IR parts for the FSR spin amplitudes is obtained:

$$\begin{aligned} \mathcal{M}_{1\{F\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= \mathfrak{s}_{\sigma_1}^{\{F\}}(k_1) \mathfrak{B} \begin{bmatrix} p \\ \lambda \end{bmatrix} + r_{\{F\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix}, \\ r_{\{F\}} \begin{pmatrix} pk_1 \\ \lambda\sigma_1 \end{pmatrix} &= + \frac{eQ_f}{2k_1 p_c} \sum_{\rho} U \begin{bmatrix} p_c k_1 k_1 \\ \lambda_c \sigma_1 \rho \end{bmatrix} \mathfrak{B}_{[ba]} \begin{bmatrix} k_1 p_d \\ \rho \lambda_d \end{bmatrix} \\ &- \frac{eQ_f}{2k_1 p_d} \sum_{\rho} \mathfrak{B}_{[ba]} \begin{bmatrix} p_c k_1 \\ \lambda_c \rho \end{bmatrix} V \begin{bmatrix} k_1 k_1 p_d \\ \rho \sigma_1 \lambda_d \end{bmatrix}, \\ \mathfrak{s}_{\sigma_1}^{\{F\}}(k_1) &= -eQ_f \frac{b_{\sigma_1}(k_1, p_c)}{2k_1 p_c} + eQ_f \frac{b_{\sigma_1}(k_1, p_d)}{2k_1 p_d}. \end{aligned} \quad (7.129)$$

For the discussion of the remaining non-IR terms, it is useful to introduce a compact tensor notation:

$$U \begin{bmatrix} p_f k_i k_j \\ \lambda_f \sigma_i \sigma_f \end{bmatrix} \equiv U_{[f,i,j]}, \quad \mathfrak{B} \begin{bmatrix} p_b p_a \\ \lambda_a \lambda_b \end{bmatrix} \begin{bmatrix} p_c p_d \\ \lambda_c \lambda_d \end{bmatrix}, \quad (7.130)$$

$$U_{[a,i,j']} V_{[j',j,b]} \equiv \sum_{\sigma'_j = \pm} U \begin{bmatrix} p_a k_i k_j \\ \lambda_a \sigma_i \sigma'_j \end{bmatrix} V \begin{bmatrix} k_j k_j p_b \\ \sigma'_j \sigma_j \lambda_b \end{bmatrix}. \quad (7.131)$$

With the help of the above notation, the complete $O(\alpha^1)$ spin amplitudes for the one-photon ISR+FSR with explicit split into IR and non-IR parts, and ISR and FSR parts reads

$$\begin{aligned} \mathfrak{M}_1^{(1)} \begin{pmatrix} p k_1 \\ \lambda \sigma \end{pmatrix} &= \mathfrak{M}_{1\{I\}}^{(1)} \begin{pmatrix} p k_1 \\ \lambda \sigma \end{pmatrix} (P - k_1) + \mathfrak{M}_{1\{F\}}^{(1)} \begin{pmatrix} p k_1 \\ \lambda \sigma \end{pmatrix} (P) \\ &= \mathfrak{s}_{[1]}^{\{I\}} \mathfrak{B} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P - k_1 + r_I \begin{pmatrix} p k_1 \\ \lambda \sigma_1 \end{pmatrix} ; P - k_1 \\ &\quad + \mathfrak{s}_{[1]}^{\{F\}} \mathfrak{B} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P + r_F \begin{pmatrix} p k_1 \\ \lambda \sigma_1 \end{pmatrix} ; P, \end{aligned} \quad (7.132)$$

where

$$\begin{aligned} r_I \begin{pmatrix} p k_1 \\ \lambda \sigma_1 \end{pmatrix} ; X &= \frac{e Q_e}{2k p_a} \mathfrak{B}_{[b1'cd]}(X) U_{[1'1a]} - \frac{e Q_e}{2k p_b} V_{[b11']} \mathfrak{B}_{[1'acd]}(X) \\ r_F \begin{pmatrix} p k_1 \\ \lambda \sigma_1 \end{pmatrix} ; X &= \frac{e Q_f}{2k p_c} U_{[c11']} \mathfrak{B}_{[ba1'd]}(X) - \frac{e Q_f}{2k p_d} \mathfrak{B}_{[bac1']}(X) V_{[1'1d]}. \end{aligned} \quad (7.133)$$

In the lowest order, the Born spin amplitudes \mathfrak{B} are defined in eq. (7.50).

With the help of the $O(\alpha^1)$ variant of eq. (7.106) we are now ready to obtain

$$\begin{aligned}\hat{\beta}_{1\{I\}}^{(1)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P - k_1 \right) &\equiv r_{\{I\}} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P - k_1 \right), \\ \hat{\beta}_{1\{F\}}^{(1)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P \right) &\equiv r_{\{F\}} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P \right) \\ &+ \left(\frac{(p_c + p_d + k_1)^2}{(p_c + p_d)^2} - 1 \right) \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right).\end{aligned}\quad (7.134)$$

The total four-momentum in the resonance propagator X is uniquely define as $X = P - k_1$ in the case of ISR and $X = P$ in the case of FSR.

In order to obtain the $\hat{\beta}_1^{(2)}$, we have to deal with the non-trivial case of the simultaneous emission of virtual and real photons. Therefore it is instructive to write the formal definition of $\hat{\beta}_1^{(2)}$ in a particular case:

$$\begin{aligned}\mathfrak{M}_{1\{\omega\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X_\omega \right) &= \left\{ \exp[-\alpha B_4 - \alpha \Delta B_4^R(X_\omega)] \mathcal{M}_{1\{\omega\}}^{(2)} \right\} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X_\omega \right) \Big|_O (\alpha^2), \\ \omega = I, F, \quad R = \gamma, Z,\end{aligned}\quad (7.135)$$

$$\begin{aligned}\hat{\beta}_{1\{I\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P - k_1 \right) &= \mathfrak{M}_{1\{I\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P - k_1 \right) - \mathfrak{s}_{\sigma_1}^{\{I\}} \hat{\beta}_0^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; P - k_1 \right), \\ \hat{\beta}_{1\{F\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P - k_1 \right) &= \mathfrak{M}_{1\{F\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; P \right) - \mathfrak{s}_{\sigma_1}^{\{F\}} \hat{\beta}_0^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; P \right).\end{aligned}\quad (7.136)$$

For this moment, we have the amplitudes corresponding to vertexlike diagrams and we miss the diagrams of the "5-box" type. More precisely, after applying the IR

virtual subtraction in eq. (7.135) we expand in the number of loops,

$$\begin{aligned}
\mathfrak{M}_{1\{\omega\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) &= \mathfrak{M}_{1\{\omega\}}^{(1)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) + \alpha Q_e^2 \mathfrak{M}_{1\{\omega\},II}^{[1]} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) \\
&+ \alpha Q_f^2 \mathfrak{M}_{1\{\omega\},FF}^{[1]} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) + \alpha Q_e Q_f \mathfrak{M}_{1\{\omega\},\text{Box5}}^{[1]} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right).
\end{aligned} \tag{7.137}$$

In the above formula the first term corresponds the tree-level single bremsstrahlung, the next two terms correspond to the vertexlike diagrams, and the last one represents the "5-box"-type diagrams. The "5-box" term is given by

$$\begin{aligned}
\hat{\beta}_{1\{\omega\},\text{Box5}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) &= \alpha Q_e Q_f \mathfrak{M}_{1\{\omega\},\text{Box5}}^{[1]} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) \\
&- \mathfrak{s}_{[1]}^{\{I\}} \mathcal{R}_{\text{Box}}^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) - \mathfrak{s}_{[1]}^{\{F\}} \mathcal{R}_{\text{Box}}^{(1)} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right).
\end{aligned} \tag{7.138}$$

As we see, the trivial IR part is proportional to the ordinary box contribution mentioned before.

From the pure "vertexlike" diagrams for one real ISR photon we have the following $O(Q_e^2 \alpha^2)$ result:

$$\begin{aligned}
\hat{\beta}_{1\{I\}}^{(2)} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) &\equiv r_{\{I\}} \left(\begin{array}{c} pk_1 \\ \lambda\sigma_1 \end{array} ; X \right) [1 + \delta_{Virt}^{(1)e}(s) + \rho_{Virt}^{(2)e}(s, \tilde{\alpha}_1, \tilde{\beta}_1)] \\
&\times [1 + \delta_{Virt}^{(1)f}(s)] + \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X \right) \mathfrak{s}_{\sigma_1}^{\{I\}}(k_1) \rho_{Virt}^{(2)e}(s, \tilde{\alpha}, \tilde{\beta}) \tag{7.139}
\end{aligned}$$

where

$$\begin{aligned}\rho_{Virt}^{(2)e}(s, \tilde{\alpha}, \tilde{\beta}) &= \frac{\alpha}{\pi} Q_e^2 \frac{1}{2} [V(s, \tilde{\alpha}, \tilde{\beta}) + V(s, \tilde{\beta}, \tilde{\alpha})], \\ V(s, \tilde{\alpha}, \tilde{\beta}) &= \log(\tilde{\alpha}) \log(1 - \tilde{\beta}) + Li_2(\tilde{\alpha}) - \frac{1}{2} \log^2(1 - \tilde{\alpha}) \\ &\quad + \frac{3}{2} \log(1 - \tilde{\alpha}) + \frac{1}{2} \frac{\tilde{\alpha}(1 - \tilde{\alpha})}{[1 + (1 - \tilde{\alpha})^2]}\end{aligned}\quad (7.140)$$

and we use the Sudakov variables

$$\tilde{\alpha}_i = \frac{2k_i p_b}{2p_a p_b}, \quad \tilde{\beta}_i = \frac{2k_i p_a}{2p_a p_b}.\quad (7.141)$$

From eq. (7.139) we have several remarks:

The terms of $O(\alpha^4)$ like $|\mathbf{s}_\sigma^{\{I\}} \rho_{Virt}^{(2)e}|^2$ in the cross section are not rejected. They are included in the process of numerical evaluation of the differential cross sections out of spin amplitudes.

The term $r_{\{I\}} \delta_{Virt}^{(1)e}$ contributes to $O(\alpha^2 L^2)$ to the integrated cross section: one logarithm is explicit from the virtual photon and another is from the integration over the angle of the real photon.

The term $\sim \log(\tilde{\alpha}) \log(1 - \hat{\beta})$ contributes a correction of $O(\alpha^2 L^2)$ to the integrated cross section. The double logarithm comes directly from the integration over the angle of the real photon:

$$\int \frac{d^3 k}{k^0} \Re[\rho_{Virt}^{(2)e}(k) \{\hat{\beta}_0 \mathbf{s}_\sigma^{\{I\}}(k)\}^*] \sim Q_e^2 \alpha^2 \int_{\frac{m_e^2}{s}} \frac{d\tilde{\alpha}}{\tilde{\alpha}} \sim Q_e^2 \alpha^2 \log^2 \left(\frac{s}{m_e^2} \right).\quad (7.142)$$

Similarly, the $O(Q_f^2\alpha^2)$ contribution for one real FSR photon is

$$\begin{aligned}
\hat{\beta}_{1\{F\}}^{(2)} \begin{pmatrix} pk \\ \lambda\sigma \end{pmatrix} ; X &\equiv r_{\{F\}} \begin{pmatrix} pk \\ \lambda\sigma \end{pmatrix} ; X [1 + \delta_{Virt}^{(1)f}(s)][1 + \delta_{Virt}^{(1)f}(s) + \rho_{Virt}^{(2)f}(s, \tilde{\alpha}', \tilde{\beta}')] \\
&+ \mathfrak{B} \begin{pmatrix} \mathbf{p} \\ \lambda \end{pmatrix} ; X \mathfrak{s}_{\sigma_1}^{\{F\}}(k_1) \rho_{Virt}^{(2)f}(s, \tilde{\alpha}', \tilde{\beta}') \\
&+ \mathfrak{B} \begin{pmatrix} \mathbf{p} \\ \lambda \end{pmatrix} ; X \mathfrak{s}_{\sigma}^{\{F\}}(k) [1 + \delta_{Virt}^{(1)e}(s)][1 + \delta_{Virt}^{(1)f}(s)] \\
&\times \left(1 - \frac{(p_c + p_d + k)^2}{(p_c + p_d)^2} \right)
\end{aligned} \tag{7.143}$$

where

$$\begin{aligned}
\rho_{Virt}^{(2)f}(s, \tilde{\alpha}, \tilde{\beta}) &= \frac{\alpha}{\pi} Q_f^2 \frac{1}{4} \bar{L}_f [\log(1 - \tilde{\alpha}'') + \log(1 - \tilde{\beta}'')], \\
\tilde{\alpha}' &= \frac{2k_i p_b}{2p_a p_b}, \quad \tilde{\beta}' = \frac{2k_i p_a}{2p_a p_b}, \\
\tilde{\alpha}'' &= \frac{\tilde{\alpha}'}{1 + \tilde{\alpha}' + \tilde{\beta}'}, \quad \tilde{\beta}'' = \frac{\tilde{\beta}'}{1 + \tilde{\alpha}' + \tilde{\beta}'}.
\end{aligned} \tag{7.144}$$

In the above FSR amplitudes averaged over the photon angles, only the double logarithmic part is kept.

7.2.6 Two Real Photons

In the $O(\alpha^2)$, the contributions from two real photons are completely at the tree level without virtual corrections. There are three types of double bremsstrahlung: two ISR photons, two FSR photons and one ISR photon plus one FSR photon. In the following, the corresponding spin amplitudes will be given without any approximation, in particular we will not use the small-mass approximation $\frac{m_f^2}{s} \ll 1$.

(i) Two real ISR photons: The second-order, two-photon, ISR matrix element for the Feynman diagrams is given by

$$\begin{aligned}
& \mathcal{M}_{2\{II\}}^{(2)} \left(\begin{array}{c} p_a p_b k_1 k_2 \\ \lambda_a \lambda_b \sigma_1 \sigma_2 \end{array} ; P - k_1 - k_2 \right) \\
&= i e^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(P - k_1 - k_2) (G_{f,\nu}^B)_{[c,d]} (e Q_e)^2 \bar{v}(p_b, \lambda_b) \\
&\quad \times \left\{ G_{e,\mu}^B \frac{(\not{p}_a + m) - \not{k}_1 - \not{k}_2}{-2k_1 p_a - 2K_2 p_a + 2k_1 k_2} \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(\not{p}_a + m) - \not{k}_2}{-2k_2 p_a} \not{\epsilon}_{\sigma_2}^*(k_2) \right. \\
&\quad + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(-\not{p}_b + m) - \not{k}_1}{-2k_1 p_b} \not{\epsilon}_{\sigma_2}^*(k_2) \frac{(-\not{p}_b + m) + \not{k}_1 + \not{k}_2}{-2k_1 p_b - 2K_2 p_b + 2k_1 k_2} G_{e,\mu}^B \\
&\quad + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(-\not{p}_b + m) - \not{k}_1}{-2k_1 p_b} G_{e,\mu}^B \frac{(\not{p}_a + m) - \not{k}_2}{-2k_2 p_a} \not{\epsilon}_{\sigma_2}^*(k_2) \\
&\quad \left. + (1 \leftrightarrow 2) \right\} u(p_a, \lambda_a). \tag{7.145}
\end{aligned}$$

Using eq. (7.106), we have

$$\begin{aligned}
& \hat{\beta}_{2\{II\}}^{(2)} \left(\begin{array}{c} p k_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{array} ; P - k_1 - k_2 \right) = \mathfrak{M}_{2\{II\}}^{(2)} \left(\begin{array}{c} p k_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{array} ; P - k_1 - k_2 \right) \\
& - \hat{\beta}_{1\{I\}}^{(1)} \left(\begin{array}{c} p k_1 \\ \lambda \sigma_1 \end{array} ; P - k_1 - k_2 \right) \mathfrak{s}_{\sigma_2}^{\{I\}}(k_2) - \hat{\beta}_{1\{I\}}^{(1)} \left(\begin{array}{c} p k_2 \\ \lambda \sigma_2 \end{array} ; P - k_1 - k_2 \right) \mathfrak{s}_{\sigma_2}^{\{I\}}(k_1) \\
& - \hat{\beta}_0^{(0)} \left(\begin{array}{c} p \\ \lambda \end{array} ; P - k_1 - k_2 \right) \mathfrak{s}_{\sigma_1}^{\{I\}}(k_1) \mathfrak{s}_{\sigma_2}^{\{I\}}(k_2). \tag{7.146}
\end{aligned}$$

We will repeat what we did in the one-photon case: we isolate the group of terms containing two factors of $(\not{p} + m)$ from the above equation first, then isolate the group containing a single factor of $(\not{p} + m)$, and isolate the rest at last. Such a treatment will almost exactly split eq. (7.67) into a contribution with two \mathfrak{s} factors, a contribution with one single \mathfrak{s} factor, and the IR-finite remnant $\hat{\beta}_2^{(2)}$. In other words, we decompose

$\mathfrak{M}_{2\{II\}}^2$ into several terms as described above and then implement the IR subtraction of eq. (7.146) term by term.

Let us first discuss the doubly IR-divergent part proportional to two factors of $(\not{p} + m)$. In order to simplify the discussion, we omit the moment $2k_1k_2$ in the propagator. Using the completeness relations (D.5) and diagonality property (D.12) in the Appendix (D), we can factorize the soft factors exactly and completely

$$\begin{aligned}
& (eQ_e)^2 \bar{v}(p_b, \lambda_b) \left\{ G_{e,\mu}^B \frac{(\not{p}_a + m) - \not{k}_1 - \not{k}_2}{-2k_1p_a - 2K_2p_a + 2k_1k_2} \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(\not{p}_a + m) - \not{k}_2}{-2k_2p_a} \not{\epsilon}_{\sigma_2}^*(k_2) \right. \\
& + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(-\not{p}_b + m) - \not{k}_1}{-2k_1p_b} \not{\epsilon}_{\sigma_2}^*(k_2) \frac{(-\not{p}_b + m) + \not{k}_1 + \not{k}_2}{-2k_1p_b - 2K_2p_b + 2k_1k_2} G_{e,\mu}^B \\
& \left. + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(-\not{p}_b + m) - \not{k}_1}{-2k_1p_b} G_{e,\mu}^B \frac{(\not{p}_a + m) - \not{k}_2}{-2k_2p_a} \not{\epsilon}_{\sigma_2}^*(k_2) + (1 \leftrightarrow 2) \right\} u(p_a, \lambda_a) \\
& = (G_{e,\mu}^B)_{[ba]} (eQ_e)^2 \left\{ \frac{b_{\sigma_1}(k_1, p_a)}{2k_1p_a + 2K_2p_a} \frac{b_{\sigma_2}(k_2, p_a)}{2k_2p_a} + \frac{b_{\sigma_1}(k_1, p_b)}{2k_1p_b} \frac{b_{\sigma_2}(k_2, p_b)}{2k_1p_b + 2k_2p_b} \right. \\
& \quad \left. - \frac{b_{\sigma_1}(k_1, p_b)}{2k_1p_b} \frac{b_{\sigma_2}(k_2, p_a)}{2k_2p_a} + (1 \leftrightarrow 2) \right\} \\
& = (G_{e,\mu}^B)_{[ba]} \mathfrak{S}_{\sigma_1}^{[I]}(k_1) \mathfrak{S}_{\sigma_2}^{[I]}(k_2), \tag{7.147}
\end{aligned}$$

where the identity

$$\frac{1}{2k_1p_a + 2K_2p_a} \frac{1}{2k_1p_a} + \frac{1}{2k_1p_a + 2K_2p_a} \frac{1}{2k_1p_a} = \frac{1}{2k_1p_a} \frac{1}{2k_2p_a} \tag{7.148}$$

is applied.

If we restore the term $2k_1k_2$ in the propagator, $\mathcal{M}_{2\{II\}}^{\text{Double IR}}$ leads to

$$\begin{aligned}
\hat{\beta}_{2\{II\}}^{(2)\text{Double}} \begin{bmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{bmatrix} &= \mathcal{M}_{2\{II\}}^{(2)\text{Double}} \begin{bmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{bmatrix} - \mathfrak{s}_{\sigma_1}^{\{I\}}(k_1)\mathfrak{s}_{\sigma_2}^{\{I\}}(k_2)\mathfrak{B} \begin{bmatrix} p \\ \lambda \end{bmatrix} \\
&= (\mathfrak{s}_{[1]}^{(a)}\mathfrak{s}_{[2]}^{(a)}\Delta_a + \mathfrak{s}_{[1]}^{(b)}\mathfrak{s}_{[2]}^{(b)}\Delta_b)\mathfrak{B} \begin{bmatrix} p \\ \lambda \end{bmatrix}, \\
\mathfrak{s}_{\sigma_i}^{(a)}(k_i) \equiv \mathfrak{s}_{\sigma_i}^{(a)} &= -eQ_e \frac{b_{\sigma_1}(k_1, p_a)}{2k_i p_a}, \\
\mathfrak{s}_{\sigma_i}^{(b)}(k_i) \equiv \mathfrak{s}_{\sigma_i}^{(b)} &= -eQ_e \frac{b_{\sigma_1}(k_1, p_b)}{2k_i p_b}, \\
\mathfrak{s}_{\sigma_i}^{\{I\}}(k_i) \equiv \mathfrak{s}_{[i]}^{(a)} + \mathfrak{s}_{[i]}^{(b)} &= \mathfrak{s}_{\sigma_i}^{(a)}(k_i) + \mathfrak{s}_{\sigma_i}^{(b)}(k_i), \\
\Delta_f &= \frac{2k_1p_f + 2k_2p_f}{2k_1p_f + 2k_2p_f \mp 2k_1k_2} - 1 \\
&= \frac{\pm 2k_1k_2}{2k_1p_f + 2k_2p_f \mp 2k_1k_2}, \quad f = a, b, c, d. \tag{7.149}
\end{aligned}$$

Clearly, $\hat{\beta}^{(2)\text{Double}}$ is IR finite due to the Δ_f term. We introduced the compact notation above. From now on, for simplicity, we use the notation below

$$r_{if} = 2k_i \cdot p_f, \quad r_{ij} = 2k_i \cdot k_j, \quad f = a, b, c, d \quad i, j = 1, 2, \dots, n. \tag{7.150}$$

The next group of terms we are going to deal with is the one containing the single factor $(\not{p} + m)$. To be more specific, we will include terms that may result in a single IR divergence (if $k_1 \ll k_2$ or $k_2 \ll k_1$), namely, with $(\not{p} + m)$ next to a spinor,

at the end of the fermion line:

$$\begin{aligned}
\mathfrak{M}_{2\{II\}}^{\text{Single IR}} \begin{bmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{bmatrix} &= ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) (G_{f,\nu}^B)_{[c,d]} (eQ_e)^2 \bar{v}(p_b, \lambda_b) \\
&\times \left\{ G_{e,\mu}^B \frac{-k_1 - k_2}{-r_{1a} - r_{2a} + r_{12}} \not{\epsilon}_{\sigma_1}^*(k_1) \frac{\not{p}_a + m}{-r_{2a}} \not{\epsilon}_{\sigma_2}^*(k_2) \right. \\
&+ \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-\not{p}_b + m}{-r_{1b}} \not{\epsilon}_{\sigma_2}^* \frac{k_1 + k_2}{-r_{1b} - r_{2b} + r_{12}} G_{e,\mu}^B \\
&+ \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-\not{p}_b + m}{-r_{1b}} G_{e,\mu}^B \frac{-k_2}{-r_{2a}} \not{\epsilon}_{\sigma_2}^* \\
&\left. + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-k_1}{-r_{1b}} G_{e,\mu}^B \frac{-\not{p}_a + m}{-r_{2a}} \not{\epsilon}_{\sigma_2}^* + (1 \leftrightarrow 2) \right\} u(p_a, \lambda_a).
\end{aligned} \tag{7.151}$$

With the help of the compact notation, we can express $\mathcal{M}_{2\{II\}}^{\text{Single IR}}$ in a form which leads to an easy numerical calculation:

$$\begin{aligned}
\mathcal{M}_{2\{II\}}^{\text{Single IR}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} &= eQ_e \frac{-\mathfrak{B}_{[b1'] [cd]} U_{[1'1a]} - \mathfrak{B}_{[b2'] [cd]} U_{[2'1a]}}{-r_{1a} - r_{2a} + r_{12}} \mathfrak{s}_{[2]}^{(a)} \\
&+ eQ_e \mathfrak{s}_{[1]}^{(b)} \frac{V_{[b22']} \mathfrak{B}_{[2'a] [cd]} + V_{[b21']} \mathfrak{B}_{[1'a] [cd]}}{-r_{1a} - r_{2a} + r_{12}} \\
&- eQ_e \mathfrak{s}_{[1]}^{(b)} \mathfrak{B}_{[b2'] [cd]} \frac{U_{[2'2a]}}{-r_{2a}} + eQ_e \frac{V_{[b11']}}{-r_{1b}} \mathfrak{B}_{[1'a] [cd]} \mathfrak{s}_{[2]}^{(a)} \\
&+ (1 \leftrightarrow 2).
\end{aligned} \tag{7.152}$$

Moreover, the single-IR part to be eliminated is

$$\begin{aligned}
\hat{\beta}_{1(1)[1]}^{(1)} \mathfrak{s}_{[2]}^{\{I\}} + \hat{\beta}_{1(1)[2]}^{(1)} \mathfrak{s}_{[1]}^{\{I\}} &= r_{[1]}^{\{I\}} \mathfrak{s}_{[2]}^{\{I\}} + r_{[2]}^{\{I\}} \mathfrak{s}_{[1]}^{\{I\}} \\
&= \left(eQ_e \mathfrak{B}_{[b1'] [cd]} \frac{U_{[1'1a]}}{r_{1a}} - eQ_e \frac{V_{[b11']}}{r_{1a}} \mathfrak{B}_{[1'a] [cd]} \right) \mathfrak{s}_{[2]}^{\{I\}} \\
&+ (1 \leftrightarrow 2).
\end{aligned} \tag{7.153}$$

To sum up, we have

$$\begin{aligned}
\hat{\beta}_{2\{II\}}^{\text{Single}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} &= \mathcal{M}_{2\{II\}}^{\text{Single IR}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} - \hat{\beta}_{1(1)[1]}^{\{I\}} \mathfrak{s}_{[2]}^{\{I\}} - \hat{\beta}_{1(1)[2]}^{\{I\}} \mathfrak{s}_{[1]}^{\{I\}} \\
&= -eQ_e \mathfrak{B}_{[b2'][cd]} \frac{U_{[2'1a]}}{-r_{1a} - r_{2a} + r_{12}} \mathfrak{s}_{[2]}^{(a)} \\
&\quad + eQ_e \mathfrak{s}_{[1]}^{(b)} \frac{U_{[2'1a]}}{-r_{1a} - r_{2a} + r_{12}} \mathfrak{B}_{[1'a][cd]} \\
&\quad - eQ_e \mathfrak{B}_{[b1'][cd]} \left(\frac{U_{[1'1a]}}{-r_{1a} - r_{2a} + r_{12}} - \frac{U_{[1'1a]}}{-r_{1a}} \right) \mathfrak{s}_{[2]}^{(a)} \\
&\quad + eQ_e \mathfrak{s}_{[2]}^{(a)} \left(\frac{V_{[b22']}}{-r_{1a} - r_{2a} + r_{12}} - \frac{V_{[b22']}}{-r_{2b}} \right) \mathfrak{B}_{[2'a][cd]} \\
&\quad + (1 \leftrightarrow 2). \tag{7.154}
\end{aligned}$$

It is straightforward to find that the expression above is IR-finite.

At last, we need to include all the remaining terms from eq. (7.147). They are IR-finite and read

$$\begin{aligned}
\hat{\beta}_{2\{II\}}^{\text{Rest}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} &= ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) (G_{f,\nu}^B)_{[cd]} (eQ_e)^2 \bar{v}(p_b, \lambda_b) \\
&\quad \times \left\{ G_{e,\mu}^B \frac{(\not{p}_a + m) - \not{k}_1 - \not{k}_2}{-r_{1a} - r_{2a} + r_{12}} \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-\not{k}_2}{-r_{2a}} \not{\epsilon}_{\sigma_2}^*(k_2) \right. \\
&\quad + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-\not{k}_2}{-r_{1b}} \not{\epsilon}_{\sigma_2}^*(k_2) \frac{(\not{p}_b + m) + \not{k}_1 + \not{k}_2}{-r_{1b} - r_{2b} + r_{12}} G_{e,\mu}^B \\
&\quad \left. + \not{\epsilon}_{\sigma_1}^*(k_1) \frac{-\not{k}_2}{-r_{1b}} G_{e,\mu}^B \frac{-\not{k}_2}{-r_{2a}} \not{\epsilon}_{\sigma_2}^*(k_2) + (1 \leftrightarrow 2) \right\} u(p_a, \lambda_a). \tag{7.155}
\end{aligned}$$

With the help of tensor notation in the fermion helicity indices, we can expre the above equation in terms of the U and V matrices in the following way:

$$\begin{aligned}
& \hat{\beta}_{2\{II\}}^{\text{Rest}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} \\
&= (eQ_e)^2 \frac{\mathfrak{B}_{[ba'] [cd]} U_{[a'12'']} - \mathfrak{B}_{[b1'] [cd]} U_{[1'12'']} - \mathfrak{B}_{[b2'] [cd]} U_{[2'12'']} - U_{[2''2a]} - r_{1a} - r_{2a} + r_{12}}{-r_{1a} - r_{2a} + r_{12}} \\
&+ (eQ_e)^2 \frac{V_{[b11'']} - V_{[1''2b']} \mathfrak{B}_{[b'a] [cd]} + V_{[1''21']} \mathfrak{B}_{[1'a] [cd]} + V_{[1''22']} \mathfrak{B}_{[2'a] [cd]}}{-r_{1b} - r_{1b} - r_{2b} + r_{12}} \\
&+ (eQ_e)^2 \frac{V_{b11'}}{-r_{[1b]}} \mathfrak{B}_{[1'2'] [cd]} \frac{-U_{[2'2a]}}{-r_{2a}} + (1 \leftrightarrow 2). \tag{7.156}
\end{aligned}$$

Therefore the total ISR $\hat{\beta}_{2\{II\}}$ reads

$$\hat{\beta}_{2\{II\}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} = \hat{\beta}_{2\{II\}}^{\text{Double}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} + \hat{\beta}_{2\{II\}}^{\text{Single}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} + \hat{\beta}_{2\{II\}}^{\text{Rest}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix}. \tag{7.157}$$

(ii) Two real FSR photons: The case of double final-state real photon emission can be treated in an analogous way. The second order two FSR photon matrix element is

$$\begin{aligned}
& \mathcal{M}_{2\{FF\}}^{(2)} \begin{pmatrix} p_a p_b k_1 k_2 \\ \lambda_a \lambda_b \sigma_1 \sigma_2 \end{pmatrix}; P \\
&= i e^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(P) (G_{e,\mu}^B)_{[ba]} (eQ_e)^2 \bar{u}(p_c, \lambda_c) \\
&\times \left\{ \not{\epsilon}_{[1]}^*(k_1) \frac{(\not{p}_c + m) + \not{k}_1}{-2k_1 p_c} \not{\epsilon}_{[2]}^*(k_2) \frac{(\not{p}_c + m) + \not{k}_1 + \not{k}_2}{2k_1 p_c + 2K_2 p_c + 2k_1 k_2} G_{f,\mu}^B \right. \\
&+ G_{f,\mu}^B \frac{(-\not{p}_d + m) - \not{k}_1 - \not{k}_2}{2k_1 p_d + 2k_2 p_d + 2k_1 k_2} \not{\epsilon}_{\sigma_1}^*(k_1) \frac{(-\not{p}_d + m) - \not{k}_2}{-2k_2 p_d} \not{\epsilon}_{\sigma_2}^*(k_2) \\
&+ \not{\epsilon}_{[1]}^*(k_1) \frac{(\not{p}_c + m) + \not{k}_1}{2k_1 p_c} G_{f,\mu}^B \frac{(-\not{p}_d + m) - \not{k}_2}{2k_2 p_d} \not{\epsilon}_{[2]}^*(k_2) \\
&\left. + (1 \leftrightarrow 2) \right\} v(p_d, \lambda_d). \tag{7.158}
\end{aligned}$$

Analogously, the expansion into soft and non-IR parts for the FSR spin amplitudes is done in a completely similar way to the ISR case. The subtraction formula is

$$\begin{aligned}
\hat{\beta}_{2\{FF\}}^{(2)} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} ; P &= \mathfrak{M}_{2\{FF\}}^{(2)} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} ; P - \hat{\beta}_{1\{F\}}^{(1)} \begin{pmatrix} pk_1 \\ \lambda \sigma_1 \end{pmatrix} ; P \mathfrak{s}_{\sigma_2}^{\{F\}}(k_2) \\
&\quad - \hat{\beta}_{1\{F\}}^{(1)} \begin{pmatrix} pk_2 \\ \lambda \sigma_2 \end{pmatrix} ; P \mathfrak{s}_{\sigma_1}^{\{F\}}(k_1) \\
&\quad - \hat{\beta}_0^{(0)} \begin{pmatrix} p \\ \lambda \end{pmatrix} ; P \mathfrak{s}_{\sigma_1}^{\{F\}}(k_1) \mathfrak{s}_{\sigma_2}^{\{F\}}(k_2). \tag{7.159}
\end{aligned}$$

First we obtain the contribution from the group of terms containing two $(\not{p}-m)$ factors:

$$\begin{aligned}
&\hat{\beta}_{2\{FF\}}^{(2)\text{Double}} \begin{bmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{bmatrix} \\
&= \mathcal{M}_{2\{FF\}}^{(2)\text{Double}} \begin{bmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{bmatrix} - \mathfrak{s}_{[1]}^{\{F\}} \mathfrak{s}_{[2]}^{\{F\}} \mathfrak{B}_{[ba][cd]} \frac{(p_c + p_d + k_1 + k_2)^2}{(p_c + p_d)^2} \\
&= (\Delta_c \mathfrak{s}_{[1]}^{(a)} \mathfrak{s}_{[2]}^{(a)} + \Delta_d \mathfrak{s}_{[1]}^{(b)} \mathfrak{s}_{[2]}^{(b)}) \mathfrak{B}_{[ba][cd]} - \mathfrak{s}_{[1]}^{\{F\}} \mathfrak{s}_{[2]}^{\{F\}} \mathfrak{B} \begin{bmatrix} p_b p_a \\ \lambda_b \lambda_a \end{bmatrix} \\
&\quad \times \left(\frac{(p_c + p_d + k_1 + k_2)^2}{(p_c + p_d)^2} - 1 \right); \\
&\mathfrak{s}_{\sigma_i}^{(c)}(k_i) \equiv \mathfrak{s}_{[i]}^{(a)} = +eQ_f \frac{b_{\sigma_1}(k_i, p_c)}{r_{ic}}, \\
&\mathfrak{s}_{\sigma_i}^{(d)}(k_i) \equiv \mathfrak{s}_{[i]}^{(d)} = -eQ_f \frac{b_{\sigma_1}(k_i, p_d)}{2k_i p_d}, \\
&\mathfrak{s}_{\sigma_i}^{\{F\}}(k_i) \equiv \mathfrak{s}_{[i]}^{(c)} + \mathfrak{s}_{[i]}^{(d)} = \mathfrak{s}_{\sigma_i}^{(c)}(k_i) + \mathfrak{s}_{\sigma_i}^{(d)}(k_i), \tag{7.160}
\end{aligned}$$

which is explicitly finite. The group of terms containing only one $(\not{p} - m)$ factor at the end of the fermion line reads

$$\begin{aligned}
& \mathcal{M}_{2\{FF\}}^{\text{Single IR}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} \\
&= eQ_f \mathfrak{s}_{[1]}^{(c)} \frac{U_{[c21']}}{r_{1c} + r_{2c} + r_{12}} \mathfrak{B}_{[ba][1'd]} + eQ_f \mathfrak{s}_{[1]}^{(c)} \frac{U_{[c22']}}{r_{1c} + r_{2c} + r_{12}} \mathfrak{B}_{[ba][2'd]} \\
&+ eQ_f \mathfrak{B}_{[ba][c1']} \frac{-V_{[1'1d]}}{r_{1d} + r_{2d} + r_{12}} \mathfrak{s}_{[2]}^{(d)} + eQ_f \mathfrak{B}_{[ba][c2']} \frac{-V_{[2'1d]}}{r_{1d} + r_{2d} + r_{12}} \mathfrak{s}_{[2]}^{(d)} \\
&+ eQ_f \mathfrak{s}_{[1]}^{(c)} \mathfrak{B}_{[ba][c2']} \frac{-V_{[2'2d]}}{r_{2d}} + eQ_f \frac{U_{[c11']}}{r_{1c}} \mathfrak{B}_{[ba][1'd]} \mathfrak{s}_{[2]}^{(d)} + (1 \leftrightarrow 2). \tag{7.161}
\end{aligned}$$

Using the matrix notation (in the fermion spin indices), we have

$$\begin{aligned}
& \mathcal{M}_{2\{FF\}}^{\text{Single IR}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} \\
&= eQ_f \mathfrak{s}_{[1]}^{(c)} \frac{U_{[c21']}}{r_{1c} + r_{2c} + r_{12}} \mathfrak{B}_{[ba][1'd]} + eQ_f \mathfrak{s}_{[1]}^{(c)} \frac{U_{[c22']}}{r_{1c} + r_{2c} + r_{12}} \mathfrak{B}_{[ba][2'd]} \\
&+ eQ_f \mathfrak{B}_{[ba][c1']} \frac{-V_{[1'1d]}}{r_{1d} + r_{2d} + r_{12}} \mathfrak{s}_{[2]}^{(d)} + eQ_f \mathfrak{B}_{[ba][c2']} \frac{-V_{[2'1d]}}{r_{1d} + r_{2d} + r_{12}} \mathfrak{s}_{[2]}^{(d)} \\
&+ eQ_f \mathfrak{s}_{[1]}^{(c)} \mathfrak{B}_{[ba][c2']} \frac{-V_{[2'2d]}}{r_{2d}} + eQ_f \frac{U_{[c11']}}{r_{1c}} \mathfrak{B}_{[ba][1'd]} \mathfrak{s}_{[2]}^{(d)} + (1 \leftrightarrow 2). \tag{7.162}
\end{aligned}$$

Moreover, the single-IR part to be eliminated is

$$\begin{aligned}
& \hat{\beta}_{1(0)[1]}^{(1)} \mathfrak{s}_{[2]}^{\{F\}} + \hat{\beta}_{1(0)[1]}^{(1)} \mathfrak{s}_{[1]}^{\{F\}} \\
&= r_{[1]}^{\{F\}} r_{[2]}^{\{F\}} + r_{[2]}^{\{F\}} r_{[1]}^{\{F\}} \\
&= \left(+eQ_e \mathfrak{B}_{[ba][1'd]} \frac{U_{[c11']}}{r_{1c}} - eQ_e \frac{V_{[1'1d]}}{r_{1d}} \mathfrak{B}_{[ba][c1']} \right) \mathfrak{s}_{[2]}^{\{F\}} + (1 \leftrightarrow 2) \\
&- \mathfrak{B}_{[ba][cd]} \left(\frac{(p_c + p_d + k_1)^2}{(p_c + p_d)^2} - 1 \right) \mathfrak{s}_{[1]}^{\{F\}} \mathfrak{s}_{[2]}^{\{F\}} + (1 \leftrightarrow 2). \tag{7.163}
\end{aligned}$$

Then we obtain

$$\begin{aligned}
& \hat{\beta}_{2\{FF\}}^{\text{Single}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} \\
&= \mathcal{M}_{2\{FF\}}^{\text{Single IR}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} - \hat{\beta}_{1(0)}^{(1)} \begin{pmatrix} pk_1 \\ \lambda \sigma_1 \end{pmatrix} \mathfrak{s}^{\{F\}} \begin{bmatrix} k_2 \\ \sigma_2 \end{bmatrix} - \hat{\beta}_{1(0)}^{(1)} \begin{pmatrix} pk_2 \\ \lambda \sigma_2 \end{pmatrix} \mathfrak{s}^{\{F\}} \begin{bmatrix} k_1 \\ \sigma_1 \end{bmatrix} \\
&= eQ_f \mathfrak{s}_{[1]}^{(c)} \left\{ \left(\frac{U_{[c22']}}{r_{2c} + r_{1c} + r_{12}} - \frac{U_{[c22']}}{r_{2c}} \right) \mathfrak{B}_{[ba][2'd]} + \frac{U_{[c21']}}{r_{2c} + r_{1c} + r_{12}} \mathfrak{B}_{[ba][1'd]} \right\} \\
&\quad + eQ_f \left\{ \mathfrak{B}_{[ba][c1']} \left(\frac{-V_{[1'1d]}}{r_{1d} + r_{2d} + r_{12}} - \frac{-V_{[1'1d]}}{r_{1d}} \right) + \frac{-V_{2'1d}}{r_{1d} + r_{2d} + r_{12}} \mathfrak{B}_{[ba][c2']} \right\} \mathfrak{s}_{[2]}^{(d)} \\
&\quad + \mathfrak{B}_{[ba][cd]} \left(\frac{(p_c + p_d + k_1)^2}{(p_c + p_d)^2} - 1 \right) \mathfrak{s}_{[1]}^{\{F\}} \mathfrak{s}_{[2]}^{\{F\}} + (1 \leftrightarrow 2). \tag{7.164}
\end{aligned}$$

Finally the remaining term in eq. (7.158) reads

$$\begin{aligned}
& \mathcal{M}_{2\{FF\}}^{\text{Rest}} \begin{pmatrix} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{pmatrix} \\
&= i e^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) (G_{e,\mu}^B)_{[b,a]} (eQ_f)^2 \bar{u}(p_c, \lambda_c) \left\{ \not{\epsilon}_{[1]}^* \frac{\not{k}_1}{r_{1c}} \not{\epsilon}_{[2]}^* \frac{(\not{p}_c + m) + \not{k}_1 + \not{k}_2}{r_{1c} + r_{2c} + r_{12}} G_{f,\nu}^B \right. \\
&\quad \left. + G_{f,\nu}^B \frac{(-\not{p}_d + m) - \not{k}_1 - \not{k}_2}{r_{1d} + r_{2d} + r_{12}} \not{\epsilon}_{[1]}^* \frac{-\not{k}_2}{r_{2d}} \not{\epsilon}_{[2]}^* + \not{\epsilon}_{[1]}^* \frac{\not{k}_1}{r_{1c}} G_{f,\nu}^B \frac{-\not{k}_2}{r_{2d}} \not{\epsilon}_{[2]}^* + (1 \leftrightarrow 2) \right\} v(p_d, \lambda_d). \tag{7.165}
\end{aligned}$$

Thus, using the matrix notation, we obtain

$$\begin{aligned}
& \hat{\beta}_{2\{FF\}}^{\text{Rest}} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} \\
&= \mathcal{M}_{2\{FF\}}^{\text{Rest}} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} \\
&= (eQ_f)^2 \frac{U_{[c11'']}}{r_{1c}} \frac{U_{[1''2c']}\mathfrak{B}_{[ba][c'd]} + U_{[1''21']}\mathfrak{B}_{[ba][1'd]} + U_{[1''22']}\mathfrak{B}_{[ba][2'd]}}{r_{1c} + r_{2c} + r_{12}} \\
&\quad + (eQ_f)^2 \frac{-\mathfrak{B}_{[ba][cd']}V_{[d'12'']} - \mathfrak{B}_{[ba][c1']}V_{[1'12'']} - \mathfrak{B}_{[ba][c2']}V_{[2'12'']} - V_{[2''2d]}}{r_{1d} + r_{2d} + r_{12}} \frac{1}{r_{2d}} \\
&\quad + (eQ_f)^2 \frac{U_{[c11']}}{r_{1c}} \mathfrak{B}_{[ba][1'2']} \frac{-V_{[2'2d]}}{r_{2d}} + (1 \leftrightarrow 2). \tag{7.166}
\end{aligned}$$

The total contribution from the double FSR real photon emission reads

$$\hat{\beta}_{2\{FF\}} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} = \hat{\beta}_{2\{FF\}}^{\text{Double}} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} + \hat{\beta}_{2\{FF\}}^{\text{Single}} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix} + \hat{\beta}_{2\{FF\}}^{\text{Rest}} \begin{pmatrix} pk_1k_2 \\ \lambda\sigma_1\sigma_2 \end{pmatrix}. \tag{7.167}$$

(iii) One real ISR photon and one real FSR photon: Compared with cases described before, the case of one real ISR photon and one real FSR photon is easier since there is at most one photon on one fermion line:

$$\begin{aligned}
& \mathcal{M}_{2\{IF\}}^{(2)} \begin{pmatrix} p_a p_b p_c p_d k_1 k_2 \\ \lambda_a \lambda_b \lambda_c \lambda_d \sigma_1 \sigma_2 \end{pmatrix}; P - k_1 \\
&= ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(P - k_1) \\
&\quad \times eQ_e \bar{v}(p_b, \lambda_b) \left(G_{e,\mu}^B \frac{\not{p}_a - \not{k}_1 + m}{-2k_1 p_a} \not{\epsilon}_{[1]}^* + \not{\epsilon}_{[1]}^* \frac{-\not{p}_b + \not{k}_1 + m}{-2k_1 p_b} G_{e,\mu}^B \right) u(p_a, \lambda_a) \\
&\quad \times eQ_f \bar{u}(p_c, \lambda_c) \left(G_{f,\nu}^B \frac{-\not{p}_d - \not{k}_2 + m}{2k_2 p_d} \not{\epsilon}_{[2]}^* + \not{\epsilon}_{[2]}^* \frac{\not{p}_c + \not{k}_2 + m}{2k_2 p_c} G_{f,\nu}^B \right) v(p_d, \lambda_d) \tag{7.168}
\end{aligned}$$

and the IR subtraction is

$$\begin{aligned}
& \hat{\beta}_{2\{IF\}}^{(2)} \left(\begin{array}{c} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{array} ; P - k_1 \right) \\
&= \mathfrak{M}_{2\{IF\}}^{(2)} \left(\begin{array}{c} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{array} ; P - k_1 \right) - \hat{\beta}_{1\{I\}}^{(1)} \left(\begin{array}{c} pk_1 \\ \lambda \sigma_1 \end{array} ; P - k_1 \right) \mathfrak{s}_{\sigma_2}^F(k_2) \\
&\quad - \hat{\beta}_{1\{F\}}^{(1)} \left(\begin{array}{c} pk_2 \\ \lambda \sigma_2 \end{array} ; P - k_1 \right) \mathfrak{s}_{\sigma_1}^I(k_1) - \hat{\beta}_0^{(0)} \left(\begin{array}{c} p \\ \lambda \end{array} ; P - k_1 \right) \mathfrak{s}_{\sigma_1}^I(k_1) \mathfrak{s}_{\sigma_2}^I(k_2).
\end{aligned} \tag{7.169}$$

And $\hat{\beta}_{2\{IF\}}$ can be obtained by a simple subtraction of all terms proportional to one or two $(\not{p} - m)$ factors

$$\begin{aligned}
& \hat{\beta}_{2\{IF\}} \left(\begin{array}{c} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{array} ; X \right) \\
&= ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) e Q_e \bar{v}(p_b, \lambda_b) \left(G_{e,\mu}^B \frac{-\not{k}_1}{-r_{1a}} \not{\epsilon}_{[1]}^* + \not{\epsilon}_{[1]}^* \frac{\not{k}_1}{-r_{1b}} G_{e,\mu}^B \right) u(p_a, \lambda_a) \\
&\quad \times e Q_f \bar{u}(p_c, \lambda_c) \left(G_{f,\nu}^B \frac{-\not{k}_2}{r_{2d}} \not{\epsilon}_{[2]}^* + \not{\epsilon}_{[2]}^* \frac{\not{k}_2}{r_{2c}} G_{f,\nu}^B \right) v(p_d, \lambda_d).
\end{aligned} \tag{7.170}$$

In the programmable matrix notation it can be written as

$$\begin{aligned}
& \hat{\beta}_{2\{IF\}} \left(\begin{array}{c} pk_1 k_2 \\ \lambda \sigma_1 \sigma_2 \end{array} ; X \right) \\
&= ie^2 \sum_{B=\gamma, Z} \Pi_B^{\mu\nu}(X) e^2 Q_e Q_f \left((G_{e\mu}^B)_{b1'} \frac{-U_{[1'1a]}}{-r_{1a}} + \frac{V_{[b11']}}{-r_{1b}} (G_{e,\mu}^B)_{[1'a]} \right) \\
&\quad \times \left((G_{f,\nu}^B)_{[c2']} \frac{-V_{[2'2d]}}{r_{2d}} + \frac{U_{[c22']}}{r_{2c}} (G_{f,\nu}^B)_{[2'd]} \right) \\
&= e^2 Q_e Q_f \left(\mathfrak{B}_{[b1'][c2']}(X) \frac{-U_{[1'1a]}}{-r_{1a}} \frac{-V_{[2'2d]}}{r_{2d}} + \frac{U_{[c22']}}{r_{2c}} \mathfrak{B}_{[b1'][2'd]}(X) \frac{-U_{[1'1a]}}{-r_{1a}} \right. \\
&\quad \left. + \frac{V_{[b11']}}{-r_{1b}} \mathfrak{B}_{[1'a][c2']}(X) \frac{-V_{[2'2d]}}{r_{2d}} + \frac{V_{[b11']}}{-r_{1b}} \frac{U_{[c22']}}{r_{2c}} \mathfrak{B}_{[1'a][2'd]}(X) \right).
\end{aligned} \tag{7.171}$$

7.3 Relations between CEEEX and EEX

We have shown the EEX and CEEEX schemes in details in the last two sections. Next we shall compare certain important and interesting features of both schemes in more detail.

Let us first investigate the limit of the CEEEX in which we drop the dependence on the partition index $X_\varphi \rightarrow P$, where $P = p_a + p_b$. Note that there is no such analogy in the EEX. In this limit, for the simplest case of the $O(\alpha)$ exponentiation, we find

$$\sum_{\varphi \in \mathcal{P}} e^{\alpha B_4(X_\varphi)} \frac{X_\varphi^2}{s_{cd}} \mathfrak{B} \left(\begin{matrix} p \\ \lambda \end{matrix}; X_\varphi \right) \prod_{i=1}^n \mathfrak{s}_{[i]}^{\{\varphi_i\}} \Rightarrow e^{\alpha B_4(X_\varphi)} \mathfrak{B} \left(\begin{matrix} p \\ \lambda \end{matrix}; X_\varphi \right) \prod_{i=1}^n (\mathfrak{s}_{[i]}^{\{I\}} + \mathfrak{s}_{[i]}^{\{F\}}), \quad (7.172)$$

because of the relation (7.87). Note that the $\text{ISR} \otimes \text{FSR}$ interference contribution is preserved in the above transition.

Next we would like to discuss the case of the very narrow resonances, which the $\text{ISR} \otimes \text{FSR}$ interference contribution to any physical observable is so small that it can be neglected. This corresponds to a well-defined limit in the CEEEX scheme. In this limit, for the simplest case of the $O(\alpha^0)$ exponentiation we have

$$\begin{aligned} |\mathcal{M}_n^{(0)}|^2 &= \sum_{\varphi \in \mathcal{P}} \sum_{\varphi' \in \mathcal{P}} \exp[\alpha B_4(X_\varphi)] \exp[\alpha B_4(X_{\varphi'})]^* \\ &\quad \times \mathfrak{B} \left(\begin{matrix} p \\ \lambda \end{matrix}; X_\varphi \right) \mathfrak{B} \left(\begin{matrix} p \\ \lambda \end{matrix}; X_{\varphi'} \right)^* \prod_{i=1}^n \mathfrak{s}_{[i]}^{\{\varphi_i\}} \mathfrak{s}_{[j]}^{\{\varphi'_j\}} \\ &\Rightarrow \exp[2\alpha \Re B_2(p_a, p_b)] \exp[2\alpha \Re B_2(p_c, p_d)] \\ &\quad \times \sum_{\varphi \in \mathcal{P}} \left| \mathfrak{B} \left(\begin{matrix} p \\ \lambda \end{matrix}; X_\varphi \right) \right|^2 \prod_{i=1}^n |\mathfrak{s}_{[i]}^{\{\varphi_i\}}|^2. \end{aligned} \quad (7.173)$$

In the above transition we omit the $\text{ISR} \otimes \text{FSR}$ interference entirely, by dropping the nondiagonal terms $\wp \neq \wp'$ in the double summation over partitions, and replace the resonance form factor by the sum of the traditional YFS form factors for the ISR and the FSR. In this way, we find $O(\alpha^0)_{\text{EEX}}$ is identical to $O(\alpha^0)_{\text{CEEX}}$.

Last but not least, it is important to find out the relations between the CEEX $\hat{\beta}$'s defined at the amplitude level and the EEX $\bar{\beta}$'s defined at the differential distribution level. Let us suppress all spin indices, that is, for every term like $|\cdots|^2$ or $\Re[AB^*]$ the corresponding spin sum or average is taken. Then traditional $\bar{\beta}$'s of the EEX/YFS scheme at the $O(\alpha^2)$ level are

$$\begin{aligned}\bar{\beta}_0^{(l)} &= \left| \mathfrak{M}_0^{(l)} \right|_{(\alpha^l)}^2, \quad l = 0, 1, 2 \\ \bar{\beta}_1^{(l)}(k) &= \left| \mathfrak{M}_1^{(l)}(k) \right|_{(\alpha^{l+1})}^2 - \bar{\beta}_0^{(l)} |\mathfrak{s}(k)|^2, \quad l = 0, 1 \\ \bar{\beta}_2^{(2)}(k_1, k_2) &= \left| \mathfrak{M}_1^{(2)}(k_1, k_2) \right|_{(\alpha^{l+1})}^2 - \bar{\beta}_0^{(l)}(k_1) |\mathfrak{s}(k_2)|^2 - \bar{\beta}_0^{(l)}(k_2) |\mathfrak{s}(k_1)|^2 \\ &\quad - \bar{\beta}_0^{(0)} |\mathfrak{s}(k_1)|^2 |\mathfrak{s}(k_2)|^2,\end{aligned}\tag{7.174}$$

where the subscript $|_{(\alpha^r)}$ means a truncation to $O(\alpha^r)$. Now for each $\mathfrak{M}_n^{(n+l)}$, after substituting its expansion in terms of $\hat{\beta}$'s according to eq. (7.65), we have

$$\begin{aligned}\bar{\beta}_0^{(l)} &= \left| \hat{\beta}_0^{(l)} \right|_{(\alpha^l)}^2, \quad l = 0, 1, 2 \\ \bar{\beta}_1^{(l)}(k) &= \left| \hat{\beta}_1^{(l)}(k) \right|^2 + 2\Re[\hat{\beta}_0^{(l)} \hat{\beta}_1^{(l)*}(k)]_{(\alpha^{l+1})}, \quad l = 0, 1, \\ \bar{\beta}_2^{(2)}(k_1, k_2) &= \left| \hat{\beta}_2^{(2)}(k_1, k_2) \right|^2 + 2\Re\{[\hat{\beta}_1^{(1)}(k_1) \mathfrak{s}(k_2)][\hat{\beta}_1^{(1)}(k_2) \mathfrak{s}(k_1)]\}^* \\ &\quad + 2\Re\{\hat{\beta}_2^{(2)}(k_1, k_2)[\hat{\beta}_1^{(1)}(k_1) \mathfrak{s}(k_2) + \hat{\beta}_1^{(1)}(k_2) \mathfrak{s}(k_1)] \\ &\quad \times \hat{\beta}_1^{(0)} \mathfrak{s}(k_1) \mathfrak{s}(k_2)\}^*.\end{aligned}\tag{7.175}$$

The relation is not completely trivial for $O(\alpha^2)$, and there are some extra IR-finite terms on the right hand side. From the above analysis it is clear that $\bar{\beta}$'s are generally

more complicated objects than the $\hat{\beta}$'s. Moreover, in the $\bar{\beta}_0$ and $\bar{\beta}_1$ some higher-order virtual terms are unnecessarily truncated, which probably undermines the perturbative convergence of the EEX scheme in comparison with that of CEEX scheme. The above relation clearly exhibits the difference between the EEX and CEEX schemes.

In the above discussion, we show how the two examples of the EEX scheme can be obtained as a limit case of the CEEX, and show the exact relation between the $\bar{\beta}$'s of the EEX and the $\hat{\beta}$'s of the CEEX. From these it is clear that the CEEX scheme is more general than the EEX scheme.

7.4 Monte Carlo Algorithm

In this section we will introduce the Monte Carlo Algorithm for the KKMC, which generates final-state four momenta, i.e, points within the Lorentz invariant phase space according to eq. (7.2) for EEX and eq. (7.36) for CEEX. The MC technique of the KKMC is generally an approach of integrating exactly over the phase space without approximation. It is based on the rigorous perturbative quantum field theory: the differential cross section is the phase-space times the scattering amplitude for the corresponding Feynman diagrams. Furthermore, the KKMC is not only the phase-space integration but also the simulation of the actual scattering process, since it requires events (lists of four momenta) to be generated with weight equal to 1. Generally speaking, the MC algorithm includes a handful of elementary techniques such as weight-rejection, mapping and multibranching [115]. We will take the notation and terminology in Ref. [115]. In the KKMC, the self-adapting MC FOAM [102] is adopted as a building block, which works for arbitrary integrand distributions. In general, it is wise to minimize the use of the multibranching and utilize the method

of reweighting, constructing several layers of weights and taking their product as the total weight. In the KKMC, there are only three multibranchings, one for the types of the final fermion type $f = e, \mu, \tau, d, u, s, c, b$, another one for the photon partitions and the last one for helicities of the emitted photons.

In the following discussions, we will introduce the algorithm of the Monte Carlo generation of the events according to CEEEX and EEX differential distributions. The algorithm is constructed with elementary technique of MC simulation and multibranching with casual use of mapping (change of integration variables). The weights are products of several component weights ordered in a chain. Their job is to simplify the very complicated differential distributions so that we could integrate manually over certain integration variables. The remaining variables that we are not able to integrate will be dealt with the self-adapting MC generator FOAM. The procedure of simplifications mentioned above which involves with weights, multibranchings and mappings will be exhibited in the following subsections.

7.4.1 *Weights and Distributions*

First, let us describe the organization of the weights and distributions in KKMC. There are four principal distributions: pure phase space, model, crude and primary. Their ratios are the principal weight in KKMC.

The pure Lorentz-invariant phase space distribution given by eq. (7.2) is the basic reference differential distribution, of which the four-momentum conversation δ will be no generated directly in the MC, so that all other differential distributions of interest can be expressed in its terms

$$d\sigma(r_1, \dots, r_n) = \rho(r_1, \dots, r_n) d\tau_n(P; r_1, \dots, r_n), \quad (7.176)$$

where the density distribution, defined as follows:

$$\rho(r_1, \dots, r_n) = \frac{d\sigma(r_1, \dots, r_n)}{d\tau_n(P; r_1, \dots, r_n)}, \quad (7.177)$$

is analytical with no δ 's.

The model distribution is the density distribution corresponding to a physical model

$$\rho^{\text{Mod}}(r_1, \dots, r_n) = \frac{d\sigma^{\text{Mod}}(r_1, \dots, r_n)}{d\tau_{n+2}(P; r_1, \dots, r_n)}, \quad (7.178)$$

with which MC events will be generated.

The crude distribution is a density distribution

$$\rho^{\text{Cru}}(r_1, r_2, \dots, k_n) = \frac{d\sigma^{\text{Cru}}(r_1, r_2, k_1, \dots, k_n)}{d\tau_{n+2}(P; r_1, r_2, k_1, \dots, k_n)}. \quad (7.179)$$

It is close to all model distributions of a certain class and it should be maximally simple. And it should be Lorentz-invariant and be a maximally simple function of dot-products of the four-momentum. Here and later r_1, r_2 will denote the four-momenta of the outgoing fermions while k_i will denote the momenta of photons. In such cases the dimension of the phase-space will be explicitly $n + 2$.

The primary distribution

$$d\rho^{\text{Pri}}(\xi_1, \xi_2, \dots, \xi_n) \quad (7.180)$$

is defined primarily in the space Σ of variables ξ_i with the following properties: (a) the integral $\int d\rho^{\text{Pri}}(\xi_1, \xi_2, \dots, \xi_n)$ is known independently from analytical integration or an independent numerical integration of the Gauss type; (b) a well-defined mapping $r \rightarrow \xi$ exists. Therefore one can define

$$\rho^{\text{Pri}}(r_1, r_2, \dots, k_n) = \frac{d\rho^{\text{Pri}}(\xi_1, \xi_2, \dots, \xi_n)}{d\tau_{n+2}(P; r_1, r_2, k_1, \dots, k_n)}, \quad (7.181)$$

which is restricted to $\xi \in \Sigma_{\text{LIPS}}$ and is the distribution generated at the lowest level of the Monte Carlo. A zero weight will be assigned to the MC points (events) $\xi \notin \Sigma_{\text{LIPS}}$. The ρ^{Pri} relates to events generated according to $d\rho^{\text{Pri}}$ with all weights equal to 1.

The choice of the intermediate crude distribution, which stands between the primary and model distribution, depends on the practical need of modularity of the MC. For example we would like to use the same low-level MC event generator for both EEX and CEEX models. Undoubtedly we would like the MC event generator to have a well-defined low-level MC module. The weighted events are generated according to the crude distribution and the weight is

$$W^{\text{Cru}}(r_1, r_2, \dots, r_n) = \begin{cases} \frac{d\sigma^{\text{Cru}}(r_i(\xi_j))}{d\sigma^{\text{Pri}}(r_i)} = \frac{\rho^{\text{Cru}}(r_i)}{\rho^{\text{Pri}}(r_i)}, & \xi \in \Sigma_{\text{LIPS}}, \\ 0, & \xi \notin \Sigma_{\text{LIPS}}. \end{cases} \quad (7.182)$$

The above weight is determined by the low-level MC numerically without any further information on how the event (r_1, r_2, \dots, r_n) was actually generated.

The model weight for the m -th model is given by the ratio

$$W^{\text{Mod}}(r_1, r_2, \dots, r_n) = \frac{d\sigma_m^{\text{Mod}}(r_1, r_2, \dots, r_n)}{d\sigma^{\text{Cru}}(r_1, r_2, \dots, r_n)} = \frac{\rho^{\text{Mod}}(r_i)}{\rho^{\text{Cru}}(r_i)}, \quad (7.183)$$

which is evaluated in a separate module. And the crude distribution ρ^{Cru} is calculated locally in the corresponding module, using an analytical expression in terms of four-momenta of the event, and without any access to information from the lower-level MC. The total weight obviously reads

$$W_m^{\text{Tot}} = W^{\text{Cru}} W_m^{\text{Mod}}, \quad (7.184)$$

and the total cross section is given by

$$\sigma_m^{\text{Tot}} = \langle W_m^{\text{Tot}} \rangle \sigma^{\text{Pri}}. \quad (7.185)$$

After giving the formal expressions for weights and distributions, we need to define the crude differential distribution explicitly for both EEX and CEEX. We first define the crude differential distribution with respect to the standard Lorentz invariant phase space as follows:

$$\begin{aligned}
& \rho_{\dot{n},n'}^{\text{Cru}}(q_1, q_2; \dot{k}_1, \dots, \dot{k}_n; k'_1, \dots, l'_{n'}) \\
& \equiv \frac{d\sigma_{\text{Cru}}}{d\tau_{n+n'+2}(P; q_1, q_2, \dot{k}_1, \dots, \dot{k}_n, k'_1, \dots, k'_n)} \\
& = \frac{1}{\dot{n}!} \frac{1}{n'} \frac{\sigma_{\text{Born}}(s_X)}{4\pi} \frac{s_X}{s_Q} \frac{2}{\beta_f} \prod_{j=1}^{\dot{n}} 2\tilde{S}_e(\dot{k}_j) \bar{\Theta}_e(\dot{k}_j) e^{\gamma_e \log \epsilon_e} \prod_{l=1}^{n'} \tilde{S}_f(k'_l) \bar{\Theta}_f(k'_l) e^{\gamma_f \log \epsilon_f},
\end{aligned} \tag{7.186}$$

where

$$\epsilon_e = \frac{2E_{\min}}{\sqrt{2p_1 p_2}}, \quad \epsilon_f = \frac{2E'_{\min}}{\sqrt{2q_1 q_2}}, \quad \beta_f = \sqrt{1 - \frac{4m_f^2}{s_q}}, \quad s_Q = 2q_1 q_2 + 2m_f^2.$$

and γ_e and γ_f are given by eqs. (7.9) and (7.10) respectively. The infrared and collinear singularities are in the soft factors \tilde{S} . The $\sigma_{\text{Born}}(s_X)$ has a resonance peak at s_X . The flux factor $\frac{s_Q}{s_X}$ is from the $O(\alpha^1)$ QED matrix element, and it can also be obtained from the leading-log approximation at any order. Note that the above crude distribution is only for EEX. It would only fit one single partition in CEEX.

Thus, according to the discussions above, we could express the differential distribution as follows:

$$\begin{aligned}
& \rho(p_c, p_d, k_1, \dots, k_n) \\
&= \frac{d\sigma^{(0)}}{dLIPS_{n+2}(P; p_c, p_d, k_1, \dots, k_n)} \\
&= \frac{1}{n!} \frac{e^{Y(\Omega; p_a, \dots, p_d)} \bar{\Theta}(\Omega)}{flux(s)} \frac{1}{4} \sum_{\sigma_i \neq \mp 1} \sum_{\lambda_i = \mp 1} \mathfrak{M}_n^{(0)} \left(\begin{array}{c} pk_1 k_2 k \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \\
& \quad \times \left[\mathfrak{M}_n^{(0)} \left(\begin{array}{c} pk_1 k_2 k \dots k_n \\ \lambda \sigma_1 \sigma_2 \dots \sigma_n \end{array} \right) \right]^* \\
&= \frac{e^Y}{4s} \Theta \frac{1}{n!} \sum_{\sigma_i, \lambda_i} \sum_{\{\varphi\}} \sum_{\{\varphi'\}} \left[\prod_{i=1}^n \mathfrak{s}_{[i]}^{\varphi_i} \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array}; X_{\varphi_i} \right) \frac{X_{\varphi_i}^2}{s''} \right] \left[\prod_{j=1}^n \mathfrak{s}_{[j]}^{\varphi'_j} \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array}; X_{\varphi_j} \right) \frac{X_{\varphi_j}^2}{s''} \right]^*
\end{aligned} \tag{7.187}$$

where $s'' = (p_c + p_d)^2$. In the crude distribution we would like to neglect IFI. That means we need to drop non-diagonal terms $\varphi' \neq \varphi$. Additionally, the YFS form factor needs to be simplified to preserve IR cancellation

$$Y(\Omega; p_a, \dots, p_d) \rightarrow \gamma_e \log \epsilon_e + \gamma_f \log \epsilon_f.$$

Therefore we have

$$\begin{aligned}
& \rho(p_c, p_d, k_1, \dots, k_n) \\
&= \frac{1}{n!} \frac{1}{4s} \exp(\gamma_e \log \epsilon_e + \gamma_f \log \epsilon_f) \sum_{\varphi} \prod_{i=1}^n \bar{\Theta}(k_i) \sum_{\sigma_i} \left| \mathfrak{s}_{[i]}^{\varphi_i} \right|^2 \sum_{\lambda_i} \left| \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array}; X_{\varphi_j} \right) \right|^2 \frac{X_{\varphi_i}^4}{(s'')^2}.
\end{aligned} \tag{7.188}$$

We can identify

$$\sum_{\sigma_i} \left| \mathfrak{s}_{[i]}^{\omega_i} \right|^2 = -8\pi^3 \tilde{S}_\omega(k_i), \quad \tilde{S}_1(k_i) \equiv \tilde{S}_I(k_i), \quad \tilde{S}_0(k_i) \equiv \tilde{S}_F(k_i),$$

and the Born-like differential cross section

$$\sum_{\lambda_i} \left| \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X_{\varphi_j} \right) \right|^2 \frac{X_{\varphi_i}^2}{s''} \sim \frac{d\sigma_{\text{Born}}}{d\Omega}(s, s'', t, u, t', u', X_{\varphi_i}^2),$$

which is dependent on $s = 2p_a p_b$, $s'' = 2p_c p_d$, $t = -2p_a p_c$, $t' = -2p_b p_d$, $u = -2p_a p_d$, $u' = -2p_b p_c$ and $X_{\varphi_i}^2$ in the Z resonance propagator. Let us convert it into an "angular average" expression

$$\sum_{\lambda_i} \left| \mathfrak{B} \left(\begin{array}{c} p \\ \lambda \end{array} ; X_{\varphi_j} \right) \right|^2 \frac{X_{\varphi_i}^2}{s''} \rightarrow \frac{\sigma_{\text{Born}}(X_{\varphi_i}^2)}{4\pi},$$

Finally, the crude distribution for CEEEX is defined as follows:

$$\begin{aligned} & \rho_{[n]}^{\text{Cru}}(p_c, p_d, k_1, \dots, k_n) \\ &= \frac{d\sigma_{\text{CEEEX}}^{\text{Cru}}}{d\tau_{n+2}(P; p_c, p_d, k_1, \dots, k_n)} \\ &= \frac{1}{n!} \sum_{\{\varphi\}} \frac{1}{s} \exp(\gamma_e \log \epsilon_e + \gamma_f \log \epsilon_f) \frac{\sigma_{\text{Born}}(X_{\varphi_i}^2)}{4\pi} \frac{X_{\varphi_i}^2}{s''} \frac{2}{\beta_f} \prod_{i=1}^n \bar{\Theta}(k_i) \tilde{S}_{\varphi_i}(k_i). \end{aligned} \tag{7.189}$$

For arbitrary photon multiplicity we have the following relations between crude distributions for CEEEX and EEX

$$\rho_{[n]}^{\text{Cru}}(k_1, \dots, k_n) = \sum_{\dot{n}+\dot{n}'=n} \rho_{[\dot{n}, \dot{n}']}^{\text{Cru}}(k_1, \dots, k_{\dot{n}}, k_1, \dots, k_{\dot{n}'}). \tag{7.190}$$

The model weight for the $O(\alpha^{(r)})$ EEX reads

$$W_{\text{EEX}}^{(r)}(q_1, q_2; \dot{k}_1, \dots, \dot{k}_{\dot{n}}; \dot{k}'_1, \dots, \dot{k}'_{\dot{n}'}) = \frac{\rho_{\text{EEX}}^{(r)}(p_1, p_2, q_1, q_2; \dot{k}_1, \dots, \dot{k}_{\dot{n}}; \dot{k}'_1, \dots, \dot{k}'_{\dot{n}'})}{\rho_{[\dot{n}, \dot{n}']}^{\text{Cru}}(q_1, q_2; \dot{k}_1, \dots, \dot{k}_{\dot{n}}; \dot{k}'_1, \dots, \dot{k}'_{\dot{n}'})}, \tag{7.191}$$

where the model distribution in the numerator is given by eq. (7.4) and the crude distribution in the denominator is given by eq. (7.186). And the model weight for the $O(\alpha^{(r)})$ CEEEX reads

$$W_{\text{CEEEX}}^{(r)}(p_a, p_b, p_c, p_d; k_1, \dots, k_n) = \frac{\rho_{\text{CEEEX}}^{(r)}(p_a, p_b, p_c, p_d; k_1, \dots, k_n)}{\rho_{[\tilde{n}, \tilde{n}']}^{\text{Cru}}(p_c, p_d; k_1, \dots, k_n)(2\pi)^{3(n+2)-4}}, \quad (7.192)$$

where the model distribution in the numerator is given by eq. (7.37) and the crude distribution in the denominator is given by eq. (7.189). Note that the factor $(2\pi)^{3(n+2)-4}$ is derived from the difference in the normalization of $d\text{LIPS}_n$ and $d\tau_n$.

Therefore, according to the previous subsection the corresponding total weight is

$$\begin{aligned} & W_{\text{CEEEX}}^{(r)\text{Tot}}(p_a, p_b, p_c, p_d; k_1, \dots, k_n) \\ &= W_{\text{CEEEX}}^{(r)}(p_a, p_b, p_c, p_d; k_1, \dots, k_n) W^{\text{Cru}}(p_a, p_b, p_c, p_d; k_1, \dots, k_n) \end{aligned} \quad (7.193)$$

and

$$\begin{aligned} & W_{\text{EEEX}}^{(r)\text{Tot}}(q_1, q_2; \dot{k}_1, \dots, \dot{k}_n; k'_1, \dots, k'_{n'}) \\ &= W_{\text{EEEX}}^{(r)}(q_1, q_2; \dot{k}_1, \dots, \dot{k}_n; k'_1, \dots, k'_{n'}) W^{\text{Cru}}(q_1, q_2; \dot{k}_1, \dots, \dot{k}_n; k'_1, \dots, k'_{n'}) \end{aligned} \quad (7.194)$$

where W^{Cru} is exactly the same since we did the proper Bosen-Einstein symmetrization for CEEEX. Among these model weights, only one can be used as the principal weight for a rejection of the events. Obviously we choose the best one, $O(\alpha^{(2)})$ CEEEX-type.

7.4.2 Phase-space Reorganization

Let us start with rewriting the phase space integral of eq. (7.186) for the crude total cross section as follows

$$\begin{aligned}
\sigma^{\text{Cru}} &= \int ds_x \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \int d\tau_{n+1}(P; k_1, \dots, k_n, X) \frac{1}{n!} \prod_{j=1}^n \tilde{S}_e(k_j) \bar{\Theta}_e(k_j) \\
&\times \int d\tau_{n'+2}(X; k'_1, \dots, k'_n, q_1, q_2) \frac{1}{n!} \prod_{j=1}^n \tilde{S}_f(k'_j) \bar{\Theta}_f(k'_j) \\
&\times \frac{\sigma_{\text{Born}}(s_X)}{4\pi} \frac{s_X}{s_Q} \frac{2}{\beta_f} \exp(\gamma_e \log \epsilon_e + \gamma_f \log \epsilon_f)
\end{aligned} \tag{7.195}$$

where $P = p_1 + p_2$. The integral above is Lorentz-invariant and can be computed in any reference frame. So we can take advantage of the Lorentz invariance of $d\tau_{n'+2}(X; k'_1, \dots, k'_n, q_1, q_2)$ and we transform all its variables to the reference frame where $X = \hat{X} = (\sqrt{s_X}, 0, 0, 0)$, the XMS frame, and rewrite eq. (7.195) as follows,

$$\begin{aligned}
\sigma^{\text{Cru}} &= \int ds_x \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \int d\tau_{n+1}(P; k_1, \dots, k_n, X) \frac{1}{n!} \prod_{j=1}^n \tilde{S}_e(k_j) \bar{\Theta}_e(k_j) \\
&\times \int d\tau_{n'+2}(\bar{X}; \bar{k}'_1, \dots, \bar{k}'_n, \bar{q}_1, \bar{q}_2) \frac{1}{n!} \prod_{j=1}^n \tilde{S}_f(\bar{k}'_j) \bar{\Theta}_f(\bar{k}'_j) \\
&\times \frac{\sigma_{\text{Born}}(s_X)}{4\pi} \frac{s_X}{s_Q} \frac{2}{\beta_f} \exp(\gamma_e \log \epsilon_e + \gamma_f \log \epsilon_f),
\end{aligned} \tag{7.196}$$

where those variables with a bar are defined in XMS. So far this operation is still ambiguous. We have to write down explicitly the Lorentz transformation L_X from XMS to CMS and back. Here we apply a so-called parallel boost B_X along the direction of the \vec{X} in PMS (a laboratory frame where $\vec{P} = 0$ and $p_1 = (p^0, 0, 0, p^3)$).

The corresponding transformation matrix is

$$B_X = \begin{bmatrix} \frac{X^0}{M_X}, & \frac{\vec{X}^T}{M_X} \\ \frac{\vec{X}}{M_X}, & I + \frac{\vec{X} \otimes \vec{X}}{M_X(M_X + X^0)} \end{bmatrix}, \quad X^2 = M_X^2, \tag{7.197}$$

where T denotes the matrix transposition and \otimes denotes the tensor product. The transformation from the XMS to CMS is

$$k'_i|_{\text{CMS}} = L_X \bar{k}'_i, \quad q_i|_{\text{CMS}} = L_X \bar{q}_i, \quad L_X = B_X. \quad (7.198)$$

The emission of the FSR photons is done in the comoving frame attached to the momenta q_i of outgoing fermions, namely, in the frame where $\vec{Q} = \vec{q}_1 + \vec{q}_2 = 0$ and $q_1 = (q_1^0, 0, 0, |q_1^3|)$, which is called QMS. In order to get from XMS to QMS we must know k'_i . This problem can be solved by reparametrization the FSR integral with the help of the integration over the Lorentz group [116]. Applying the result of Ref. [116], we have

$$\begin{aligned} \sigma^{\text{Cru}} &= \int ds_X \sum_{n=0}^{\infty} \frac{1}{n!} \prod_{j=1}^n \frac{d^3 k_j}{2k_j^0} 2\tilde{S}_e(k_j) \bar{\Theta}_e(k_j) \delta\left(s_X - \left(P - \sum_{j=0}^n k_j\right)^2\right) e^{\gamma_e \log \epsilon_e} \\ &\times \int d\psi d\cos\omega \frac{\sigma_{\text{Born}}(s_X)}{4\pi} \sum_{n'=0}^{\infty} \frac{1}{n'!} \int ds_Q \prod_{l=1}^{n'} \frac{d^3 \tilde{k}'_l}{2\tilde{k}'_l{}^0} \tilde{S}_f(\tilde{k}'_l) \bar{\Theta}_f(\tilde{k}'_l) \\ &\times \delta\left(s_X - \left(\hat{Q} - \sum_{j=0}^{n'} \tilde{k}'_j\right)^2\right) e^{\gamma_f \log \epsilon_f}, \end{aligned} \quad (7.199)$$

where those variables with a tilde are defined in QMS. Note that the Jacobian from the reparametrization of the FSR integral cancels exactly the factor $\frac{s_X}{s_Q} \frac{2}{\beta_f}$. The explicit transformation from QMS to XMS defines the new integration variables ψ and ω :

$$\bar{k}_i = L_A k_i, \quad \bar{q}_i = L_A \hat{q}_i, \quad L_A = R_3(\psi) R_2(\omega) B_{\hat{X}}^{-1}, \quad \hat{X} = \hat{Q} - \sum \tilde{k}'_j. \quad (7.200)$$

Notice that the explicit integration over q_1 and q_2 has disappeared completely after the operation above, which leads a great simplification of the crude integral. Note that ψ and ω are not polar angles of a certain momentum in a certain frame but parameters in the Lorentz transformation.

Therefore the crude integral can be rewritten in the following way:

$$\begin{aligned}
\sigma^{\text{Cru}} &= \sum_{f=\mu,\tau,u,s,c,b} \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \int d\tau_{n+n'+2}(P; q_1, q_2, k_1, \dots, k_n, k'_1, \dots, k'_{n'}) \\
&\quad \times \rho_{[n,n']}^{\text{Cru}}(q_1, q_2; k_1, \dots, k_n; k'_1, \dots, k'_{n'}) \\
&= \sum_{f=\mu,\tau,u,s,c,b} \int ds_X \sigma_{\text{Born}}^f(s_X) \int d\psi \frac{d\cos\omega}{4\pi} \\
&\quad \times \sum_{n=0}^{\infty} \frac{1}{n!} \prod_{j=1}^n \frac{d^3k_j}{2k_j^0} 2\tilde{S}_e(k_j) \bar{\Theta}_e(k_j) \delta\left(s_X - \left(P - \sum_{j=0}^n k_j\right)^2\right) e^{\gamma_e \log \epsilon_e} \\
&\quad \times \sum_{n'=0}^{\infty} \frac{1}{n'!} \int ds_Q \prod_{l=1}^{n'} \frac{d^3\tilde{k}'_l}{2\tilde{k}'_l{}^0} \tilde{S}_f(\tilde{k}'_l) \bar{\Theta}_f(\tilde{k}'_l) \delta\left(s_X - \left(\hat{Q} - \sum_{j=0}^{n'} \tilde{k}'_j\right)^2\right) e^{\gamma_f \log \epsilon_f},
\end{aligned} \tag{7.201}$$

Obviously this factorizes into independent ISR and FSR parts. The above integral is ready for the MC generation.

7.4.3 MC generation of the FSR photon momenta

Next we will describe the MC algorithm for the generation of the FSR photon momenta. Let us consider FSR part of the crude integral of eq. (7.201)

$$\mathfrak{F}_{n'} = \frac{1}{n'!} \int_{4m_f^2}^{s_X} ds_Q \prod_{j=1}^{n'} \int \frac{d^3\tilde{k}'_j}{\tilde{k}'_j{}^0} \Theta(\tilde{k}'_j - E'_{\min}) \delta\left(s_X - \left(\hat{Q} + \sum_{l=0}^{n'} \tilde{k}'_l\right)^2\right) e^{\gamma_f \log \epsilon_f}, \tag{7.202}$$

where

$$\begin{aligned}
\gamma_f &= Q_f^2 \frac{\alpha}{\pi} \frac{1 + \beta_f^2}{\beta_f} \left(\log \frac{1 + \beta_f}{1 - \beta_f} - 1 \right) = Q_f^2 \frac{\alpha}{\pi} \frac{1 + \beta_f^2}{\beta_f} \left(\log \frac{(1 + \beta_f)^2}{\mu_f^2} - 1 \right), \\
\beta_f &= \sqrt{1 - \mu_f^2}, \quad \mu_f^2 = \frac{4m_f^2}{s_Q}, \quad \epsilon_f = \frac{2E'_{\min}}{\sqrt{s_Q}}, \quad \hat{Q} = (\sqrt{s}, 0, 0, 0)
\end{aligned} \tag{7.203}$$

where we restored finite fermion mass m_f , photon momenta \tilde{k}'_l in the QMS rest frame of the outgoing fermions and E'_{\min} is the minimum energy of the real photon in this frame. Let us express the photon momenta in units of $\frac{1}{2}\sqrt{s_Q}$ and introduce polar

parametrization and other auxiliary notation:

$$\begin{aligned}\tilde{k}'_j &\equiv \frac{\sqrt{s_Q}}{2} \bar{k}_j \equiv \frac{\sqrt{s_Q}}{2} x_j (1, \sin \theta_j \cos \phi_j, \sin \theta_j \sin \phi_j, \cos \theta_j), \\ \tilde{K}' &= \sum_{l=0}^{n'} \tilde{k}'_l \equiv \frac{\sqrt{s_Q}}{2} \bar{K}.\end{aligned}\tag{7.204}$$

Then the δ -function can be eliminated:

$$\begin{aligned}\int_{4m_f^2}^{s_X} ds_Q \delta\left(s_X - \left(\hat{Q} + \sum_{l=0}^{n'} \tilde{k}'_l\right)^2\right) &= \int_{4m_f^2}^{s_X} ds_Q \delta\left(s_X - s_Q \left(1 + \bar{K}^0 + \frac{1}{4} \bar{K}^2\right)\right) \\ &= \frac{\Theta(s_Q(\bar{k}_1, \dots, \bar{k}_{n'} - 4m_f^2))}{1 + \bar{K}^0 + \frac{1}{4} \bar{K}^2},\end{aligned}\tag{7.205}$$

and from now on

$$s_Q = s_Q(\bar{k}_1, \dots, \bar{k}_{n'}) = \frac{s_X}{1 + \bar{K} + \frac{1}{4} \bar{K}^2}.\tag{7.206}$$

And the single-photon distribution is transformed as follows:

$$\begin{aligned}\frac{d^3 \tilde{k}'_j}{\tilde{k}'_j{}^0} \tilde{S}_f(\tilde{k}'_j) &= \frac{dx_j}{x_j} \frac{d\phi_j}{2\pi} d\cos \theta_j \frac{\alpha}{\pi} f\left(\theta_j, \frac{m_f^2}{s_Q}\right), \\ f\left(\theta_j, \frac{m_f^2}{s_Q}\right) &= \frac{1 + \beta_f^2}{\delta_{1j} \delta_{2j}} - \frac{\mu_f^2}{2} \frac{1}{\delta_{1j}^2} - \frac{\mu_f^2}{2} \frac{1}{\delta_{2j}^2}, \\ \delta_{1j} &= 1 - \beta_f \cos \theta_j, \quad \delta_{2j} = 1 + \beta_f \cos \theta_j.\end{aligned}\tag{7.207}$$

and the whole FSR integral is transformed into the semi-factorized expression:

$$\begin{aligned}\mathfrak{F}_{n'} &= \frac{1}{n!} \prod_{j=1}^{n'} \int_{\epsilon_f}^{\infty} \frac{dx_j}{x_j} \int_0^{2\pi} \frac{d\phi_j}{2\pi} \int_{-1}^{+1} d\cos \theta_j \frac{\alpha}{\pi} f\left(\theta_j, \frac{m_f^2}{s_Q}\right) \\ &\quad \times \frac{\Theta(s_Q - 4m_f^2)}{1 + \bar{K} + \frac{1}{4} \bar{K}^2} e^{\gamma_f \log \epsilon_f}.\end{aligned}\tag{7.208}$$

Note that the integral above is not factorized into a product of independent integrals since the dependence on all photon momenta \bar{k}_j is entering everywhere through the variable s_Q . So we call it semi-factorized.

Moreover, the introduction of the factor $\frac{1}{2}\sqrt{s_Q}$ leads another problem: the upper bound of x_j extends to a large values but not really to infinity due to the

$\Theta(s_Q - 4m_f^2)$. And this problem can be solved by the following change of variables:

$$\begin{aligned} y_i &= \frac{x_i}{1 + \sum x_j}, & x_i &= \frac{y_i}{1 - \sum y_j}, \\ 1 + \sum_j x_j &= \frac{1}{1 - \sum y_j} = 1 + \bar{K}^0 = 1 + \frac{2K' \cdot Q}{s_Q} = \frac{s_X}{s_Q} \left(1 - \frac{K'^2}{s_X}\right), \end{aligned} \quad (7.209)$$

which leads to

$$\begin{aligned} \mathfrak{F}_{n'} &= \frac{1}{n'!} \prod_{j=1}^{n'} \int_{\frac{\epsilon_f}{1+\bar{K}^0}}^1 \frac{dy_j}{y_j} \int_0^{2\pi} \frac{d\phi_j}{2\pi} \int_{-1}^{+1} d\cos\theta_j \frac{\alpha}{\pi} f\left(\theta_j, \frac{m_f^2}{s_Q}\right) \\ &\quad \times \frac{1 + \bar{K}^0}{1 + \bar{K}^0 + \frac{1}{4}\bar{K}^2} \Theta(s_Q - 4m_f^2) e^{\gamma_f \log \epsilon_f}. \end{aligned} \quad (7.210)$$

With the help of new variables the condition $s_Q > 4m_f^2$ (easily executable in the MC) translates approximately into $\sum_j y_j < 1$. Then we have

$$\frac{1 + \bar{K}^0}{1 + \bar{K}^0 + \frac{1}{4}\bar{K}^2} \leq 1,$$

which is perfect for the MC. However, the new IR limit $y_j > \epsilon_f/(1 + \bar{K}^0)$ is inconvenient for the MC. This issue can be solved by substituting

$$\epsilon_f = \delta_f(1 + \bar{K}^0). \quad (7.211)$$

where $\delta_f \ll 1$ is the new IR regulator for the FSR real photon. Note that this gives a new lower bound for the photon energy in the QMS:

$$E''_{\min} = \delta_f \frac{1}{2} \sqrt{s_Q} (1 + \bar{K}^0) = \delta_f \frac{1}{2} \sqrt{s_Q} \left(1 + \frac{2K' \cdot \hat{Q}}{s_Q}\right), \quad (7.212)$$

which is higher than the previous $E'_{\min} = \frac{1}{2} \sqrt{s_Q} \delta_f$. Therefore, we must keep the value of δ_f very low.

So far the FSR integral (7.202) has been transformed without any approximations and the integrals were conveniently parametrized for the MC generation:

$$\begin{aligned} \mathfrak{F}_{n'} &= \frac{1}{n'!} \prod_{j=1}^{n'} \int_{\delta_f}^1 \frac{dy_j}{y_j} \int_0^{2\pi} \frac{d\phi_j}{2\pi} \int_{-1}^{+1} d\cos\theta_j \frac{\alpha}{\pi} f\left(\theta_j, \frac{m_f^2}{s_Q}\right) \\ &\times \frac{1 + \bar{K}^0}{1 + \bar{K}^0 + \frac{1}{4}\bar{K}^2} \Theta(s_Q - 4m_f^2) e^{\gamma_f \log(\delta_f(1+\bar{K}^0))}. \end{aligned} \quad (7.213)$$

There is a one-to-one correspondence between the points in the Lorentz-invariant phase space and the points in space of the new variables:

$$\{n', (\tilde{k}'_1, \dots, \tilde{k}'_{n'})\} \leftrightarrow \{n', (y_j, \theta_j, \phi_j), \quad j = 1, \dots, n'\}. \quad (7.214)$$

Besides, we can write explicitly the differential distributions in the two equivalent parametrizations

$$\begin{aligned} &\frac{d\mathfrak{F}_{n'}}{ds_Q \delta(s_X - (\hat{Q} + \sum_{l=0}^{n'} \tilde{k}'_l)^2) \prod_{j=1}^{n'} \frac{d^3\tilde{k}'_j}{2\tilde{k}'_j}} \\ &= \frac{\Theta(s_Q - 4m_f^2)}{n'!} \exp\left[\gamma_f \log\left(\frac{2E''_{\min}}{\sqrt{s_X}}\right)\right] \prod_{j=1}^{n'} 2\tilde{S}_f(\tilde{k}'_j) \Theta(\hat{k}'_j - E''_{\min}) \end{aligned} \quad (7.215)$$

$$\begin{aligned} &\frac{d\mathfrak{F}_{n'}}{\prod_{j=1}^{n'} dy_j d\cos\theta_j d\phi_j} \\ &= \frac{\Theta(s_Q - 4m_f^2)}{n'!} e^{\gamma_f \log(\delta_f(1+\bar{K}^0))} \left(\frac{\alpha}{2\pi^2}\right)^{n'} \prod_{j=1}^{n'} \frac{\Theta(y_j - \delta_f)}{y_j} f\left(\theta_j, \frac{m_f^2}{s_Q}\right). \end{aligned} \quad (7.216)$$

Now it is time to introduce the simplifications that lead to a primary distribution. And the primary distribution can be integrated analytically and generated

using standard uniform random numbers. The simplifications are given as

$$\begin{aligned}
f\left(\theta_j, \frac{m_f^2}{s_Q}\right) &\rightarrow \bar{f}\left(\theta_j, \frac{m_f^2}{s_X}\right) = \frac{1 + \bar{\beta}_f^2}{\bar{\beta}_f} \frac{1}{1 - \bar{\beta}_f^2 \cos^2 \theta_j}, \\
\frac{1 + \bar{K}^0}{1 + \bar{K}^0 + \frac{1}{4}\bar{K}^2} \Theta(s_Q - 4m_f^2) &\rightarrow 1, \\
e^{\gamma_f \log(\delta_f(1+\bar{K}^0))} &\rightarrow e^{\bar{\gamma}_f \log \delta_f},
\end{aligned} \tag{7.217}$$

where

$$\bar{\beta}_f = \left[1 - \left(\frac{m_f^2}{s_X}\right)^2\right]^{\frac{1}{2}}, \quad \bar{\gamma}_f = Q_f^2 \frac{\alpha}{\pi} \frac{1 + \bar{\beta}_f^2}{\bar{\beta}_f} \log \frac{1 + \bar{\beta}_f}{1 - \bar{\beta}_f}. \tag{7.218}$$

With the help of the simplifications above we could remove any complicated dependence on the momenta of all photons through s_Q , replacing s_Q with s_X . Then hard FSR photons get stronger collinear peaks at $\cos \theta_j = \pm 1$ in the primary differential distribution. Thus the FSR primary differential distribution is:

$$\frac{d\mathfrak{F}_{n'}^{\text{Pri}}}{\prod_{j=1}^{n'} dy_j d \cos \theta_j d \phi_j} = e^{\bar{\gamma}_f \log(\delta_f)} \left(\frac{\alpha}{2\pi^2}\right)^{n'} \prod_{j=1}^{n'} \frac{\Theta(y_j - \delta_f)}{y_j} \bar{f}\left(\theta_j, \frac{m_f^2}{s_X}\right), \tag{7.219}$$

and the compensating weight which transforms the primary distribution into the crude distribution is

$$w_{\text{FSR}}^{\text{Cru}} = \frac{d\mathfrak{F}_{n'}}{d\mathfrak{F}_{n'}^{\text{Pri}}} = \frac{1 + \bar{K}^0}{1 + \bar{K}^0 + \frac{1}{4}\bar{K}^2} e^{\gamma_f \log(\delta_f(1+\bar{K}^0)) - \bar{\gamma}_f \log(\delta_f)} \prod_{j=1}^{n'} \frac{f\left(\theta_j, \frac{m_f^2}{s_Q}\right)}{\bar{f}\left(\theta_j, \frac{m_f^2}{s_X}\right)}. \tag{7.220}$$

Events $\{n', (y_j, \cos \theta_j, \phi_j), \quad j = 1, \dots, n'\}$ generated according to $d\mathfrak{F}_{n'}^{\text{Pri}}$, defined in eq. (7.221) with weight $w_{\text{FSR}}^{\text{Cru}}$, will be distributed according to the differential distribution (7.216).

Finally we conclude that the integral over the FSR primary distribution can be evaluated analytically:

$$\begin{aligned}
\sum_{n'=0}^{\infty} \mathfrak{F}_{n'}^{\text{Pri}} &= \sum_{n'=0}^{\infty} \frac{1}{n'!} \prod_{j=1}^{n'} \int_{\delta_f}^1 \frac{dy_j}{y_j} \int_0^{2\pi} \frac{d\phi_j}{2\pi} \int_{-1}^{+1} d \cos \theta_j \frac{\alpha}{\pi} \bar{f} \left(\theta_j, \frac{m_f^2}{s_X} \right) e^{\bar{\gamma}_f \log(\delta_f)} \\
&= \sum_{n'=0}^{\infty} e^{-\bar{\gamma}_f \log(1/\delta_f)} \frac{1}{n'!} \left(\bar{\gamma}_f \log \frac{1}{\delta_f} \right)^{n'} \\
&= \sum_{n'=0}^{\infty} e^{-\langle n' \rangle} \frac{\langle n' \rangle^{n'}}{n'!} = 1.
\end{aligned} \tag{7.221}$$

The photon multiplicity for the primary distribution is the standard Poisson distribution, with the average

$$\langle n' \rangle = \bar{\gamma}_f \log \frac{1}{\delta_f}, \tag{7.222}$$

and the overall normalization is equal to 1.

The MC generation of the distribution (7.219) is fully factorized, and the variables $\cos \theta_j$, ϕ_j and y_j can be generated independently. The distribution of ϕ_j is flat and the distribution of y_j is trivial to generate

$$\phi_j = 2\pi r_{1j}, \quad y_j = \delta_f^{r_{2j}}, \tag{7.223}$$

where r_{ij} are the standard uniform random numbers $0 < r_{ij} < 1$. The distribution of $\cos \theta_j$ needs using the branching approach: it is split into two components

$$\frac{2}{1 - \bar{\beta}_f \cos^2 \theta_j} = \frac{1}{1 - \bar{\beta}_f \cos \theta_j} + \frac{1}{1 + \bar{\beta}_f \cos \theta_j}, \tag{7.224}$$

and $\cos \theta_j$ is generated according to one component, chosen with the equal odd between these two. For instance, if we choose the first component as $1/(1 - \bar{\beta}_f \cos \theta_j)$, then

$$\cos \theta_j = \frac{1}{\bar{\beta}_j} \left\{ 1 - (1 + \bar{\beta}_f) \left(\frac{1 - \bar{\beta}_f}{1 + \bar{\beta}_f} \right)^{r_{3j}} \right\}, \tag{7.225}$$

where r_{3j} is another uniform random number.

Next we introduce the MC algorithm of the generation of the ISR photon momenta. Let us begin with the ISR part of the crude integral (7.201) for one final fermion type f

$$\begin{aligned} \mathfrak{I}_n &= \frac{1}{n!} \int ds_X \sigma_{\text{Born}}^f(s_X) \prod_{j=1}^n \int \frac{d^3 k_j}{k_j^0} \tilde{S}_e(k_j) \Theta(k_j^0 - E_{\min}) \\ &\quad \times \delta\left(s_X - \left(P - \sum_{j=0}^n k_j\right)^2\right) e^{\gamma_e \log \epsilon_e}, \end{aligned} \quad (7.226)$$

where $E_{\min} = \epsilon_e \frac{1}{2} \sqrt{s}$ is the minimum energy of the real ISR photon in the laboratory CMS. We first introduce the variable $v = 1 - \frac{s_X}{s}$ and order energies of the photons

$$\begin{aligned} \mathfrak{I}_n &= \int_0^{v_{\max}} dv \sigma_{\text{Born}}^f(s(1-v)) \prod_{j=1}^n \int \frac{d^3 k_j}{k_j^0} \tilde{S}_e(k_j) \\ &\quad \times \Theta(k_1^0 - k_2^0) \Theta(k_2^0 - k_3^0) \dots \Theta(k_n^0 - E_{\min}) \delta\left(v - \frac{2KP - K^2}{s}\right) e^{\gamma_e \log \epsilon_e}, \end{aligned} \quad (7.227)$$

where $K = \sum_{j=0}^n k_j$ and $v_{\max} = 1 - \frac{4m_f^2}{s}$. Then we rescale all momenta and introduce a polar parametrization

$$k_i = \eta \bar{k}_i = \eta x_i (1, \sin \theta_i \sin \phi_i, \sin \theta_i \cos \phi_i, \cos \theta_i); \quad (7.228)$$

We fix the scaling factor η so that $\bar{k}_1^0 = x_1 = v$:

$$\begin{aligned} \mathfrak{I}_n &= \int d\eta \delta\left(\eta - \frac{k_1^0}{v}\right) \int_0^{v_{\max}} dv \sigma_{\text{Born}}^f(s(1-v)) \prod_{j=1}^n \frac{d^3 k_j}{k_j^0} \tilde{S}_e(k_j) \\ &\quad \times \Theta(k_1^0 - k_2^0) \Theta(k_2^0 - k_3^0) \dots \Theta(k_n^0 - E_{\min}) \delta\left(v - \frac{2KP - K^2}{s}\right) e^{\gamma_e \log \epsilon_e} \\ &= \int_0^{v_{\max}} dv \sigma_{\text{Born}}^f(s(1-v)) \prod_{j=1}^n \int_0^1 \frac{dx_j}{x_j} \int_0^{2\pi} \frac{d\phi_j}{2\pi} \int_{-1}^{+1} d \cos \theta_j \frac{\alpha}{\pi} f(\cos \theta_j) \\ &\quad \times \delta(v - x_1) \Theta(x_1 - x_2) \Theta(x_2 - x_3) \dots \Theta(\lambda_0 x_n - \epsilon) e^{\gamma_e \log \epsilon_e} \mathcal{J}(\bar{K}, v), \end{aligned} \quad (7.229)$$

where η_0 is the root of the equation $v - \frac{2\bar{K}P}{s}\eta + \frac{\bar{K}^2}{s}\eta = 0$ and $\mathcal{J}(\bar{K}, v)$ is an overall Jacobian factor:

$$\begin{aligned}\mathcal{J}(\bar{K}, v) &= \frac{v}{\eta_0} \frac{1}{\frac{2\bar{K}P}{s} - 2\eta_0 \frac{\bar{K}^2}{s}} = \frac{1}{2} \left(1 + \frac{1}{\sqrt{1 - Av}} \right), \\ \eta_0 &= \frac{\sqrt{s}}{2} \frac{v}{\bar{K}^0} \frac{2}{1 + \sqrt{1 - Av}} \equiv \frac{\sqrt{s}}{2} \lambda_0, \\ A &= \frac{\bar{K}^2 P^2}{(\bar{K}P)^2} = \frac{\bar{K}^2}{(\bar{K}^0)^2} \leq 1, \quad 0 \leq \lambda_0 \leq 1,\end{aligned}\tag{7.230}$$

and the photon angular distribution is determined by

$$f(\cos \theta_j) = \frac{2}{(1 - \beta \cos \theta_j)(1 + \beta \cos \theta_j)} - \frac{2m_e^2}{s} \frac{1}{(1 - \beta \cos \theta_j)^2} - \frac{2m_e^2}{s} \frac{1}{(1 + \beta \cos \theta_j)^2}.\tag{7.231}$$

Up till now, the ISR integral (7.226) has been transformed without any approximation and there is a one-to-one correspondence of the points in the Lorentz-invariant phase space and the points in the space of new variables:

$$\{n, (\tilde{k}_1, \dots, \tilde{k}_n)\} \leftrightarrow \{n, (y_j, \theta_j, \phi_j), \quad j = 1, \dots, n\}.\tag{7.232}$$

Analogously, we can write the differential distributions in two equivalent parametrizations of the IRS crude differential distribution:

$$\begin{aligned}\frac{d\mathcal{J}_n}{ds_X \prod_{j=1}^n \frac{d^3 k_j}{2k_j^0}} &= \frac{1}{n!} \sigma_{\text{Born}}^f(s_X) \prod_{j=1}^n 2\tilde{S}_e(k_j) \Theta(k_j^0 - E_{\min}) e^{\gamma_e \log \epsilon_e} \quad n > 0 \\ \frac{d\mathcal{J}_n}{dv \prod_{j=1}^n dx_j d \cos \theta_j d \phi_j} &= \sigma_{\text{Born}}^f(s(1-v)) \left(\frac{\alpha}{2\pi^2} \right)^n \delta(v - x_1) \frac{\Theta(\lambda_0 x_n - \epsilon)}{x_n} \\ &\times \prod_{j=1}^{n-1} \frac{\Theta(x_j - x_{j-1})}{x_j} \prod_{j=1}^n f(\cos \theta_j) e^{\gamma_e \log \epsilon_e} \mathcal{J}(\bar{K}, v), \quad n > 0, \\ \frac{d\mathcal{J}_0}{ds_X} &= \sigma_{\text{Born}}^f(s) \delta(s_X), \quad \frac{d\mathcal{J}_0}{dv} = \sigma_{\text{Born}}^f(s) \delta(s_X), \quad n = 0.\end{aligned}\tag{7.233}$$

Now we are ready to introduce the simplification leading to the ISR primary differential distribution:

$$\begin{aligned}
f(\cos \theta_j) &\rightarrow \bar{f}(\cos \theta_j) = \frac{2}{(1 - \cos \theta_j)(1 + \cos \theta_j)}, \\
\mathcal{J}(\bar{K}, v) &\rightarrow \mathcal{J}_0(v) = \frac{1}{2} \left(1 + \frac{1}{\sqrt{1-v}} \right), \\
\Theta(\lambda_0 x_n - \epsilon) &\rightarrow \Theta(x_n - \epsilon),
\end{aligned} \tag{7.234}$$

where

$$\bar{\gamma}_e = 2 \frac{\alpha}{\pi} \log \left(\frac{s}{m_e^2} \right). \tag{7.235}$$

Thus, the ISR primary differential distribution reads

$$\begin{aligned}
\frac{d\mathcal{J}_n}{dv \prod_{j=1}^n dx_j d \cos \theta_j d\phi_j} &= \sigma_{\text{Born}}^f(s(1-v)) \left(\frac{\alpha}{2\pi^2} \right)^n \delta(v - x_1) \frac{\Theta(\lambda_0 x_n - \epsilon)}{x_n} \\
&\times \prod_{j=1}^{n-1} \frac{\Theta(x_j - x_{j-1})}{x_j} \prod_{j=1}^n f(\cos \theta_j) e^{\gamma_e \log \epsilon_e} \mathcal{J}_0(v), \quad n > 0, \\
\frac{d\mathcal{J}_0^{\text{Pri}}}{dv} &= \sigma_{\text{Born}}^f \delta(v), \quad n = 0,
\end{aligned} \tag{7.236}$$

and the corresponding weight is

$$w_{\text{ISR}}^{\text{Cru}} = \frac{d\mathcal{J}_n}{d\mathcal{J}_n^{\text{Pri}}} = \Theta(\lambda_0 x_n - \epsilon) \frac{\mathcal{J}(\bar{K}, v)}{\mathcal{J}_0(v)} \prod_{j=1}^n \frac{f(\cos \theta_j)}{\bar{f}(\cos \theta_j)}. \tag{7.237}$$

Note that the $\Theta(\lambda_0 x_n - \epsilon)$ contribution to the weight leads directly to a characteristic factor $F(\gamma_e) = e^{-C\gamma_e} / \Gamma(1 + \gamma_e)$ [88, 117]

$$f(\cos \theta_j) = \frac{2 \sin^2 \theta_j}{[(1 - \beta \cos \theta_j)(1 + \beta \cos \theta_j)]^2}, \tag{7.238}$$

Finally, we can integrate analytically the ISR primary differential distribution

$$\begin{aligned}
\mathfrak{J}^{\text{Pri}} &= \sum_{n=0}^{\infty} \mathfrak{J}_n^{\text{Pri}} \\
&= \sum_{n=0}^{\infty} \int_0^{v_{\text{max}}} dv \sigma_{\text{Born}}^f(s(1-v)) \prod_{j=1}^n \int_0^1 \frac{dx_j}{x_j} \int_0^{2\pi} \frac{d\phi_j}{2\pi} \int_{-1}^{+1} d\cos\theta_j \\
&\quad \times \frac{\alpha}{\pi} \bar{f}(\cos\theta_j) \delta(v-x_1) \Theta(x_1-x_2) \Theta(x_2-x_3) \dots \Theta(\lambda_0 x_n - \epsilon) e^{\gamma_e \log \epsilon_e} \mathcal{J}_0(v), \\
&= \int_0^{v_{\text{max}}} dv \sigma_{\text{Born}}^f(s(1-v)) \mathcal{J}_0(v) e^{\gamma_e \log \epsilon_e} \\
&\quad \times \left(\delta(v) + \Theta(v-\epsilon) \frac{1}{v} \sum_{n=1}^{\infty} \frac{1}{(n-1)!} \left(\bar{\gamma}_e \log \frac{v}{\epsilon} \right)^{n-1} \right) \\
&= \int_0^{\epsilon} dv \gamma_e v^{\gamma_e-1} \sigma_{\text{Born}}^f(s) + \int_{\epsilon}^{v_{\text{max}}} dv \sigma_{\text{Born}}^f(s(1-v)) \mathcal{J}_0(v) \bar{\gamma}_e v^{\bar{\gamma}_e-1} \epsilon^{\gamma_e-\bar{\gamma}_e}.
\end{aligned} \tag{7.239}$$

For the generation of the primary differential distribution $d\mathfrak{J}^{\text{Pri}}$, we start with the generation of v according to

$$\frac{d\mathfrak{J}^{\text{Pri}}}{dv} = \sigma_{\text{Born}}^f(s(1-v)) \mathcal{J}_0(v) \bar{\gamma}_e v^{\bar{\gamma}_e-1} \epsilon^{\gamma_e-\bar{\gamma}_e}, \tag{7.240}$$

which is done by using FOAM. Photon multiplicity n is generated in the next step.

For $v < \epsilon$ we have simply

$$\mathfrak{J}_n^{\text{Pri}} = \text{const} \times \frac{1}{(n-1)!} \left(\bar{\gamma}_e \log \frac{v}{\epsilon} \right)^{n-1}, \tag{7.241}$$

which is just the shifted-by-one Poisson distribution P_{n-1} , with the average $\langle n-1 \rangle = \bar{\gamma}_e \log \frac{v}{\epsilon}$. The angles $\cos\theta_j$ and ϕ_j are generated in the same way as in the case of FSR.

7.4.4 Common IR Boundary For ISR and FSR

As discussed above, IRS photons are generated in CMS, while the FSR ones are generated in QMS. It is therefore the easiest to introduce the IR cut for the real photons in terms of minimum energy in these two frames. This defines the IR

boundary, and IR domains inside them, which are differential for ISR and FSR real photons. As long as the ISR-FSR interference (IFI) is omitted, this will not be an issue. However, the IFI is present and the IR boundary has to be common for the CEEEX. For the case of events with weight 1, this can be solved by taking the common IR domain which contains both ISR and FSR domains. For each event, we "remove from the record" all photons that are inside the new common IR domain. However, for the case of weighted events, the above approach has to be modified and it needs to be accompanied by the additional weight that is analytically calculable. We shall introduce this approach in the following.

Let us consider the case of EEX

$$\sigma_{\text{EEX}}^{(r)} = \int W_{\text{EEX}}^{(r)} d\sigma^{\text{Cru}}, \quad (7.242)$$

,where the model weight $W_{\text{EEX}}^{(r)}$ is defined in eq. (7.191) in terms of the $O(\alpha^r)$ EEX differential distributions (7.4). We arrive at

$$\begin{aligned} \sigma_{\text{EEX}}^{(r)} \{A\} &= \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \int d\sigma_{[n,n']}^{\text{Cru}}(\Omega_I, \Omega_F) \\ &\quad \times A(n, k_1, \dots, k_n; n', k'_1, \dots, k'_{n'}; p_i, q_i) \\ &\quad \times W_{\text{EEX}}^{(r)}(n, k_1, \dots, k_n; n', k'_1, \dots, k'_{n'}; p_i, q_i), \\ d\sigma_{[n,n']}^{\text{Cru}}(\Omega_I, \Omega_F) &\equiv ds_X \frac{\sigma_{\text{Born}}(s_X)}{4\pi} d\tau_{n+1}(P; k_1, \dots, k_n, X) \\ &\quad \times e^{\gamma_e \log \epsilon} \frac{1}{n!} \prod_{j=1}^n 2\tilde{S}_e(k_j) \bar{\Theta}(\Omega_I, k_j) d\tau_{n'+2}(X; k'_1, \dots, k'_{n'}, q_1, q_2) \\ &\quad \times \frac{s_X}{s_Q} \frac{2}{\beta_f} e^{\gamma_f \log \left(\delta_f \frac{s_Q + 2K'Q}{s_Q} \right)} \frac{1}{n'} \prod_{l=1}^{n'} 2\tilde{S}_f(k'_l) \bar{\Theta}(\Omega_F, k'_l + l), \end{aligned} \quad (7.243)$$

with the help of eq. (7.195) and

$$\epsilon_f = \delta_f \left(1 + \frac{2QK'}{s_Q} \right), \quad K' = \sum_{i=0}^{n'} k'_i. \quad (7.244)$$

Here we have introduced a general acceptance function A to discuss the IR cancellations. Each IR-safe observable corresponds uniquely to one or more such acceptance functions. The acceptance function corresponding to a physically meaningful, IR-safe, observable must follow the important rule

$$\lim_{k_i \rightarrow 0} A(n, k_1, \dots, k_{i-1}, k_i, k_{i+1}, \dots, k_n) = A(n-1, k_1, \dots, k_{i-1}, k_{i+1}, \dots, k_n), \quad (7.245)$$

and there should be a similar rule for FSR photons.

So far we have kept the IR domains different for ISR and FSR. For ISR, Ω_I was defined by: $k_j^0 < \epsilon_e \frac{1}{2} \sqrt{s}$ in the laboratory CMS system where $\vec{p}_1 + \vec{p}_2 = 0$. For FSR, Ω_F was defined by $k_j'^0 < \delta_f ((s_Q + 2K'Q)/s_Q)^{\frac{1}{2}} \sqrt{s_Q}$ in the QMS system where $\vec{q}_1 + \vec{q}_2 = 0$. Next, we are about to bring the two IR domains together

$$d\sigma_{[n,n']}^{\text{Cru}^*}(\Omega_I, \Omega_F). \quad (7.246)$$

It is known that the total cross section and any IR-safe observable should be independent of Ω_I and Ω_F . The intuitive solution is to set δ_f so small that $\Omega_F \subset \Omega_I$ always holds, and to neglect all FSR photons $k'_i \in \delta\Omega = \Omega_I \setminus \Omega_F$, i.e., removing them from the list of the generated momenta in the MC. Note that since $(s_Q + 2K'Q)/s_Q \sim s_X/s_Q \ll s_X/(4m_f^2)$. Next we will prove the validity of the approach described above. Let us consider the internal FSR subintegral in eq. (7.244),

fixing all ISR photon momenta

$$\begin{aligned} \mathfrak{J} &= \sum_{n'=0}^{\infty} \int d\tau_{n'+2}(X; k'_1, \dots, k'_n, q_1, q_2) \\ &\quad \times \frac{1}{n'!} \prod_{l=1}^{n'} 2\tilde{S}_f(k'_l) \Theta(\Omega_F^-, k'_l) b(k'_1, \dots, k'_{n'}; p_i, q_i), \end{aligned} \quad (7.247)$$

where

$$\begin{aligned} b(k'_1, \dots, k'_{n'}; p_i, q_i) &\equiv e^{\gamma_f \log\left(\delta_f \frac{s_Q + 2K'Q}{s_Q}\right)} \frac{s_X}{s_Q} \frac{2}{\beta_f} \\ &\quad \times W_{\text{EEX}}^{(r)}(n, k_1, \dots, k_n; n', k'_1, \dots, k'_{n'}; p_i, q_i) \\ &\quad \times A(n, k_1, \dots, k_n; n', k'_1, \dots, k'_{n'}; p_i, q_i). \end{aligned}$$

Given that $\Omega_I = \Omega_F \cup \delta\Omega$ we can split every photon integral into two parts and reorganize the sum factorizing out the integral over $\delta\Omega$

$$\begin{aligned} \mathfrak{J}\{A\} &= \sum_{n'=0}^{\infty} \frac{1}{n'!} \prod_{l=1}^{n'} \left\{ \int \frac{d^3 k'_l}{k'_l{}^0} \Theta(\delta\Omega, k'_l) \tilde{S}_f(l'_l) + \int \frac{d^3 k'_l}{k'_l{}^0} \Theta(\Omega_I, k'_l) \tilde{S}_f(l'_l) \right\} \\ &\quad \times \int d\tau_{n'+2}(X, k'_i; q_1, q_2) b(k'_1, \dots, k'_{n'}; p_i, q_i) \\ &= \sum_{n'=0}^{\infty} \frac{1}{n'!} \sum_{s=0}^{n'} \binom{n'}{s} \left\{ \int \frac{d^3 k}{2k^0} \Theta(\delta\Omega, k') \tilde{S}_f(k') \right\}^s \\ &\quad \times d\tau_{n'+2-s} \left(X - \sum_1^s; k'_1, \dots, k'_{n'-s}, q_1, q_2 \right) \\ &\quad \times \prod_{l=1}^{n'-s} \bar{\Theta}(\Omega_I, k'_l) \tilde{S}_f(k'_l) b(k'_1, \dots, k'_{n'-s}; p_i, q_i), \end{aligned} \quad (7.248)$$

where

$$\Theta(\delta\Omega, k') = \begin{cases} = 1 & \text{for } k' \in \delta\Omega, \\ = 0 & \text{otherwise.} \end{cases}$$

Note that because of the specific expansion (7.4) of $\rho_{\text{EEX}}^{(r)}$ into $\bar{\beta}$ -components the model weight $W_{\text{EEX}}^{(r)}$ is the most important ingredient in the above algebraic transformation.

The model weight $W_{\text{EEX}}^{(r)}$ satisfies the IR-safeness condition

$$\begin{aligned} & \lim_{k'_i \rightarrow 0} W_{\text{EEX}}^{(r)}(n', k'_1, \dots, k'_{i-1}, k'_i, k'_{i+1}, \dots, k'_{n'}) \\ &= W_{\text{EEX}}^{(r)}(n' - 1, k'_1, \dots, k'_{i-1}, k'_{i+1}, \dots, k'_{n'}), \end{aligned} \quad (7.249)$$

and so does the function $b(k'_1, \dots, k'_{n'}; p_i, q_i)$. Thus the integral becomes

$$\begin{aligned} \mathcal{J}\{A\} &= \sum_{n'=0}^{\infty} \int d\tau_{n'+2}(X; k'_1, \dots, k'_{n'}, q_1, q_2) \frac{1}{n!} \prod_{l=1}^{n'} 2\tilde{S}_f(k'_l) \Theta(\Omega_I^-, k'_l) \\ &\times \exp\left(\int \frac{d^3k}{2k^0} \Theta(\delta\Omega, k) 2\tilde{S}_f(k)\right) b(k'_1, \dots, k'_{n'}; p_i, q_i), \end{aligned} \quad (7.250)$$

getting an additional exponential factor.

Therefore, by the explicit calculation, it is valid to skip photons that fall into

$$\delta\Omega = \Omega_I \setminus \Omega_F$$

$$\begin{aligned} & d\sigma_{[n, n']}^{\text{Cru}*}(\Omega_I, \Omega_I) \\ &= d s_X \frac{\sigma_{\text{Born}}(s_X)}{4\pi} d\tau_{n+1}(P; k_1, \dots, k_n, X) \\ &\times e^{\gamma_e \log \epsilon} \frac{1}{n!} \prod_{j=1}^n 2\tilde{S}_e(k_j) \bar{\Theta}(\Omega_I, k_j) d\tau_{n'+2}(X; k'_1, \dots, k'_{n'}; q_1, q_2) \frac{s_X}{s_Q} \frac{2}{\beta_f} \\ &\times e^{R_F(\Omega_I)} \frac{1}{n!} \prod_{l=1}^{n'} 2\tilde{S}_f(k'_l) \Theta(\Omega_I^-, k'_l), \end{aligned} \quad (7.251)$$

where

$$R_F = \gamma_f \log\left(\delta_f \frac{s_Q + 2K'Q}{s_Q}\right) + 2Q_f^2 \alpha \tilde{B}(\Omega_I, q_1, q_2) - 2Q_f^2 \alpha \tilde{B}(\Omega_F, q_1, q_2). \quad (7.252)$$

Note that the integral is preserved by construction

$$\sum_{n, n'} \int d\sigma_{[n, n']}^{\text{Born}*}(\Omega_I, \Omega_F) = \sum_{n, n'} \int d\sigma_{[n, n']}^{\text{Born}}(\Omega_I, \Omega_F).$$

Now, we can not keep using $\rho_{\text{EEX}}^{(r)}$ of eq. (7.4) since the IR boundary in the new above distribution has changed for FSR photons. We have to use another distribution $\rho_{\text{EEX}}^{*(r)}$

in which $\widehat{B}(\Omega_F)$ is replaced by $\widehat{B}(\Omega_I)$ in the YFS form factor,

$$\rho_{\text{EEX}}^{*(r)} = \rho_{\text{EEX}}^{(r)} e^{2Q_f^2(\widetilde{B}(\Omega_I, q_1, q_2) - \widetilde{B}(\Omega_F, q_1, q_2))}.$$

And since the model weight is the ratio of the model distribution and the crude one, the new exponential factor cancel out. Thus the new model weight is functionally exactly the same

$$W_{\text{EEX}}^{*(r)} = W_{\text{EEX}}^{(r)}.$$

In the new MC calculation, we have

$$\sigma_{\text{EEX}}^{(r)} = \int W_{\text{EEX}}^{*(r)} d\sigma^{\text{Cru}*} = \int W_{\text{EEX}}^{(r)} d\sigma^{\text{Cru}*}. \quad (7.253)$$

This result is trivial since in the MC program for the EEX model we change almost nothing, only neglecting hidden photons in the evaluation of the model weight. This feature implies that very soft photons are not important for all IR-safe integrand functions.

The term $\gamma_f \log(\dots)$ in R_F is canceled by $\widetilde{B}(\Omega_F)$ and there is actually no dependence on Ω_F or δ_f in $d\sigma_{[n, n']}^{\text{Cru}*}$ any more. The IR cancellation is now ensured by the term below

$$2\frac{\alpha}{\pi} \left(\log \frac{2q_1 q_2}{m_f^2} - 1 \right) \log \epsilon.$$

However, the situation is still not as good as expected. We have to deal with the complication due to the use of the weighted events at the level of the crude distribution. Let us return to the EEX case

$$\sigma_{\text{EEX}}^{(r)} = \int W_{\text{EEX}}^{(r)} W_{\text{FSR}}^{\text{Cru}} W_{\text{ISR}}^{\text{Cru}} d\sigma^{\text{Pri}}. \quad (7.254)$$

Now the issue is that photons in $\delta\Omega$ cannot be hidden, because $W_{\text{FSR}}^{\text{Cru}}$ does not follow the IR-saftness condition

$$\begin{aligned} & \lim_{k'_i \rightarrow 0} W_{\text{FSR}}^{\text{Cru}}(n', k'_1, \dots, k'_i, \dots, k'_{n'}) \\ &= W_{\text{FSR}}^{\text{Cru}}(n' - 1, k'_1, \dots, k'_{i-1}, k'_{i+1}, \dots, k'_{n'}) \frac{f\left(\theta_i, \frac{m_f^2}{s_Q}\right)}{\bar{f}\left(\theta_i, \frac{m_f^2}{s}\right)}. \end{aligned}$$

Soft photons contribute the finite ratio (f/\bar{f}) , and this condition is essential for the IR-cancellations and for the overall normalization.

In order to save the validity of the approach of replacing Ω_F with Ω_I , we repeat the calculation of eq. (7.248) and assume that photons hidden inside $\delta\Omega$ do not contribute the factor (f/\bar{f}) to the overall weight. Then we obtain an expression with the modified exponential factor

$$\begin{aligned} \mathcal{I}'\{A\} &= \sum_{n'=0}^{\infty} \int d\tau_{n'+2}(X'; k'_1, \dots, k'_{n'}, q_1, q_2) \frac{1}{n'!} \prod_{l=1}^{n'} 2\tilde{S}_f(k'_l) \bar{\Theta}(\Omega_I, k'_l) \\ &\times \exp\left(\int_{\delta\Omega} \frac{d^3k}{k^0} \tilde{S}_f(k) \frac{\bar{f}\left(\theta_i, \frac{m_f^2}{s}\right)}{f\left(\theta_i, \frac{m_f^2}{s_Q}\right)}\right) b(k'_1, \dots, k'_{n'}; p_i, q_i). \end{aligned} \quad (7.255)$$

It is important that the effect from neglecting (f/\bar{f}) in the overall weight can be calculated analytically. If so, we can compensate analytically for the missing average contribution to $W_{\text{FSR}}^{\text{Cru}}$ from the hidden photons. The evaluation of the integral over $\delta\Omega$ is based on the relation

$$\tilde{S}_f^*(k) = \tilde{S}_f(k) \frac{\bar{f}\left(\theta_i, \frac{m_f^2}{s}\right)}{f\left(\theta_i, \frac{m_f^2}{s_Q}\right)} = -Q_f^2 \frac{\alpha}{4\pi} \left(\frac{q_1^*}{kq_1^*} - \frac{q_2^*}{kq_2^*} \right).$$

where q_i^* , $i = 1, 2$, are defined so that $(q_i^*)^2 = m_f^2(s_Q/s)$. Moreover, they have the same directions as the original \vec{q}_i and the same total energy, $q_1^{*0} + q_2^{*0} = \sqrt{s_Q}$ in the

QMS. Thus we have

$$I_{\delta\Omega} = \int_{\delta\Omega} \frac{d^3k}{k^0} \tilde{S}_f(k) \frac{\bar{f}\left(\theta_i, \frac{m_f^2}{s}\right)}{f\left(\theta_i, \frac{m_f^2}{s_Q}\right)} = 2\alpha Q_f [\tilde{B}(\Omega_I, q_1^*, q_2^*) - \tilde{B}(\Omega_F, q_1^*, q_2^*)]. \quad (7.256)$$

To sum up, for the case of the weighted events, the method of hiding photons in $\delta\Omega = \Omega_I \setminus \Omega_F$ leads to a new crude distribtuion similar to that in eq. (7.251) with the new

$$R_f = \gamma_f \log\left(\delta_f \frac{s_Q + 2K'Q}{s_Q}\right) + 2Q_f^2 \alpha \tilde{B}(\Omega_I, q_1^*, q_2^*) - 2Q_f^2 \alpha \tilde{B}(\Omega_F, q_1^*, q_2^*). \quad (7.257)$$

Consequently the above exponential factor does not cancel not exactly in the model weight with the corrections to the YFS form factor as before. And we have the correcting factor in the model weight:

$$W_{\text{hide}} = \exp\{-2\alpha Q_f [\tilde{B}(\Omega_I, q_1^*, q_2^*) - \tilde{B}(\Omega_F, q_1^*, q_2^*)] + 2\alpha Q_f [\tilde{B}(\Omega_I, q_1, q_2) - \tilde{B}(\Omega_F, q_1, q_2)]\}. \quad (7.258)$$

The important asset from the approach of hiding photons in $\delta\Omega = \Omega_I \setminus \Omega_F$ is that with the above correcting factor we could do calculation for the CEEEX model with the ISR-FSR interference switched on.

7.4.5 Entire MC Algorithm Top-to-Bottom

For the CEEEX model, according to the results of the previous subsections, we obtain

$$\sigma_{\text{CEEEX}}^{(r)}\{A\} = \sum_{f=\mu,\tau,d,u,s,c,b} \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \int AW_{\text{CEEEX}}^{(r)} W_{\text{FSR}}^{\text{Cru}} W_{\text{ISR}}^{\text{Cru}} W_{\text{hide}} d\sigma_{[n,n']}^{\text{Pri}*}(\Omega_I). \quad (7.259)$$

And $d\sigma_{[n,n']}^{\text{Pri}}(\Omega_I)$ is derived from the product of the ISR and FSR primary differential distributions

$$d\sigma_{[n,n']}^{\text{Pri}}(\Omega_I, \Omega_F) = d\mathcal{J}_n^{\text{Pri}}(\Omega_I) d\mathcal{J}_{n'}^{\text{Pri}}(\Omega_F) \quad (7.260)$$

by means of hiding FSR photons in $\delta\Omega$. Therefore, only momenta outside the common IR-domain enter into the evaluation of $W_{\text{CEEX}}^{(r)}$ and of all other weights.

The integrated cross section with the acceptance function A is obtained in the MC run in a standard way

$$\sigma_{\text{CEEX}}^{(r)}\{A\} = \langle AW_{\text{CEEX}}^{(r)} W_{\text{FSR}}^{\text{Cru}} W_{\text{ISR}}^{\text{Cru}} W_{\text{hide}} \rangle \sigma^{\text{Pri*}}. \quad (7.261)$$

The overall normalization is based on

$$\begin{aligned} \sigma^{\text{Pri*}} &= \sum_{f=\mu,\dots,b} \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \int d\sigma_{[n,n']}^{\text{Pri*}}(\Omega_I) = \sum_{n=0}^{\infty} \int d\mathfrak{J}_n^{\text{Pri}}(\Omega_I) \sum_{n'=0}^{\infty} \int d\mathfrak{F}_{n'}^{\text{Pri}}(\Omega_F) \\ &= \sum_{f=\mu,\dots,b} \sum_{n=0}^{\infty} \int d\mathfrak{J}_n^{\text{Pri}}(\Omega_I) \\ &= \sum_{f=\mu,\dots,b} \int_0^1 dv \sigma_{\text{Born}}^f(s(1-v)) \mathcal{J}_0(v) \bar{\gamma}_e v^{\bar{\gamma}_e-1} \epsilon^{\gamma_e-\bar{\gamma}_e}, \end{aligned} \quad (7.262)$$

where we have used the property $\int \sum d\mathfrak{F}^{\text{Pri}}(\Omega_F) \equiv 1$ of eq. (7.221), and the ISR part is taken from eq. (7.239).

Now we have the entire MC algorithm from the top to the bottom. It starts from the generation of v describing the total energy loss due to the IRS, the type of final fermion f and the photon multiplicities n and n' , and then generate photon energies and angles using the method described in the previous subsections.

CHAPTER EIGHT

Interface between KKMC-hh and MG5_aMC@NLO

With the help of KKMC-hh [90–92], the coherent exclusive exponentiation (CEEX) electroweak (EW) exact $O(\alpha^2 L)$ correction for the Drell-Yan process (please read Appendix E) has been achieved. In order to realize the EW+the next-to-leading order (NLO) QCD correction for the Drell-Yan process, we will first apply the Madgraph5_aMC@NLO (MG5_aMC@NLO) [118] to obtain the next-to-leading order QCD correction, and then we will interface KKMC-hh with MG5_aMC@NLO via merging their LHE files to achieve the EW and NLO QCD corrections. In this chapter, we will first introduce the overview of MG5_aMC@NLO. Next we will describe the approach to interface KKMC-hh with MG5_aMC@NLO. Finally we will exhibit and discuss our results.

8.1 Overview of Madgraph_aMC@NLO

MADGRAPH [119] is a powerful tool for automatically generating matrix elements for high energy physics process, such as $2 \rightarrow n$ scatterings and decays. First the user inputs a specific process in terms of initial and final particles, allowing some refined criteria. As a result, MADGRAPH generates all Feynman diagrams for the process, and yields the computer code to compute the matrix element at a given phase space point. The matrix element calculation is done using the helicity amplitudes technique which was first implemented in the package HELAS [120]. The application of the helicity amplitudes is efficient because it allows the helicity amplitudes corresponding to identical subdiagrams to be reused between the diagrams, leading

a considerable optimization. The computer code generate by MADGRAPH can then be used for the cross section and decay width evaluation and event generation.

The essential idea of MADGRAPH5_aMC@NLO is the same as the MADGRAPH family. The structure of a cross section is essentially independent of the process regardless of the theory and of the perturbative order, and thus it can be written as a computer code once and for all. For example, phase phases can be defined in full generality, leaving only particle masses and number as free parameters. Conversely, matrix elements which are obviously dependent on the theory and process can be calculated starting from a limited number of formal instructions, such as Feynman rules and recursion relations. MADGRAPH5_aMC@NLO is written in a meta-code, in which a Python code writes a Python, C++ or Fortan code. The latter code is specific to the desired process. MADGRAPH5_aMC@NLO includes two ingredients. The first one is a theory model, which is equivalent to the Lagrangian of the theory and its parameters, such as masses and coupling constant. The second one is a set of process-indepdent building blocks for automation of calculations. The automation of NLO computation involves the FKS subtraction block, which carrys out the generation of the real corrections with the proper subtractions automatically [121–126], by interfacing MadFKS [126]. Besides the module for real corrections, the automation of NLO computation requires a specific module for virtual corrections. In MG5_aMC@NLO, the virtual contribution to an NLO cross section is achieved through the module MADLOOP [127], which is based on the OPP integrand reduction technique [128]. These two module above together allow a fully automatic computation of infrared-safe observables at NLO in QCD. After the integration of

the matrix element hard process, the full parton shower and hadronization infrastructures, such as Herwig and Pythia, etc., are also critical for an accurate simulation of a hadronic process. In this thesis, we have applied MG5_aMC@NLO + Herwig to obtain the simulation of the Drell-Yan process $pp \rightarrow Z/\gamma \rightarrow \mu^+\mu^- + X$ at the next-to-leading order in QCD.

8.2 Interfacing KKMC-hh with MG5_aMC@NLO

We will now introduce our approach to interface KKMC-hh and MG5_aMC@NLO. Our essential idea is merging their LHE [129] files to achieve the EW and NLO QCD corrections. Namely, given that LHE file from the MG5_aMC@NLO contains all the information of the events at the partonic level, we extract the next-to-leading order contribution in QCD for Drell-Yan process from the LHE file and combine it with the weight of KKMC-hh to achieve the EW and NLO QCD corrections of the Drell-Yan process. We are exhibiting our method in details in the following.

In the KKMC-hh, the basic event $(x_1, x_2, v, W_{\text{Basic}})$ is generated by the distribution

$$\begin{aligned} \rho = & 2N_q \left[\frac{Q_s^2}{z^2 s} \left(\frac{1}{Q_{smin}} - \frac{1}{Q_{samx}} \right) \right] \log\left(\frac{s}{Q_s}\right) f_{q_1}(\sqrt{Q_s}, x_1) f_{\bar{q}_2}(\sqrt{Q_s}, x_2) \\ & \times \frac{1}{2} \left(\frac{\bar{\gamma}}{\gamma} \right) \left(\frac{v}{v_{min}} \right)^{\bar{\gamma}-\gamma} \left(1 + \frac{1}{\sqrt{1-v}} \right) v_{max}^{\gamma} \\ & \times \frac{\sigma_{Born}((1-v)Q_s)}{3(1-v)}. \end{aligned} \quad (8.1)$$

which includes three components:

$$\begin{aligned} \rho(x_1, x_2) = & 2N_q \left[\frac{Q_s^2}{z^2 s} \left(\frac{1}{Q_{smin}} - \frac{1}{Q_{samx}} \right) \right] \log\left(\frac{s}{Q_s}\right) f_{q_1}(\sqrt{Q_s}, x_1) f_{\bar{q}_2}(\sqrt{Q_s}, x_2), \\ \rho(v) = & \frac{1}{2} \left(\frac{\bar{\gamma}}{\gamma} \right) \left(\frac{v}{v_{min}} \right)^{\bar{\gamma}-\gamma} \left(1 + \frac{1}{\sqrt{1-v}} \right) v_{max}^{\gamma}, \text{ and } \frac{\sigma_{Born}((1-v)Q_s)}{3(1-v)}, \end{aligned}$$

where x_1 and x_2 describe beamstrahlung, v describes the total energy loss due to ISR, and W_{Basic} is the weight corresponding to the basic event.

The weight **XWGTUP** derived from the LHE file of MG5_aMC@NLO represents the cross section of the process in units of pb, which includes the partonic cross section, contributions from parton distribution functions (PDF) and the perturbative QCD corrections. To be more specific, for the Drell-Yan process $pp \rightarrow Z/\gamma^* \rightarrow l^+l^- + X$, the differential cross section is

$$\frac{d\sigma}{dydM} = \sum_{i,j} \hat{\sigma}_{\text{Born}} \int_{x_1}^1 dx_i \int_{x_2}^1 dx_j f_i(x_i, Q^2) f_j(x_j, Q^2) \Delta(x_1, x_2, x_i, x_j, Q^2) \quad (8.2)$$

where $\hat{\sigma}_{\text{Born}}$ is the partonic cross section of the Drell-Yan process, Δ_{ij} is the perturbative QCD coefficient function for the Drell-Yan process. And the partonic cross section of the Drell-Yan process can be written as

$$\hat{\sigma}_{\text{Born}} = \frac{1}{3} \cdot \frac{4\pi\alpha^2}{3Q^2} \sum q_f^2. \quad (8.3)$$

In order to interface MG5_aMC@NLO with KKMC-hh, we need to replace the basic weight W_{Basic} with the weight from **XWGTUP** after removing the intersections between W_{Basic} and **XWGTUP**, and use momentums of initial quark pairs derived from LHE file of MG5_aMC@NLO to generate a new pair of parton momentum fractions x_1 and x_2 .

We see **XWGTUP** has three components: partonic cross section $\hat{\sigma}_{\text{Born}}$, the function of x_i and x_j and QCD corrections. So the intersections between W_{Basic} and **XWGTUP** are $\hat{\sigma}_{\text{Born}}$ and the function of x_1 and x_2 . Therefore, we need to remove $\rho(x_1, x_2)$ from ρ . Since KKMC-hh would calculate the crude Born cross section after ISR generation, we remove $\hat{\sigma}_{\text{Born}}$ from **XWGTUP**. Then, we could have the ρ'

$$\rho' = \frac{\text{XWGTUP}}{\hat{\sigma}_{\text{Born}}} \times \frac{1}{2} \left(\frac{\bar{\gamma}}{\gamma} \right) \left(\frac{v}{v_{\text{min}}} \right)^{\bar{\gamma}-\gamma} \left(1 + \frac{1}{\sqrt{1-v}} \right) v_{\text{max}}^\gamma \frac{\sigma_{\text{Born}}((1-v)Q_s)}{3(1-v)} \quad (8.4)$$

In $\rho(x_1, x_2)$, the factor $2N_q$ comes from summation over quarks, where N_q is determined by the subroutine **hh_Quarks**. Since we need to replace $\rho(x_1, x_2)$ with the **LHE** weight removing the Born cross section calculated by MG5_aMC@NLO, **XWGTUP**/ $\hat{\sigma}_{Born}$, we would replace this factor together with $\rho(x_1, x_2)$ in the program.

In order to realize our approach in the computer programming, we first coded a new subroutine **UPYVNT**, which read event information from the LHE file of MG5_aMC@NLO. Then we coded another new subroutine **hh_MakeLHE** (called before the subroutine **hhFoam_Make**) to use momentums \hat{p}_1 and \hat{p}_2 of initial quark pairs from LHE file to compute the pair of x_1 and x_2 as follows:

$$\begin{aligned} Q_s &= Q^2 = (\hat{p}_1 + \hat{p}_2)^2, \\ z &= \frac{Q_s}{s}, \end{aligned} \quad (8.5)$$

where $s = E_{\text{CMS}}^2$. Then we have

$$x_1 = z^{r_2}, x_2 = z^{1-r_2},$$

where r_2 is a random number. And the corresponding weight is

$$W_{\text{LHE}} = \frac{\mathbf{XWGTUP}}{\hat{\sigma}_{\text{Born}}}.$$

Ater generating x_1 and x_2 , we modified the subroutine **hhBornV_RhoFoam** so that it would only calculate $\bar{\gamma}$ and γ with the help of new Q_s and generate a new v . In sum, the new W_{Basic} is calculated according to eq. (8.4).

With the help of the new basic weight of eq. (8.4), the cross section for interfacing KKMC-hh with MG5_aMC@NLO can be evaluated by

$$\sigma = \rho' \langle \prod_k w_k \rangle. \quad (8.6)$$

In the program, this cross section is calculated by the average of main weight W_{Main} .

The main weight has two components W_{Crud} and W_{Best} :

$$W_{\text{Main}} = W_{\text{Crud}} \times W_{\text{Best}}.$$

The crude weight W_{Crud} is calculated in subroutine **KK2f_Make**. The crude weight has two components, the ISR components W_{ISR} and the FSR components W_{FSR} .

$$W_{\text{Crud}} = W_{\text{ISR}} \times W_{\text{FSR}},$$

where the ISR components W_{ISR} is calculated by subroutine **KarLud_Make** and the FSR components W_{FSR} is calculated by subroutine **KarFin_Make**

$$W_{\text{ISR}} = W_{\text{Basic}} \times W_{\text{Mass}} \times W_{\text{Dil}} \times W_{\text{Cut}} \times W_{\text{KF}},$$

$$W_{\text{FSR}} = W_1 \times W_2 \times W_3, \quad (8.7)$$

The brief explanations for components are as follows:

(i) W_{Basic} : the basic weight W_{Basic} is calculated by eq. (8.4), which generates v , x_1 and x_2 . It includes not only the electroweak contribution but the NLO QCD corrections as well.

(ii) W_{Mass} : the weight W_{Mass} corresponds the simplification made on the photon angular distribution by dropping mass terms. If there is no photon above detectability threshold, then

$$W_{\text{Mass}} = 1.$$

If not,

$$W_{\text{Mass}} = \prod_{i=1}^n \frac{f(\theta_i)}{\bar{f}(\theta_i)}. \quad (8.8)$$

where

$$f(\theta_i) = \frac{\alpha}{\pi^2} \left[\frac{1}{(1 - \beta \cos \theta_i)(1 + \beta \cos \theta_i)} - \frac{m_e^2}{s} \frac{1}{(1 - \beta \cos \theta_i)^2} - \frac{m_e^2}{s} \frac{1}{(1 + \beta \cos \theta_i)^2} \right],$$

$$\bar{f}(\theta_i) = \frac{\alpha}{\pi^2} \frac{1}{(1 - \beta \cos \theta_i)(1 + \beta \cos \theta_i)}. \quad (8.9)$$

(iii) W_{Dil} : this weight corresponds to the simplification made on the dilatation Jacobian.

$$J(\bar{k}, v) = \frac{1}{2} \left(1 + \frac{1}{\sqrt{1 - Av}} \right) \longrightarrow J_0(v) = \frac{1}{2} \left(1 + \frac{1}{\sqrt{1 - v}} \right). \quad (8.10)$$

We therefore have

$$W_{\text{Dil}} = \frac{J(\bar{k}, v)}{J_0(v)} = \frac{1 + \sqrt{1 - Av}}{1 + \sqrt{1 - v}}. \quad (8.11)$$

(iv) W_{Cut} : this weight corresponds to the lower photon energy boundary:

$$W_{\text{Cut}} = \theta(\lambda_0(\bar{k}, v)x_n - \epsilon) = \theta\left(\frac{2k_0^n}{\sqrt{s}} - \epsilon\right). \quad (8.12)$$

(v) W_{KF} : this weight corresponds the generation of **KF** codes, generated by subroutine **MBrA_GenKF**.

(vi) W_1 : it is the weight corresponding to phase space limits for very hard photon, generated by subroutine **KarFin_YFSfin**,

$$W_1 = 1. \quad (8.13)$$

(vii) W_2 : it corresponds to the weight for translation Jacobian, generated by subroutine **KarFin_YFSfin**,

$$W_2 = \frac{1 + \bar{K}^0}{1 + \bar{K}^0 + \frac{1}{4}\bar{K}^2}. \quad (8.14)$$

(viii) W_3 : it is the weight that corresponds to the following simplifications:

$$f\left(\theta_j, \frac{m_f^2}{s_Q}\right) \longrightarrow \bar{f}\left(\theta_j, \frac{m_f^2}{s_X}\right), \quad e^{\gamma_f \log(\delta_f((1 + \bar{K}^0)))} \longrightarrow e^{\gamma_f \log(\delta_f)} \quad (8.15)$$

where

$$\begin{aligned}
f\left(\theta_j, \frac{m_f^2}{s_Q}\right) &= \frac{1 + \beta_f^2}{\delta_{1j}\delta_{2j}} - \frac{\mu_f^2}{2} \frac{1}{\delta_{1j}^2} - \frac{\mu_f^2}{2} \frac{1}{\delta_{2j}^2}, \\
\delta_{1j} &= 1 - \beta_f \cos\theta_j, \\
\delta_{2j} &= 1 + \beta_f \cos\theta_j, \\
\bar{f}\left(\theta_j, \frac{m_f^2}{s_X}\right) &= \frac{1 + \bar{\beta}_f^2}{\bar{\beta}_f^2} \frac{1}{1 - \bar{\beta}_f^2 \cos^2\theta_j}, \\
\bar{\beta}_f &= \sqrt{1 - \left(\frac{m_f^2}{s_X}\right)^2}, \\
\bar{\gamma}_f &= Q_f^2 \frac{\alpha}{\pi} \frac{1 + \bar{\beta}_f^2}{\bar{\beta}_f} \log \frac{1 + \bar{\beta}_f}{1 - \bar{\beta}_f}. \tag{8.16}
\end{aligned}$$

Therefore, we have

$$W_3 = e^{\gamma_f \log(\delta_f((1+\bar{K}^0))) - \gamma_f \log(\delta_f)} \prod_{j=1}^{n'} \frac{f\left(\theta_j, \frac{m_f^2}{s_Q}\right)}{\bar{f}\left(\theta_j, \frac{m_f^2}{s_X}\right)} \tag{8.17}$$

And the model weight W_{Best} is of the $O(\alpha^{(2)})$, calculated by the sub-routine **GPS_Make**,

$$W_{\text{CEEX}}^{(2)}(p_a, p_b, p_c, p_d; k_1, \dots, k_n) = \frac{\rho_{\text{CEEX}}^{(2)}(p_a, p_b, p_c, p_d; k_1, \dots, k_n)}{\rho_{[i,n]'}^{\text{Cru}}(p_c, p_d; k_1, \dots, k_n) (2\pi)^{3(n+2)-4}}. \tag{8.18}$$

Please read Section (7.2) and Subsection (7.4.1) for details.

Therefore, with the help of the W_{Main} , the cross section which includes electroweak and NLO QCD corrections will be evaluated. Our results will be exhibited in the next section.

8.3 Results for Interfacing KKMC-hh with MG5_aMC@NLO

We will now discuss the results of interfacing KKMC-hh with MG5_aMC@NLO. Specifically, we compared the results of the Drell-Yan process $pp \rightarrow Z/\gamma^* \rightarrow \mu^+\mu^- + X$ obtained by KKMC-hh, MG5_aMC@NLO and MG5_aMC@NLO interfaced with

KKMC-hh respectively. We made these comparisons at $\sqrt{s} = 13$ TeV with the ATLAS cuts on the Z/γ^* production and decay to lepton pairs [130]:

$$80 \text{ GeV} < M_{\ell\ell} < 100 \text{ GeV}, \quad P_T^{\ell\ell} < 30 \text{ GeV}$$

where both members of the decay lepton pair satisfy

$$P_T^\ell > 25 \text{ GeV}, \quad |\eta_\ell| < 2.4.$$

We here defined $M_{\ell\ell}$ as the lepton pair invariant mass, $P_T^{\ell\ell}$ as the transverse momentum of the lepton pair, P_T^ℓ as the transverse momentum of the lepton or antilepton ℓ , and η_ℓ as the pseudorapidity of the lepton or antilepton ℓ . We take the quark masses as $m_u = 6.0$ MeV, $m_d = 10.0$ MeV, $m_s = 0.15$ GeV, $m_c = 1.67$ GeV and $m_b = 4.78$ GeV [131].

The results calculated by three methods based on 1 million events are listed as follows:

Table 8.1: Cross Sections obtained by MG5_aMC@NLO \otimes KKMC-hh, KKMC-hh, MG5_aMC@NLO, respectively

Generator	Cross Section (pb)
MG5_aMC@NLO \otimes KKMC-hh	2144.72 ± 7.46
KKMC-hh	1707.68 ± 2.44
MG5_aMC@NLO	1816.00 ± 2.20

The first quantity that we compared were the transverse momentum distributions of muon. As we see, in Figure 8.1, the result obtained by KKMC-hh is larger than that obtained by MG5_aMC@NLO for $P_T < 35$ GeV and $P_T > 52$ GeV but smaller for $35 \text{ GeV} < P_T < 46 \text{ GeV}$. In the range $46 \text{ GeV} < P_T < 52 \text{ GeV}$,




Figure 8.1: Muon transverse momentum distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.

the results derived from KKMC-hh and MG5_aMC@NLO overlap. However, the result derived from MG5_aMC@NLO interfaced with KKMC-hh did not exhibit the enhancement for the muon transverse momentum distributions.

The next quantity we compared is the muon pseudorapidity distribution. The pseudorapidity η is a spatial coordinate describing the angle of a particle relative to the beam axis, commonly used in the experimental particle physics. It is defined as

$$\eta = \log \left(\tan \frac{\theta}{2} \right), \quad (8.19)$$

where the angle θ is angle between the particle three-momentum \vec{p} and the positive direction of the beam axis. The pseudorapidity can also be expressed in terms of three-momentum:

$$\eta \equiv \frac{1}{2} \log \left(\frac{|p| + p_L}{|p| - p_L} \right), \quad (8.20)$$




Figure 8.2: Muon pseudorapidity distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.

where p_L is the component of the momentum along the beam axis, namely, the longitudinal momentum. From Figure 8.2, we find that interfacing MG5_aMC@NLO with KKMC-hh results an apparent enhancement on the muon pseudorapidity distribution compared with that derived from MG5_aMC@NLO.

We compared not only quantities of the single lepton but those of lepton pairs as well. The dimuon transverse momentum distributions obtained by these three approaches are given in the Figure 8.3. As we can see, the differential cross section calculated by MG5_aMC@NLO interfaced with KKMC-hh is larger than that obtained by MG5_aMC@NLO only, and the enhancement is from the EW corrections calculated by KKMC-hh.

And the Figure 8.4 described the dimuon invariant mass distributions. By comparing the dimuon invariant mass distribution derived from MG5_aMC@NLO interfaced with KKMC-hh with that from MG5_aMC@NLO only, we find there is also




Figure 8.3: Dimuon transverse momentum distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.

an enhancement that is due to the EW corrections provided by KKMC-hh. Besides, we see the resonance peaks near 91 GeV derived from these three generators.

Finally, let us see the dimuon rapidity distributions in the Figure 8.5. The rapidity is defined as

$$y \equiv \frac{1}{2} \log \left(\frac{E + p_L}{E - p_L} \right). \quad (8.21)$$

We see that MG5_aMC@NLO interfaced with KKMC-hh amplified the dimuon rapidity distribution obtained by MG5_aMC@NLO only. The amplification can be viewed as a consequence of the EW corrections.

In sum, we exhibited the comparisons of the results obtained by MG5_aMC@NLO⊗KKMC-hh, MG5_aMC@NLO and KKMC-hh. We find that MG5_aMC@NLO interfaced with KKMC-hh would enhance the results obtained by MG5_aMC@NLO only and the enhancement is due to the EW corrections derived from KKMC-hh.




Figure 8.4: Dimuon invariant mass distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.




Figure 8.5: Dimuon rapidity distributions for KKMC-hh (blue), MG5_aMC@NLO (violet) and MG5_aMC@NLO interfaced with KKMC-hh (red) with the cuts specified in the text.

CHAPTER NINE

Overall Summary

In this dissertation, we have developed a new numerical method to calculate the general five-point function, which is important for evaluating one-loop radiative corrections. Our method is developed from the magic spinor product approach in loop integrals proposed by B. F. L. Ward originally, which applied the "Chinese magic" spinor technique to simplify the loop integral so that the E_0 could be expressed in terms of n -point one-loop integrals ($n \leq 4$). And the n -point one-loop integrals ($n \leq 4$) can be calculated numerically by the package LoopTools. Theoretically, the magic spinor product method should provide more efficiency and numerical stability for the evaluation of the general five point function. By comparing the results obtained by our method with those directly obtained from LoopTools, we find that they agreed with each other overall. Such agreements are encouraging.

Additionally, we also developed an approach to achieve the next-to-leading order and the electroweak (EW) exact $O(\alpha_s \otimes \alpha^2 L)$ corrections, interfacing MG5_aMC@NLO with KKMC-hh by merging their LHE files. We first coded a program to read the event information from the LHE file of MG5_aMC@NLO, and then extracted the next-to-leading QCD $O(\alpha_s)$ correction. Combining the NLO QCD corrections computed by MG5_aMC@NLO with the basic weight for generating events in the KKMC-hh, we obtained a new basic weight including both the NLO QCD $O(\alpha_s)$ corrections and the EW $O(\alpha^2 L)$ corrections. With the help of new basic weight, the new events

with $O(\alpha_s \otimes \alpha^2 L)$ were generated. We compared the muon transverse momentum distributions, muon pseudorapidity distributions, dimuon invariant mass distributions, dimuon rapidity distributions obtained by KKMC-hh, MG5_aMC@NLO and KKMC-hh interfaced with MG5_aMC@NLO , at $\sqrt{s} = 13$ TeV with the ATLAS cuts on the Z/γ^* production and decay to lepton pairs, respectively. By comparing the results of the Drell-Yan process obtained by these three generators, we find that the results derived from KKMC-hh interfaced with MG5_aMC@NLO bring enhancements from those derived from MG5_aMC@NLO, which are due to the EW corrections provided by KKMC-hh. We conclude that interfacing MG5_aMC@NLO with KKMC-hh would provide a way to achieve the exact $O(\alpha_s \otimes \alpha^2 L)$ corrections.

APPENDICES

APPENDIX A

Feynman Rules of the Electroweak SM

In this appendix we outline the Feynman rules of electroweak SM in the 't Hooft-Feynman gauge including the counterterms. In the vertices all momenta are set up as incoming.

Propagators:

for gauge bosons $V = \gamma, Z, W$ in the 't Hooft-Feynman gauge ($\xi_i = 1$)

$$V_\mu \text{---} \overset{k}{\text{~~~~~}} \text{---} V_\nu = \frac{-ig_{\mu\nu}}{k^2 - M_V^2},$$

for Faddeev-Popov ghosts $G = u^\gamma, u^Z, u^W$

$$G \text{---} \overset{k}{\text{-----}} \text{---} \bar{G} = \frac{i}{k^2 - M_G^2},$$

for scalar fields $S = H, \chi, \phi$

$$S \text{---} \overset{k}{\text{-----}} \text{---} S = \frac{i}{k^2 - M_S^2},$$

and for fermion fields $F = f_i$

$$F \text{---} \overset{p}{\text{-----}} \text{---} \bar{F} = \frac{i}{\not{p} - m_f}.$$

In the 't Hooft-Feynman gauge we have the following relations:

$$M_{u^\gamma} = 0, \quad M_{u^Z} = M_\chi = M_Z, \quad M_{u^\pm} = M_\phi = M_W. \quad (\text{A.1})$$

Tadpole:

$$\textcircled{\otimes} \text{---} \overset{S}{\text{---}} = i\delta t.$$

VV -counterterm:

$$V_\mu \text{---} \overset{k}{\textcircled{\otimes}} \text{---} V_\nu - ig_{\mu\nu}[C_1 k^2 - C_2]$$

with the actual values of V_1 , V_2 and C_1 , C_2

$$\begin{aligned} W^+W^- : \quad C_1 &= \delta Z_W, \quad C_2 = M_W^2 \delta Z_W + \delta M_W^2 \\ ZZ : \quad C_1 &= \delta Z_{ZZ}, \quad C_2 = M_Z^2 \delta Z_{ZZ} + \delta M_Z^2 \\ AZ : \quad C_1 &= \frac{1}{2} \delta Z_{AZ} + \frac{1}{2} \delta Z_{ZA}, \quad C_2 = M_Z^2 \frac{1}{2} \delta Z_{ZA} \\ AA : \quad C_1 &= \delta Z_{AA}, \quad C_2 = 0. \end{aligned} \tag{A.2}$$

SS -counterterm:

$$S_1 \leftarrow \text{---} \overset{k}{\textcircled{\otimes}} \text{---} \rightarrow S_2 = i[C_1 k^2 - C_2],$$

with the actual values of S_1 , S_2 and C_1 , C_2

$$HH : C_1 = \delta Z_H, \quad C_2 = M_H^2 \delta Z_H + \delta M_H^2. \tag{A.3}$$


FF -counterterm:

$$\begin{aligned}
& F_1 \xrightarrow{p} \otimes \xrightarrow{\quad} \bar{F}_2 \\
& = i[C_L \not{p} \frac{1}{2}(1 - \gamma_5) + C_R \not{p} \frac{1}{2}(1 + \gamma_5)] - C_S^- \not{p} \frac{1}{2}(1 - \gamma_5 - C_S^+ \not{p} \frac{1}{2}(1 + \gamma_5)],
\end{aligned}$$

with the actual values of F_1 , \bar{F}_2 and C_L, C_R, C_S^-, C_S^+

$$f_i \bar{f}_i : \begin{cases} C_L = \frac{1}{2}(\delta Z_{ij}^{f,L} + \delta Z_{ij}^{f,L,\dagger}), \\ C_R = \frac{1}{2}(\delta Z_{ij}^{f,R} + \delta Z_{ij}^{f,R,\dagger}) \\ C_S^- = m_{f,i} \frac{1}{2} \delta Z_{ij}^{f,L} + m_{f,i} \frac{1}{2} \delta Z_{ij}^{f,R,\dagger} + \delta_{ij} \delta m_{f,i}, \\ C_S^+ = m_{f,i} \frac{1}{2} \delta Z_{ij}^{f,R} + m_{f,i} \frac{1}{2} \delta Z_{ij}^{f,L,\dagger} + \delta_{ij} \delta m_{f,i}. \end{cases} \quad (\text{A.4})$$

VVVV-coupling:




$$= ie^2 C [2g_{\mu\nu}g_{\sigma\rho} - g_{\nu\rho}g_{\mu\sigma} - g_{\rho\mu}g_{\nu\sigma}]$$

with the actual values of V_1, V_2, V_3, V_4 and C

$$\begin{aligned}
W^+W^+W^-W^- : \quad C &= \frac{1}{s_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + 2\delta Z_W \right] \\
W^+W^-ZZ : \quad C &= -\frac{c_W^2}{s_W^2} \left[1 + 2\delta Z_e - 2\frac{1}{c^2} \frac{\delta s_W}{s_W} + 2\delta Z_W + \frac{1}{2}\delta Z_{ZZ} \right] + \frac{1}{2} \frac{c_W}{s_W} \\
W^+W^-AZ : \quad C &= \frac{c_W}{s_W} \left[1 + 2\delta Z_e - \frac{1}{c^2} \frac{\delta s_W}{s_W} + \delta Z_W + \frac{1}{2}\delta Z_{ZZ} + \frac{1}{2}\delta Z_{AA} \right] \\
&\quad - \frac{1}{2}\delta Z_{AZ} - \frac{1}{2} \frac{c_W^2}{s_W^2} \delta Z_{ZA} \\
W^+W^-AA : \quad C &= -[1 + 2\delta Z_e + \delta Z_W + \delta Z_{AA}] + \frac{1}{2} \frac{c_W}{s_W} \delta Z_{ZA}. \quad (\text{A.5})
\end{aligned}$$

VVV-coupling:




$$= -ieC[g_{\mu\nu}(k_2 - k_1)_\rho - g_{\nu\rho}(k_3 - k_2)_\mu - g_{\rho\mu}(k_1 - k_3)_\nu]$$

with the actual values of V_1, V_2, V_3 and C

$$\begin{aligned}
 AW^+W^- : \quad C &= 1 + \delta Z_e + \delta Z_W + \frac{1}{2}\delta Z_{AA} - \frac{1}{2}\frac{c_W}{s_W}\delta Z_{ZA}, \\
 ZW^+W^- : \quad C &= -\frac{c_W}{s_W} \left[1 + \delta Z_e - \frac{1}{c^2}\frac{\delta s_W}{s_W} + \delta Z_W + \frac{1}{2}\delta Z_{ZZ} \right] + \frac{1}{2}\delta Z_{AZ}.
 \end{aligned}
 \tag{A.6}$$


SSSS-coupling:



with the actual values of S_1, S_2, S_3, S_4 and C

$$\begin{aligned}
HHHH : \quad C &= -\frac{3}{4s_W^2} \frac{M_H^2}{M_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{\delta M_H^2}{M_H^2} + 2\delta Z_H \right], \\
HH\chi\chi : \quad C &= -\frac{1}{4s_W^2} \frac{M_H^2}{M_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{\delta M_H^2}{M_H^2} + \delta Z_H \right], \\
HH\phi\phi : \quad C &= -\frac{1}{4s_W^2} \frac{M_H^2}{M_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{\delta M_H^2}{M_H^2} + \delta Z_H \right], \\
\chi\chi\chi\chi : \quad C &= -\frac{3}{4s_W^2} \frac{M_H^2}{M_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{\delta M_H^2}{M_H^2} \right], \\
\chi\chi\phi\phi : \quad C &= -\frac{1}{4s_W^2} \frac{M_H^2}{M_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{\delta M_H^2}{M_H^2} \right], \\
\phi\phi\phi\phi : \quad C &= -\frac{1}{2s_W^2} \frac{M_H^2}{M_W^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{\delta M_H^2}{M_H^2} \right].
\end{aligned} \tag{A.7}$$


SSS-coupling:



with the actual values of S_1, S_2, S_3 and C

$$\begin{aligned}
 HHH : \quad C &= -\frac{3}{2s} \frac{M_H^2}{m_W} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{3}{2} \delta Z_H \right], \\
 H\chi\chi : \quad C &= -\frac{1}{2s} \frac{M_H^2}{m_W} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_H \right], \\
 H\chi\chi : \quad C &= -\frac{1}{2s} \frac{M_H^2}{m_W} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta M_H^2}{M_H^2} - \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_H \right].
 \end{aligned}
 \tag{A.8}$$

$VVVV$ -couping:



with the actual values of V_1, V_2, S_1, S_2 and C

$$\begin{aligned}
W^+W^-HH : \quad C &= \frac{1}{2s^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \delta Z_W + \delta Z_H \right], \\
W^+W^-\chi\chi : \quad C &= \frac{1}{2s^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \delta Z_W \right], \\
W^+W^-\phi\phi : \quad C &= \frac{1}{2s^2} \left[1 + 2\delta Z_e - 2\frac{\delta s_W}{s_W} + \delta Z_W \right], \\
ZZ\phi^+\phi^- : \quad C &= \frac{s_W^2 - c_W^2}{2s_W^2 c_W^2} \left[1 + 2\delta Z_e + \frac{2}{(s_W^2 - c_W^2)c_W^2} \frac{\delta s_W}{s_W} + \delta Z_{ZZ} \right] \\
&\quad + \frac{s_W^2 - c_W^2}{s_W c_W} \frac{1}{2} \delta Z_{AZ}, \\
ZA\phi^+\phi^- : \quad C &= \frac{s_W^2 - c_W^2}{2s_W^2 c_W^2} \left[1 + 2\delta Z_e + \frac{1}{(s_W^2 - c_W^2)c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_{ZZ} \right. \\
&\quad \left. + \frac{1}{2} Z_{AA} \right] + \frac{1}{2} \frac{(s_W^2 - c_W^2)^2}{2s_W^2 c_W^2} \delta Z_{ZA} + \delta Z_{AZ}, \\
AA\phi^+\phi^- : \quad C &= 2[1 + 2\delta Z_e + \delta Z_{AA}] + \frac{1}{2} \frac{s_W^2 - c_W^2}{s_W c_W} \delta Z_{ZA}, \\
ZZHH : \quad C &= \frac{1}{2s_W^2 c_W^2} \left[1 + 2\delta Z_e + 2\frac{s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} + \delta Z_Z + \delta Z_H \right], \\
ZZ\chi\chi : \quad &= \frac{1}{2s_W^2 c_W^2} \left[1 + 2\delta Z_e + 2\frac{s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} + \delta Z_Z \right], \\
ZAHH : \quad C &= \frac{1}{2s_W^2 c_W^2} \frac{1}{2} \delta Z_{ZA}, \\
ZA\chi\chi : \quad C &= \frac{1}{2s_W^2 c_W^2} \frac{1}{2} \delta Z_{ZA},
\end{aligned}$$

$$\begin{aligned}
W^\pm Z \phi^\mp H : \quad C &= -\frac{1}{2c_w} \left[1 + \delta Z_e - \frac{\delta c_W}{c_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_H + \frac{1}{2} \delta Z_{ZZ} \right] \\
&\quad - \frac{1}{2} \frac{1}{s_W} \delta Z_{AZ}, \\
W^\pm A \phi^\mp H : \quad C &= -\frac{1}{2s_w} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_H + \frac{1}{2} \delta Z_{AA} \right] \\
&\quad - \frac{1}{2} \frac{1}{c_W} \delta Z_{ZA}, \\
W^\pm Z \phi^\mp \chi : \quad C &= \mp \frac{i}{2c_w} \left[1 + \delta Z_e - \frac{\delta c_W}{c_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_{ZZ} \right] \mp \frac{1}{2} \frac{i}{2s_W} \delta Z_{AZ}, \\
W^\pm Z \phi^\mp \chi : \quad C &= \mp \frac{i}{2s_w} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_{AA} \right] \mp \frac{1}{2} \frac{i}{2c_W} \delta Z_{ZA}.
\end{aligned} \tag{A.9}$$

VSS-coupling:

$$= ieC(k_1 - k_2)_\mu$$

with the actual values of V , S_1, S_2 and C

$$\begin{aligned}
A\chi H : \quad C &= -\frac{1}{2} \frac{i}{2c_w s_w} \delta Z_{ZA}, \\
Z\chi H : \quad C &= -\frac{i}{2c_w s_w} \left[1 + \delta Z_e + \frac{s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_H + \frac{1}{2} \delta Z_{ZZ} \right], \\
A\phi^+ \phi^- : \quad C &= -\left[1 + \delta Z_e + \frac{1}{2} \delta Z_{AA} + \frac{1}{2} \frac{s_W^2 - c_W^2}{2s_W c_W} \delta Z_{ZA} \right], \\
Z\phi^+ \phi^- : \quad C &= -\frac{s_w^2 - c_w^2}{2s_w c_w} \left[1 + \delta Z_e + \frac{1}{(s_W^2 - c_W^2) c_W^2} \frac{\delta s_w}{s_w} + \frac{1}{2} \delta Z_{ZZ} \right] - \frac{1}{2} \delta Z_{AZ}, \\
W^\pm \phi^\pm H : \quad C &= \mp \frac{1}{2s_w} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_H \right], \\
W^\pm \phi^\pm \chi : \quad C &= -\frac{i}{2s_w} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W \right].
\end{aligned} \tag{A.10}$$

SVV -coupling:

$$= ieg_{\mu\nu}C$$

with the actual values of S , V_1, V_2 and C

$$\begin{aligned}
HW^+W^- : \quad C &= M_W \frac{1}{s_W} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_H + \delta Z_W \right] \\
HZZ : \quad C &= M_W \frac{1}{s_W c_W^2} \left[1 + \delta Z_e + \frac{2s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_H + \delta Z_{ZZ} \right], \\
HZA : \quad C &= M_W \frac{1}{s_W c_W^2} \frac{1}{2} \delta Z_{ZA}, \\
\phi^\pm W^\mp Z : \quad C &= -M_W \frac{s_W}{c_W} \left[1 + \delta Z_e + \frac{1}{c_W} \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_{ZZ} \right], \\
\phi^\pm W^\mp A : \quad C &= -M_W \left[1 + \delta Z_e + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_W + \frac{1}{2} \delta Z_{AA} \right] - M_W \frac{s_W}{c_W} \frac{1}{2} \delta Z_{ZA}.
\end{aligned} \tag{A.11}$$

VFF -coupling:

$$= ie\gamma_\mu(C^-\omega_- + C^+\omega_+)$$

with the actual values of V , \bar{F}_1, F_2 and C^+ and C^-

$$\begin{aligned}
\gamma \bar{f}_i f_j : & \begin{cases} C^+ = -Q_f [\delta_{ij} (1 + \delta Z_e + \frac{1}{2} \delta Z_{AA}) + \frac{1}{2} (\delta Z_{ij}^{f,R} + \delta Z_{ij}^{f,R^\dagger})] + \delta_{ij} g_f^+ \frac{1}{2} \delta Z_{ZA}, \\ C^- = -Q_f [\delta_{ij} (1 + \delta Z_e + \frac{1}{2} \delta Z_{AA}) + \frac{1}{2} (\delta Z_{ij}^{f,L} + \delta Z_{ij}^{f,L^\dagger})] + \delta_{ij} g_f^- \frac{1}{2} \delta Z_{ZA}, \end{cases} \\
Z \bar{f}_i f_j : & \begin{cases} C^+ = g_f^+ [\delta_{ij} (1 + \frac{\delta g_f^+}{g_f^+} + \frac{1}{2} \delta Z_{ZZ}) + \frac{1}{2} (\delta Z_{ij}^{f,R} + \delta Z_{ij}^{f,R^\dagger})] - \delta_{ij} Q_f \frac{1}{2} \delta Z_{AZ}, \\ C^- = g_f^+ [\delta_{ij} (1 + \frac{\delta g_f^-}{g_f^-} + \frac{1}{2} \delta Z_{ZZ}) + \frac{1}{2} (\delta Z_{ij}^{f,L} + \delta Z_{ij}^{f,L^\dagger})] - \delta_{ij} Q_f \frac{1}{2} \delta Z_{AZ}, \end{cases} \\
W^+ \bar{u}_i d_j : & \begin{cases} C^+ = 0, \\ C^- = \frac{1}{\sqrt{2} s_W} \left[V_{ij} \left(1 + \delta Z_e - \frac{\delta S_W}{s_W} + \frac{1}{2} \delta Z_W \right) + \delta V_{ij} + \frac{1}{2} \sum_k (\delta Z_{jk}^{u,L^\dagger} V_{kj} + V_{ik} \delta Z_{kj}^{d,L}) \right], \end{cases} \\
W^- \bar{d}_j u_i : & \begin{cases} C^+ = 0, \\ C^- = \frac{1}{\sqrt{2} s_W} \left[V_{ji}^\dagger \left(1 + \delta Z_e - \frac{\delta S_W}{s_W} + \frac{1}{2} \delta Z_W \right) + \delta V_{ji}^\dagger + \frac{1}{2} \sum_k (\delta Z_{jk}^{d,L^\dagger} V_{ki}^\dagger + V_{jk}^\dagger \delta Z_{ki}^{u,L}) \right], \end{cases} \\
W^+ \bar{\nu}_i l_j : & \begin{cases} C^+ = 0, \\ C^- = \frac{1}{\sqrt{2} s_W} \delta_{ij} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} (\delta Z^{\nu,L^\dagger})_{ii} + \delta Z_{ii}^{\nu,L} \right], \end{cases} \\
W^- \bar{l}_j \nu_i : & \begin{cases} C^+ = 0, \\ C^- = \frac{1}{\sqrt{2} s_W} \delta_{ij} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W + \frac{1}{2} (\delta Z^{l,L^\dagger})_{ii} + \delta Z_{ii}^{l,L} \right], \end{cases} \tag{A.12}
\end{aligned}$$


where

$$\begin{aligned}
g_f^+ &= -\frac{s_W^0}{c_W} Q_f, & \delta g_f^+ &= -\frac{s_W}{c_W} Q_f \left[\delta Z_e + \frac{1}{c_W^2} \frac{\delta s_W}{s_W} \right], \\
g_f^- &= \frac{I_{W,f}^3 - s_W^2 Q_f}{s_W c_W}, & \delta g_f^+ &= \frac{I_{W,f}^2}{s_W c_W} Q_f \left[\delta Z_e + \frac{s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} \right] + \delta g_f^+.
\end{aligned} \tag{A.13}$$

The vector and axial vector couplings for the Z -boson are given by

$$v_f = \frac{1}{2}(g^- + g^+) = \frac{I_{W,f}^3 - 2s_W^2 Q_f}{2s_W c_W}, \quad a_f = \frac{1}{2}(g^- - g^+) = \frac{I_{W,f}^3}{2s_W c_W}. \tag{A.14}$$

SFF-coupling:




with the actual values of S , \bar{F}_1, F_2 and C^+ and C^-

$$\begin{aligned}
H\bar{f}_i f_j : & \left\{ \begin{aligned} C^+ &= -\frac{1}{2s_W} \frac{m_{f,i}}{M_W} \left[\delta_{ij} \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{f,i}}{m_{f,i}} - \frac{\delta M_W}{M_W} + \frac{1}{2} \delta Z_H \right) \right. \\ & \quad \left. + \frac{1}{2} (\delta Z_{ij}^{f,R} + \delta Z_{ij}^{f,R\dagger}) \right], \\ C^- &= -\frac{1}{2s_W} \frac{m_{f,i}}{M_W} \left[\delta_{ij} \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{f,i}}{m_{f,i}} - \frac{\delta M_W}{M_W} + \frac{1}{2} \delta Z_H \right) \right. \\ & \quad \left. + \frac{1}{2} (\delta Z_{ij}^{f,L} + \delta Z_{ij}^{f,L\dagger}) \right], \end{aligned} \right. \\
\chi\bar{f}_i f_j : & \left\{ \begin{aligned} C^+ &= i \frac{1}{2s_W} 2I_{W,f}^3 \frac{m_{f,i}}{M_W} \left[\delta_{ij} \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{f,i}}{m_{f,i}} - \frac{\delta M_W}{M_W} \right) \right. \\ & \quad \left. + \frac{1}{2} (\delta Z_{ij}^{f,R} + \delta Z_{ij}^{f,R\dagger}) \right], \\ C^- &= -i \frac{1}{2s_W} 2I_{W,f}^3 \frac{m_{f,i}}{M_W} \left[\delta_{ij} \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{f,i}}{m_{f,i}} - \frac{\delta M_W}{M_W} \right) \right. \\ & \quad \left. + \frac{1}{2} (\delta Z_{ij}^{f,L} + \delta Z_{ij}^{f,L\dagger}) \right], \end{aligned} \right. \\
\phi^+ \bar{u}_i d_j : & \left\{ \begin{aligned} C^+ &= -\frac{1}{\sqrt{2}s_W} \frac{m_{d,j}}{M_W} \left[V_{ij} \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{d,j}}{m_{d,j}} - \frac{\delta M_W}{M_W} + \delta V_{ij} \right) \right. \\ & \quad \left. + \frac{1}{2} \sum_k (\delta Z_{ik}^{u,R\dagger} V_{kj} + V_{ik} \delta Z_{kj}^{d,R}) \right], \\ C^- &= \frac{1}{\sqrt{2}s_W} \frac{m_{u,j}}{M_W} \left[V_{ij} \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{u,j}}{m_{u,j}} - \frac{\delta M_W}{M_W} + \delta V_{ij} \right) \right. \\ & \quad \left. + \frac{1}{2} \sum_k (\delta Z_{ik}^{u,L\dagger} V_{kj} + V_{ik} \delta Z_{kj}^{d,L}) \right], \end{aligned} \right.
\end{aligned}$$

$$\begin{aligned}
\phi^- \bar{d}_j u_i : & \left\{ \begin{aligned} C^+ &= \frac{1}{\sqrt{2}s_W} \frac{m_{u,i}}{M_W} \left[V_{ji}^\dagger \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{u,j}}{m_{u,j}} - \frac{\delta M_W}{M_W} + \delta V_{ji}^\dagger \right. \right. \\ & \left. \left. + \frac{1}{2} \sum_k (\delta Z_{jk}^{d,R\dagger} V_{ki}^\dagger + V_{jk}^\dagger \delta Z_{ki}^{u,R}) \right) \right], \\ C^- &= -\frac{1}{\sqrt{2}s_W} \frac{m_{d,i}}{M_W} \left[V_{ji}^\dagger \left(1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{d,j}}{m_{d,j}} - \frac{\delta M_W}{M_W} + \delta V_{ji}^\dagger \right. \right. \\ & \left. \left. + \frac{1}{2} \sum_k (\delta Z_{jk}^{d,L\dagger} V_{ki}^\dagger + V_{jk}^\dagger \delta Z_{ki}^{u,L}) \right) \right], \end{aligned} \right. \\
\phi^+ \bar{\nu}_i l_j : & \left\{ \begin{aligned} C^+ &= -\frac{1}{\sqrt{2}s_W} \frac{m_{i,j}}{M_W} \delta_{ij} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{i,j}}{m_{i,j}} - \frac{\delta M_W}{M_W} + \frac{1}{2} \left(\delta Z_{ii}^{\nu,R\dagger} + \delta Z_{ii}^{l,R} \right) \right], \\ C^- &= 0, \end{aligned} \right. \\
\phi^- \bar{l}_j \nu_i : & \left\{ \begin{aligned} C^+ &= 0, \\ C^- &= -\frac{1}{\sqrt{2}s_W} \frac{m_{i,j}}{M_W} \delta_{ij} \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{\delta m_{i,j}}{m_{i,j}} - \frac{\delta M_W}{M_W} + \frac{1}{2} \left(\delta Z_{ii}^{l,L\dagger} + \delta Z_{ii}^{\nu,L} \right) \right]. \end{aligned} \right.
\end{aligned} \tag{A.15}$$


VGG-coupling:



with the actual values of V , \bar{G}_1, G_2 and C

$$\begin{aligned}
A\bar{u}^\pm u^\pm : \quad C &= \pm \left[1 + \delta Z_e + \frac{1}{2} \delta Z_{AA} \right] \mp \frac{c_W}{s_W} \frac{1}{2} \delta Z_{AA}, \\
Z\bar{u}^\pm u^\pm : \quad C &= \mp \left[1 + \delta Z_e - \frac{1}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_{ZZ} \right] \pm \frac{1}{2} \delta Z_{AZ}, \\
W^\pm \bar{u}^\pm u^Z : \quad C &= \pm \left[1 + \delta Z_e - \frac{1}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W \right], \\
W^\pm \bar{u}^Z u^\mp : \quad C &= \mp \left[1 + \delta Z_e - \frac{1}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \delta Z_W \right], \\
W^\pm \bar{u}^\pm u^\gamma : \quad C &= \mp \left[1 + \delta Z_e + \frac{1}{2} \delta Z_W \right], \\
W^\pm \bar{u}^\gamma u^\pm : \quad C &= \pm \left[1 + \delta Z_e + \frac{1}{2} \delta Z_W \right].
\end{aligned} \tag{A.16}$$

SGG-coupling:



with the actual values of S , \bar{G}_1, G_2 and C

$$\begin{aligned}
H\bar{u}^Z u^Z : \quad C &= -\frac{1}{2s_W c_W^2} M_W \left[1 + \delta Z_e + \frac{2s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_H \right], \\
H\bar{u}^\pm u^\pm : \quad C &= -\frac{1}{2s_W} M_W \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} + \frac{1}{2} \delta Z_H \right], \\
\chi\bar{u}^\pm u^\pm : \quad C &= \mp i \frac{1}{2s_W} M_W \left[1 + \delta Z_e - \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} \right], \\
\phi^\pm \bar{u}^Z u^\mp : \quad C &= \frac{1}{2s_W c_W} M_W \left[1 + \delta Z_e + \frac{s_W^2 - c_W^2}{c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} \right], \\
\phi^\pm \bar{u}^\pm u^Z : \quad C &= \frac{s_W^2 - c_W^2}{2s_W c_W} M_W \left[1 + \delta Z_e + \frac{1}{(s_W^2 - c_W^2)c_W^2} \frac{\delta s_W}{s_W} + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} \right], \\
\phi^\pm \bar{u}^\pm u^\gamma : \quad C &= M_W \left[1 + \delta Z_e + \frac{1}{2} \frac{\delta M_W^2}{M_W^2} \right].
\end{aligned} \tag{A.17}$$

APPENDIX B

Feynman Rules of Quantum Chromodynamics

In this appendix we outline the Feynman rules of quantum chromodynamics including counterterms.

$$V_{\mu_1\mu_2\mu_3}(k_1, k_2, k_3) = (k_1 - k_2)_{\mu_3} g_{\mu_1\mu_2} + (k_2 - k_3)_{\mu_1} g_{\mu_2\mu_3} + (k_3 - k_1)_{\mu_2} g_{\mu_3\mu_1} \quad (\text{B.1})$$

$$\begin{aligned} W_{\mu_1\cdots\mu_4}^{a_1\cdots a_4} = & (f^{13,24} - f^{14,32}) g_{\mu_1\mu_2} g_{\mu_3\mu_4} + (f^{12,34} - f^{14,23}) g_{\mu_1\mu_3} g_{\mu_2\mu_4} \\ & + (f^{13,42} - f^{12,34}) g_{\mu_1\mu_4} g_{\mu_3\mu_2} \end{aligned} \quad (\text{B.2})$$

$$f^{ij,kl} = f^{a_i a_j a} f^{a k a l a} \quad (\text{B.3})$$

Gluons A

$$a\mu \cdot \text{---} \overset{k}{\text{---}} \text{---} b\nu = \delta_{ab} \frac{1}{k^2} \left(g_{\mu\nu} - (1 - \alpha) \frac{k_\mu k_\nu}{k^2} \right),$$

Faddeev-Popov ghosts χ

$$a \cdot \text{---} \overset{k}{\leftarrow} \text{---} \bar{b} = \delta_{ab} \frac{-1}{k^2},$$

Quark fields ψ

$$a \cdot \text{---} \overset{p}{\leftarrow} \text{---} b = \delta_{ij} \frac{1}{m_f - \not{p}}.$$

Gluon-counterterm

$$a\mu \cdot \text{---} \overset{k}{\text{---}} \otimes \text{---} b\nu = (Z_3 - 1) \delta_{ab} (k_\mu k_\nu - k^2 g_{\mu\nu})$$


Faddeev-Popov ghost counterterm

$$a \cdots \leftarrow \text{---} \otimes \text{---} \leftarrow \cdots b = (\tilde{Z}_3 - 1) \delta_{ab} k^2,$$

Fermion counterterm


$$i \cdots \leftarrow \text{---} \otimes \text{---} \leftarrow \cdots j = [(Z_2 - 1) \not{p} - (Z_2 Z_m - 1) m_R] \delta_{ij},$$

Three-gluon vertex




$$= -igf^{a_1 a_2 a_3} V_{\mu_1 \mu_2 \mu_3}(k_1, k_2, k_3)$$

Gluon-ghost-ghost vertex



$$= -igf^{a_1 a_2 a_3} k_\mu$$

Gluon-quark-quark vertex



$$= g \gamma_\mu T_{ij}^a$$

Four-gluon vertex

$$= -g^2 W_{\mu_1 \dots \mu_4}^{a_1 \dots a_4}$$

Three-gluon vertex counterterm

$$= (Z_1 - 1)(-i)g_R f^{a_1 a_2 a_3} V_{\mu_1 \mu_2 \mu_3}(k_1, k_2, k_3)$$


Gluon-ghost-ghost vertex counterterm

$$= (\tilde{Z}_1 - 1)(-i)g_R f^{abc} k_\mu$$

Gluon-quark-quark vertex

$$= (\tilde{Z}_{1F} - 1)g_R T_{ij}^a \gamma_\mu$$

Four-gluon vertex




$$= (Z_4 - 1)(-1)g_R^2 W_{\mu_1 \dots \mu_4}^{a_1 \dots a_4}.$$

The gluon loop




$$\int \frac{d^4 k}{(2\pi)^4} \delta^{ab} g^{\mu\nu}.$$

The ghost loop




$$- \int \frac{d^4 k}{(2\pi)^4} \delta^{ab}.$$

The quark loop




$$- \int \frac{d^4 k}{(2\pi)^4 i} \delta^{ij} \delta^{\alpha\beta}.$$

The gluon-quark loop




$$\int \frac{d^4 k}{(2\pi)^4 i}.$$

The gluon-ghost loop




$$\int \frac{d^4 k}{(2\pi)^4 i}.$$


Symmetry factors



$$\sim \frac{1}{2!}$$



$$\sim \frac{1}{2!}$$



$$\sim \frac{1}{3!}$$

APPENDIX C

The $SU(3)$ Group

The $SU(3)$ generators T_a are hermitian, traceless matrices which generate the closed $SU(3)$ algebra

$$[T_a, T_b] = if_{abc}T_c, \tag{C.1}$$

where f_{abc} are the antisymmetric $SU(3)$ structure constants with non-zero values given by

a	b	c	f_{abc}
1	2	3	1
1	4	7	$\frac{1}{2}$
1	5	6	$-\frac{1}{2}$
2	4	6	$\frac{1}{2}$
2	5	7	$\frac{1}{2}$
3	4	5	$\frac{1}{2}$
3	6	7	$-\frac{1}{2}$
4	5	8	$\frac{\sqrt{3}}{2}$
6	7	8	$\frac{\sqrt{3}}{2}$

A convenient representation of the T_a matrices is the one introduced by Gell-Mann [132] in which

$$\begin{aligned}
T_1 &= \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & T_2 &= \frac{1}{2} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\
T_3 &= \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & T_4 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\
T_5 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, & T_6 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \\
T_7 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, & T_8 &= \frac{1}{2\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}.
\end{aligned} \tag{C.2}$$

The fundamental representation is 3-dimensional where T_a satisfy the relation,

$$\{T_a, T_b\} = \frac{1}{3}\delta_{ab} + d_{abc}T_c, \tag{C.3}$$

which is consistent with the normalization

$$\text{Tr}[T_a T_b] = \frac{1}{2}\delta_{ab}. \tag{C.4}$$

Here d_{abc} is totally symmetric in a , b and c and is given by

$$d_{abc} = 2\text{Tr}[\{T_a, T_b\}T_c]. \tag{C.5}$$

According to eqs. (C.1) and (C.3), we find that

$$T_a T_b = \frac{1}{2} \left[\frac{1}{3}\delta_{ab} + (d_{abc} + if_{abc})T_c \right]. \tag{C.6}$$

The traces of the products of generators T_a in the fundamental representation are given by

$$\text{Tr}[T_a T_b T_c] = \frac{1}{4}(d_{abc} + if_{abc}), \quad (\text{C.7})$$

$$\text{Tr}[T_a T_b T_c T_d] = \frac{1}{12}\delta_{ab}\delta_{cd} + \frac{1}{8}(d_{abe} + if_{abe})(d_{cde} + if_{cde}) \quad (\text{C.8})$$

In the adjoint representation the generator T_a is a 8×8 matrix and its matrix element reads

$$(T_a)_{bc} = -if_{abc}. \quad (\text{C.9})$$

The traces of products of generators T^a yield

$$\text{Tr}[T_a T_b T_c] = \frac{3}{2}if_{abc}, \quad (\text{C.10})$$

$$\text{Tr}[T_a T_b T_c T_d] = \delta_{ab}\delta_{cd} + \delta_{ad}\delta_{bc} + \frac{3}{4}(d_{abe}d_{cde} - d_{ace}d_{bde} + d_{ade}d_{bce}). \quad (\text{C.11})$$

The eq. (C.10) reduces to

$$f_{ade}f_{bef}f_{cfd} = \frac{3}{2}f_{abc}. \quad (\text{C.12})$$

The Jacob identities

$$[T_a, [T_b, T_c]] + \text{cyclic permutations} = 0, \quad (\text{C.13})$$

$$[T_a, \{T_b, T_c\}] + \text{cyclic permutations} = 0, \quad (\text{C.14})$$

leads to the following relations:

$$f_{abe}f_{cde} + f_{cbe}f_{dae} + f_{dbe}f_{ace} = 0, \quad (\text{C.15})$$

$$f_{abe}f_{cde} + f_{cbe}f_{dae} + f_{dbe}f_{ace} = 0. \quad (\text{C.16})$$

APPENDIX D

The Global Positioning of Spin GPS Scheme

In Chapter Three, we introduced the Kleiss and Stirling spinor method. In the KS scheme, the massless spinors and massive spinors are defined [52]. The definitions in the KS scheme will be supplemented in Ref. [54] with the precise prescription of the spin quantization axes, the translation from spin amplitudes to density matrices, and the methodology of connecting production and decay for unstable fermions.

The GPS rules determining the spin quantization frame for the $u(p, \pm)$ and $v(p, \pm)$ of eq. (3.99) are summarized as follows:

- (i) In the rest frame of the fermion, take the z -axis along $-\vec{k}$.
- (ii) Place the x axis in the plane define by the z -axis from the previous point and the vector $\vec{\eta}$, in the same half-plane as $\vec{\eta}$.
- (iii) With the y -axis, complete the right-handed system of coordinates. The rest frame defined in this way we call the GPS frame of the particular fermion.

Next we will assume that polarization vectors of beams and of outgoing fermions are defined in their corresponding GPS frames.

For the definitions of inner product of the spinors are the same as those described in Chapter Three.

For a circularly polarization vector with four-momentum k and helicity $\sigma = \pm 1$ we take the following convention [51]:

$$[\epsilon_{\sigma}^{\mu}(\beta)]^{*} = \frac{\bar{u}_{\sigma}(k)\gamma^{\mu}u_{\sigma}(\beta)}{\sqrt{2}u_{-\sigma}(k)u_{\sigma}(\beta)}$$

$$[\epsilon_\sigma^\mu(\zeta)]^* = \frac{\bar{u}_\sigma(k)\gamma^\mu u_\sigma(\zeta)}{\sqrt{2}u_{-\sigma}(k)u_\sigma(\zeta)} \quad (\text{D.1})$$

where β is an arbitrary light-like four-vector $\beta^2 = 0$. The second choice with $u_\sigma(\zeta)$ (constant basic spinors) often simplifies the resulting photon emission amplitudes.

With the help of the Chisholm identity

$$\bar{u}_\sigma(k)\gamma_\mu u_\sigma(\beta)\gamma^\mu = 2u_\sigma(\beta)\bar{u}_{-\sigma}(k) + 2u_\sigma(k)\bar{u}_{-\sigma}(\beta), \quad (\text{D.2})$$

$$\bar{u}_\sigma(k)\gamma_\mu u_\sigma(\zeta)\gamma^\mu = 2u_\sigma(\zeta)\bar{u}_{-\sigma}(k) + 2u_\sigma(k)\bar{u}_{-\sigma}(\zeta), \quad (\text{D.3})$$

we obtain two useful formula, equivalent to eq. (D.1)

$$\begin{aligned} \not{\epsilon}_\sigma^*(k, \beta) &= \frac{\sqrt{2}}{\bar{u}_{-\sigma}(k)u_\sigma(\beta)} [u_\sigma(\beta)\bar{u}_{-\sigma}(k) + u_\sigma(k)\bar{u}_{-\sigma}(\beta)], \\ \not{\epsilon}_\sigma^*(k, \zeta) &= \frac{\sqrt{2}}{\sqrt{2}\zeta k} [u_\sigma(\zeta)\bar{u}_{-\sigma}(k) - u_\sigma(k)\bar{u}_{-\sigma}(\zeta)]. \end{aligned} \quad (\text{D.4})$$

While calculating photon emission spin amplitudes, we will use the following important building blocks, i.e., the elements of the ‘‘transition matrices’’ U and V defined as

$$\begin{aligned} \bar{u}(p_1, \lambda_1)\not{\epsilon}_\sigma^*(k, \beta)u(p_2, \lambda_2) &= U \begin{pmatrix} k \\ \sigma \end{pmatrix} \begin{bmatrix} p_1 p_2 \\ \lambda_1 \lambda_2 \end{bmatrix} = U_{\lambda_1, \lambda_2}^\sigma(k, p_1, m_1, p_2, m_2), \\ \bar{v}(p_1, \lambda_1)\not{\epsilon}_\sigma^*(k, \zeta)v(p_2, \lambda_2) &= V \begin{pmatrix} k \\ \sigma \end{pmatrix} \begin{bmatrix} p_1 p_2 \\ \lambda_1 \lambda_2 \end{bmatrix} = V_{\lambda_1, \lambda_2}^\sigma(k, p_1, m_1, p_2, m_2). \end{aligned} \quad (\text{D.5})$$

In the case of $u_\sigma(\zeta)$ the above transition matrices reads

$$U^+(k, p_1, m_1, p_2, m_2) = \sqrt{2} \begin{bmatrix} \sqrt{\frac{2\zeta p_2}{2\zeta k}} s_+(k, \hat{p}_1), & 0 \\ m_2 \sqrt{\frac{2\zeta p_1}{2\zeta p_2}} - m_1 \sqrt{\frac{2\zeta p_1}{2\zeta p_2}}, & \sqrt{\frac{2\zeta p_1}{2\zeta k}} s_+(k, \hat{p}_2) \end{bmatrix}, \quad (\text{D.6})$$

$$U_{\lambda_1, \lambda_2}^-(k, p_1, m_1, p_2, m_2) = [-U_{\lambda_2, \lambda_1}^+(k, p_2, m_2, p_1, m_1)]^*, \quad (\text{D.7})$$

$$V_{\lambda_1, \lambda_2}^\sigma(k, p_1, m_1, p_2, m_2) = -U_{-\lambda_1, -\lambda_2}^\sigma(k, p_1, -m_1, p_2, -m_2). \quad (\text{D.8})$$

Compared with the case of $u_\sigma(\zeta)$, the more general case $u(\beta)$ is a little bit more complicated

$$\begin{aligned}
& U^+(k, p_1, m_1, p_2, m_2) \\
&= \sqrt{\frac{2}{s_-(k, \beta)}} \times \\
& \left[\begin{array}{cc}
s_+(\hat{p}_1, k) s_-(\beta, \hat{p}_2) + m_1 m_2 \sqrt{\frac{2\zeta\beta}{2\zeta p_1} \frac{2\zeta k}{2\zeta p_2}}, & m_1 \sqrt{\frac{2\zeta\beta}{2\zeta p_1}} s_+(k, \hat{p}_2) + m_2 \sqrt{\frac{2\zeta\beta}{2\zeta p_2}} s_+(\hat{p}_1, k) \\
m_1 \sqrt{\frac{2\zeta k}{2\zeta p_1}} s_-(\beta, \hat{p}_2) + m_2 \sqrt{\frac{2\zeta k}{2\zeta p_2}} s_-(\hat{p}_1, \beta), & s_-(\hat{p}_1, \beta) s_+(k, \hat{p}_2) + m_1 m_2 \sqrt{\frac{2\zeta\beta}{2\zeta p_1} \frac{2\zeta k}{2\zeta p_2}}
\end{array} \right].
\end{aligned} \tag{D.9}$$

The numbering of elements in matrices U and V is

$$\{(\lambda_1, \lambda_2)\} = \begin{bmatrix} (++) & (+-) \\ (-+) & (--) \end{bmatrix}. \tag{D.10}$$

When computing bremsstrahlung amplitudes we will adopt the following compact notation:

$$\begin{aligned}
U \begin{bmatrix} pkp \\ \lambda_1 \sigma \lambda_2 \end{bmatrix} &\equiv U_{\lambda_1, \lambda_2}^\sigma(k, p_1, m_1, p_2, m_2) \\
V \begin{bmatrix} pkp \\ \lambda_1 \sigma \lambda_2 \end{bmatrix} &\equiv V_{\lambda_1, \lambda_2}^\sigma(k, p_1, m_1, p_2, m_2).
\end{aligned} \tag{D.11}$$

When dealing with the soft real photon limit we will implement the following important diagonality property:

$$U \begin{bmatrix} pkp \\ \lambda_1 \sigma \lambda_2 \end{bmatrix} = V \begin{bmatrix} pkp \\ \lambda_1 \sigma \lambda_2 \end{bmatrix} = b_\sigma(k, p) \delta_{\lambda_1, \lambda_2}, \tag{D.12}$$

$$b_\sigma(k, p) = \sqrt{2} \frac{\bar{u}(k) \not{p} u_\sigma(\zeta)}{\bar{u}(k) u_\sigma(\zeta)} = \sqrt{2} \sqrt{\frac{2\zeta p}{2\zeta k}} s_\sigma(k, \hat{p}), \tag{D.13}$$

which also holds in the general case of $u_\sigma(\beta)$, where

$$b_\sigma(k, p) = \frac{\sqrt{2}}{s_{-\sigma}(k, \beta)} \left(s_{-\sigma}(\beta, \hat{\beta}) s_\sigma(\hat{p}, k) + \frac{m^2}{2\zeta\hat{p}} \sqrt{(2\beta\zeta)(2\zeta k)} \right). \quad (\text{D.14})$$

APPENDIX E

The Drell-Yan Process

The Drell-Yan process is a model for the production of massive lepton pair in hadron-hadron collision developed by Drell and Yan in 1970 [93]. In the model a quark from one incident hadron annihilates with an antiquark from the other hadron incident hadron producing a virtual gauge boson which in turn decays into a massive lepton pair. It provides many interesting tests of perturbative QCD. We will make a brief introduction of Drell-Yan process here [133].

First, let us begin with the parton model, in which large mass muon pairs are created in the proton-proton collision via the subprocess $q + \bar{q} \rightarrow \gamma^* \rightarrow \mu^+ + \mu^-$. The experimental cross section reads as follows

$$d\sigma = G_{p \rightarrow q}(x_a) dx_a G_{p \rightarrow \bar{q}}(x_b) dx_b \hat{\sigma}(q + \bar{q} \rightarrow \gamma^* \rightarrow \mu^+ + \mu^-), \quad (\text{E.1})$$

where $G_{p \rightarrow q}(x_a) dx_a$ is the probability of finding a quark with momentum

$$p_q = x_a P_A, \quad (\text{E.2})$$

and $G_{p \rightarrow \bar{q}}(x_b) dx_b$ is the probability of finding a quark with momentum

$$p_{\bar{q}} = x_b P_B, \quad (\text{E.3})$$

where P_A and P_B are the momentum of the initial two protons. It is convenient to define the dimensionless variables


$$\tau = \frac{M^2}{s}, \quad \hat{\tau} = \frac{M^2}{\hat{s}}, \quad (\text{E.4})$$

where M is the mass of the muon pair and where s is the external proton-proton CMS energy squared

$$s = (P_A + P_B)^2 = 2P_{\text{CM}}^2, \quad (\text{E.5})$$

and \hat{s} is the internal parton parton CMS energy squared

$$\hat{s} = (p_q + p_{\bar{q}})^2 = 2p_q \cdot p_{\bar{q}}. \quad (\text{E.6})$$



$$p + p \rightarrow \mu^+ \mu^- + X$$




Figure. E.1. The proton-proton collisions (Drell Yan process), $p + p \rightarrow \gamma^* + X$

Then we have

$$\hat{s} = x_a x_b s, \quad \tau = x_a x_b \hat{t}. \quad (\text{E.7})$$

The Longitudinal momentum of the muon pair are

$$P_L = p_q - p_{\bar{q}}, \quad (\text{E.8})$$

and if we assume that the incoming partons are parallel to the incident protons then the total energy is

$$E^2 = P_L^2 + M^2. \quad (\text{E.9})$$

eq. (E.8) leads to

$$x_L = x_a - x_b \quad (\text{E.10})$$

where

$$x_L \equiv \frac{2P_L}{\sqrt{s}}, \quad (\text{E.11})$$

and eq. (E.9) implies

$$x_E^2 = x_L^2 + 4\tau \quad (\text{E.12})$$

where

$$x_E \equiv \frac{2E}{\sqrt{s}}. \quad (\text{E.13})$$

The total cross section for a quark and anti-quark to annihilate into a muon pair, $q\bar{q} \rightarrow \mu^+\mu^-$ reads

$$\hat{\sigma}(q\bar{q} \rightarrow \mu^+\mu^-) \equiv \sigma_0 = \frac{1}{3} \frac{4\pi\alpha e_q^2}{3M^2}, \quad (\text{E.14})$$

where M is the virtual photon invariant mass with

$$\hat{s} = M^2. \quad (\text{E.15})$$

According to eq. (E.7) and (E.10) we find that x_a and x_b can be specified in terms of τ and x_L ,

$$x_a x_b = \tau, \quad (\text{E.16})$$

$$x_a - x_b = x_L, \quad (\text{E.17})$$

and the experimental cross section can be rewritten as

$$\frac{d\sigma_{\text{DY}}}{d\tau dx_L}(s, M^2, x_L) = \frac{4\pi\alpha^2}{9M^2} \frac{1}{(x_a + x_b)} P_{q\bar{q}}(x_a, x_b), \quad (\text{E.18})$$

with the joint $q\bar{q}$ probability function

$$P_{q\bar{q}}(x_a, x_b) = \sum_{i=1}^{n_f} e_{q_i}^2 [G_{p \rightarrow q_i}(x_a) G_{p \rightarrow \bar{q}_i}(x_b) + G_{p \rightarrow \bar{q}_i}(x_a) G_{p \rightarrow q_i}(x_b)], \quad (\text{E.19})$$

where the subscript DY denotes the "Drell-Yan" process $pp \rightarrow \mu^+ \mu^- + X$. And eqs. (E.16) and (E.17) lead to

$$x_a = \frac{1}{2}(x_E + x_L) = \sqrt{\tau} e^y, \quad (\text{E.20})$$

$$x_b = \frac{1}{2}(x_E - x_L) = \sqrt{\tau} e^{-y}, \quad (\text{E.21})$$

where y is the rapidity of the muon pair defined by

$$y \equiv \frac{1}{2} \log \left(\frac{E + p_L}{E - p_L} \right). \quad (\text{E.22})$$

Next, we consider the possibility that the initial quark or antiquark can radiate a gluon before annihilating into a virtual photon. The differential cross section for the subprocess $q + \bar{q} \rightarrow \gamma^* + g$ is

$$\begin{aligned} \frac{d\hat{\sigma}_{\text{DY}}^q}{d\hat{t}}(\hat{s}, \hat{t}) &= \frac{1}{64\pi\hat{s}\hat{p}_{\text{CM}}^2} |\bar{\mathcal{M}}(q + \bar{q} \rightarrow \gamma^* + g)|^2 \\ &= \frac{1}{16\pi\hat{s}^2} |\bar{\mathcal{M}}(q + \bar{q} \rightarrow \gamma^* + g)|^2. \end{aligned} \quad (\text{E.23})$$

and the amplitude squared is

$$|\bar{\mathcal{M}}(q + \bar{q} \rightarrow \gamma^* + g)|^2 = e^2 e_q^2 g^2 \frac{4}{9} \frac{1}{4} 8 \left[\frac{\hat{u}}{\hat{t}} + \frac{\hat{t}}{\hat{u}} + \frac{2M^2(M^2 - \hat{t} - \hat{u})}{\hat{t}\hat{u}} \right], \quad (\text{E.24})$$

where the superscript q denotes the process $q + \bar{q} \rightarrow \gamma^* + g$ the invariant mass of the virtual is timelike,

$$M^2 = q_\gamma^2, \quad (\text{E.25})$$




Figure E.2. Leading order diagrams for the subprocess $q + \bar{q} \rightarrow \gamma^* + g$.

The invariants are given by

$$\begin{aligned} \hat{s} &= (p_q + p_{\bar{q}})^2, \\ \hat{t} &= (q_\gamma - p_q)^2, \\ \hat{u} &= (q_g + p_q)^2, \end{aligned} \quad (\text{E.26})$$

with

$$\hat{s} + \hat{t} + \hat{u} = M^2. \quad (\text{E.27})$$

Therefore we have

$$\frac{d\hat{\sigma}_{\text{DY}}^q}{d\hat{t}}(\hat{s}, \hat{t}) = \frac{\pi\alpha\alpha_s e_q^2}{\hat{s}^2} \left[\frac{\hat{u}}{\hat{t}} + \frac{\hat{t}}{\hat{u}} + \frac{2M^2(M^2 - \hat{t} - \hat{u})}{\hat{t}\hat{u}} \right]. \quad (\text{E.28})$$

The integral over \hat{t} is given by

$$\hat{\sigma}_{\text{DY}}^q(\hat{s}) = \int_{\hat{t}_{\min}}^{\hat{t}_{\max}} \frac{d\hat{\sigma}_{\text{DY}}^q}{d\hat{t}}(\hat{s}, \hat{t}) d\hat{t}, \quad (\text{E.29})$$

where

$$\hat{t}_{\min} = 0, \quad \hat{t}_{\max} = M^2 - \hat{s} = -(1 - \hat{\tau})\hat{s}. \quad (\text{E.30})$$

Besides we have to include the "Compton" subprocess $q + g \rightarrow \gamma^* + q$ for correcting the parton model. The corresponding differential cross section

$$\frac{d\hat{\sigma}_{\text{DY}}^g}{d\hat{t}}(\hat{s}, \hat{t}) = \frac{1}{16\pi\hat{s}^2} |\bar{\mathcal{M}}(q + g \rightarrow \gamma^* + q)|^2, \quad (\text{E.31})$$

where

$$|\bar{\mathcal{M}}(q + g \rightarrow \gamma^* + q)|^2 = e^2 e_q^2 g_s^2 \frac{4}{24} \frac{1}{4} 8 \left[-\frac{\hat{t}}{\hat{s}} - \frac{\hat{s}}{\hat{t}} + \frac{2M^2(M^2 + \hat{s} + \hat{t})}{\hat{s}\hat{t}} \right]. \quad (\text{E.32})$$




Figure E.3. Leading order diagrams for the "Compton" subprocess $q + g \rightarrow \gamma^* + q$.

The invariants here are defined by

$$\begin{aligned} \hat{s} &= (p_q + q_g)^2, \\ \hat{t} &= (q_\gamma - p_q)^2, \\ \hat{u} &= (q_\gamma - q_g)^2. \end{aligned} \quad (\text{E.33})$$

Inserting eq. (E.32) into eq. (E.31) yields

$$\frac{d\hat{\sigma}_{\text{DY}}^g}{d\hat{t}}(\hat{s}, \hat{t}) = \frac{\pi\alpha\alpha_s e_q^2}{\hat{s}^2} \frac{1}{3} \left[-\frac{\hat{t}}{\hat{s}} - \frac{\hat{s}}{\hat{t}} + \frac{2M^2(M^2 + \hat{s} + \hat{t})}{\hat{s}\hat{t}} \right], \quad (\text{E.34})$$

and the integral over \hat{t} is

$$\hat{\sigma}_{\text{DY}}^g(\hat{s}) = \int_{\hat{t}_{\min}}^{\hat{t}_{\max}} \frac{d\hat{\sigma}_{\text{DY}}^g(\hat{s}, \hat{t})}{d\hat{t}} d\hat{t}. \quad (\text{E.35})$$

Note that eqs. (E.29) and (E.35) are divergent with $\hat{t}_{\min} = 0$ and we must regulate the divergences. The divergences can be regulated either by giving the gluon a fictitious mass $q_g^2 = m_g^2$ or by using the dimensional regularization. In the following we apply the dimensional regularization.

Let us consider the 2-to-2 scattering subprocesses $\gamma^* + q \rightarrow q + g$ and $\gamma^* + g \rightarrow q + \bar{q}$ occur in the N rather than 4 spacetime dimension. In the N spacetime dimensions the 2-to-2 cross section is given by

$$d\hat{\sigma} = \frac{1}{4(p_1 \cdot p_2)} |\bar{\mathcal{M}}|^2 d^{2N-2}R_2, \quad (\text{E.36})$$

where

$$d^{2N-2}R_2 = \frac{d^{N-1}p_3}{(2\pi)^{N-1}(2E_3)} \frac{d^{N-1}p_4}{(2\pi)^{N-1}(2E_4)} (2\pi)^N \delta^N(p_3 + p_4 - p_1 - p_2). \quad (\text{E.37})$$

Integrating over p_4 gives

$$\int d^{N-1}p_4 \delta^N(p_3 + p_4 - p_1 - p_2) = \delta(E_3 + E_4 - E_1 - E_2). \quad (\text{E.38})$$

Now let $y \equiv \cos \theta_{13}$, where θ_{13} is the scattering angle between particles 1 and 3 then

$$d^{N-1}p_3 = \frac{2\pi^{\frac{N-2}{2}}}{\Gamma(\frac{N}{2} - 1)} p_3^{N-2} dp_3 (1 - y^2)^{\frac{N-4}{2}} dy. \quad (\text{E.39})$$

Integrating over p_3 yields

$$\int dp_3 \frac{1}{4E_3 E_4} p_3^{N-2} \delta(E_3 + E_4 - E_{\text{CM}}) = \frac{(\hat{p}'_{\text{CM}})^{N-3}}{4\sqrt{\hat{s}}}, \quad (\text{E.40})$$

where

$$(\hat{p}'_{\text{CM}})^2 = \frac{1}{4\hat{s}} [\hat{s} - (m_3 + m_4)^2][\hat{s} - (m_3 - m_4)^2], \quad (\text{E.41})$$

and

$$p_1 \cdot p_2 = \sqrt{s} \hat{p}_{\text{CM}}, \quad (\text{E.42})$$

with

$$(\hat{p}_{\text{CM}})^2 = \frac{1}{4\hat{s}} [\hat{s} - (m_1 + m_2)^2] [\hat{s} - (m_1 - m_2)^2], \quad (\text{E.43})$$

Thus we have

$$\frac{d\hat{\sigma}}{dy}(\hat{s}, \hat{t}) = \frac{1}{32\pi\hat{s}} \frac{(\hat{p}'_{\text{CM}})^{N-3}}{\hat{p}_{\text{CM}}} |\bar{\mathcal{M}}|^2 \frac{(1-y^2)^{\frac{N-4}{2}}}{2^{N-4} \pi^{\frac{N-4}{2}} \Gamma(\frac{N}{2}-1)}. \quad (\text{E.44})$$

For the case,

$$\hat{p}_{\text{CM}} = \frac{1}{2} \sqrt{\hat{s}}, \quad (\text{E.45})$$

$$\hat{p}'_{\text{CM}} = \frac{1}{2} (1 - \hat{\tau}) \sqrt{\hat{s}}, \quad (\text{E.46})$$

we have

$$\hat{\sigma}_{\text{DY}}(\hat{\tau}) = \frac{1 - \hat{\tau}}{32\pi\hat{s}} \left(\frac{M^2(1 - \hat{\tau}^2)^2}{4\pi\hat{\tau}} \right)^{\frac{\epsilon}{2}} \frac{I}{2^\epsilon \Gamma(1 + \frac{\epsilon}{2})}, \quad (\text{E.47})$$

where

$$I = \int_{-1}^1 dy (1 - y^2)^{\frac{\epsilon}{2}} |\bar{\mathcal{M}}|^2, \quad (\text{E.48})$$

with $N = 4 + \epsilon$ and

$$\hat{t} = -\frac{1}{2} (\hat{s} - M^2)(1 - y) = -\frac{\hat{s}}{2} (1 - \hat{\tau})(1 - y), \quad (\text{E.49})$$

$$\hat{u} = -\frac{1}{2} (\hat{s} - M^2)(1 + y) = -\frac{\hat{s}}{2} (1 - \hat{\tau})(1 + y), \quad (\text{E.50})$$

In $N = 4 + \epsilon$ dimensions the matrix element squared for the subprocesses $q + \bar{q} \rightarrow \gamma^* + g$ reads

$$\begin{aligned} |\bar{\mathcal{M}}(q + \bar{q} \rightarrow \gamma^* + g)|^2 &= 16\pi^2 \alpha_N^{\text{QED}} \alpha_N^{\text{QCD}} e_q^2 \frac{8}{9} \left(1 + \frac{\epsilon}{2}\right) \\ &\times \left\{ \frac{2(\hat{\tau}^2 y^2 - 2\hat{\tau} y^2 + y^2 + \hat{\tau}^2 + 2\hat{\tau} + 1)}{(1 - \hat{\tau})^2 (1 - y^2)} + \frac{2}{1 - y^2} \epsilon \right\}, \quad (\text{E.51}) \end{aligned}$$

where

$$\alpha_N^{\text{QED}} = \frac{\alpha}{(m_D^2)^{\epsilon/2}}, \quad \alpha_N^{\text{QCD}} = \frac{\alpha_s}{(m_D^2)^{\epsilon/2}}, \quad (\text{E.52})$$

and m_D is the "dimensional regularization mass". Thus we arrive at

$$\begin{aligned} \sigma_{\text{DY}}^q(\hat{\tau}) &= \frac{\pi \alpha_N^{\text{QED}} \alpha_s e_q^2}{\hat{s}} \frac{16}{9} \left(\frac{M^2(1-\hat{\tau})^2}{\hat{\tau} 4\pi m_D^2} \right)^{\frac{\epsilon}{2}} \frac{\Gamma(1+\frac{\epsilon}{2})}{\Gamma(1+\epsilon)} \left(1+\frac{\epsilon}{2}\right) \\ &\times \left\{ \frac{1+\hat{\tau}^2}{1-\hat{\tau}} \frac{2}{\epsilon} + \frac{\epsilon(1-\hat{\tau})}{1+\epsilon} \right\} \end{aligned} \quad (\text{E.53})$$

Since

$$\left(\frac{1}{\sigma_0} \frac{d\hat{\sigma}^q}{d\hat{\tau}} \right) = \frac{3}{4\pi^2 \alpha e_q^2 (1+\frac{\epsilon}{2})} \hat{s} \hat{\sigma}_{\text{DY}}^q, \quad (\text{E.54})$$

then we have

$$\left(\frac{1}{\sigma_0} \frac{d\hat{\sigma}^q}{d\hat{\tau}} \right)_{\text{DY}} = 2 \frac{2\alpha_s}{3\pi} \left(\frac{(1-\hat{\tau})^2 M^2}{\hat{\tau} 4\pi m_D^2} \right)^{\frac{\epsilon}{2}} \frac{\Gamma(1+\frac{\epsilon}{2})}{\Gamma(1+\epsilon)} \left\{ \frac{1+\hat{\tau}^2}{1-\hat{\tau}} \frac{2}{\epsilon} + \frac{\epsilon(1-\hat{\tau})}{1+\epsilon} \right\}, \quad (\text{E.55})$$

where σ_0 is N -dimensional Born cross section. Integrating over $\hat{\tau}$ yields

$$(\hat{\sigma}(\text{real}))_{\text{DY}} = \frac{2\alpha_s}{3\pi} \sigma_0 \left(\frac{M^2}{4\pi m_D^2} \right)^{\frac{\epsilon}{2}} \Gamma\left(1-\frac{\epsilon}{2}\right) \left\{ \frac{8}{\epsilon^2} - \frac{6}{\epsilon} + \frac{9}{2} + \dots \right\}. \quad (\text{E.56})$$




Figure E.4. Virtual gluon corrections to the quark-antiquark annihilation $q + \bar{q} \rightarrow \gamma^*$.

The virtual corrections are given by

$$\begin{aligned}
(\hat{\sigma}(\text{virtual}))_{\text{DY}} &= \frac{2\alpha_s}{3\pi}\sigma_0 \left(\frac{M^2}{4\pi m_D^2} \right)^{\frac{\epsilon}{2}} \frac{\Gamma(1 - \frac{\epsilon}{2})\Gamma^2(1 + \frac{\epsilon}{2})}{\Gamma(1 + \epsilon)} \\
&\times \left\{ -\frac{8}{\epsilon^2} + \frac{6}{\epsilon} - 8 + \pi^2 + \dots \right\}.
\end{aligned} \tag{E.57}$$

From the expansions

$$\begin{aligned}
\frac{\Gamma(1 - \frac{\epsilon}{2})\Gamma^2(1 + \frac{\epsilon}{2})}{\Gamma(1 + \epsilon)} &= 1 + \frac{1}{2}\gamma_E\epsilon + \frac{1}{48}(6\gamma_E^2 - \pi^2)\epsilon^2 + \dots, \\
\Gamma\left(1 - \frac{\epsilon}{2}\right) &= 1 + \frac{1}{2}\gamma_E\epsilon + \frac{1}{8}\left(\frac{\pi^2}{6} + \gamma_E^2\right)\epsilon^2 + \dots,
\end{aligned} \tag{E.58}$$

where γ_E is the Euler constant, we find that

$$(\hat{\sigma}(\text{real}) + \hat{\sigma}(\text{virtual})) = \sigma_0\alpha_s \left(\frac{8\pi}{9} - \frac{7}{3\pi} \right). \tag{E.59}$$

For the Drell-Yan case, the perturbation series behaves like

$$\sigma_{\text{tot}}^{\text{DY}} = \sigma_0(1 + \alpha_s I_q^{\text{DY}} + \dots) \tag{E.60}$$

with

$$\alpha_s I_q^{\text{DY}} = \alpha_s \left(\frac{8\pi}{9} - \frac{7}{3\pi} \right). \tag{E.61}$$

We now define " + functions" and have

$$\frac{1}{\sigma_0} \left(\frac{d\hat{\sigma}_{\text{DY}}^q}{d\hat{\tau}} \right)_+ = 2\frac{\alpha_s}{2\pi} P_{q \rightarrow qg}(\hat{\tau}) \log\left(\frac{M^2}{m_D^2}\right) + 2\alpha_s f^{q,\text{DY}}(\hat{\tau}), \tag{E.62}$$

where the splitting function

$$P_{q \rightarrow qg}(\hat{\tau}) = \frac{4}{3} \left(\frac{1 + \hat{\tau}^2}{1 - \hat{\tau}} \right), \tag{E.63}$$

and

$$\begin{aligned}
\alpha_s f^{q,\text{DY}}(\hat{\tau}) &= \frac{2\alpha_s}{3\pi} \left\{ 2(1 + \hat{\tau}^2) \left(\frac{\log(1 - \hat{\tau})}{1 - \hat{\tau}} \right)_+ - \frac{1 + \hat{\tau}^2}{1 - \hat{\tau}} \log(\hat{\tau}) \right. \\
&\left. - \left(\frac{\pi^2}{3} + \frac{9}{4} \right) \delta(1 - \hat{\tau}) \right\} + \frac{\alpha_s}{2\pi} P_{q \rightarrow qg}(\hat{\tau}) \left(\frac{2}{\epsilon} + \gamma_E - \log(4\pi) \right).
\end{aligned} \tag{E.64}$$

Note that the "little f " functions is regularization scheme dependent and the integral of $f^{q,\text{DY}}$ over $\hat{\tau}$ vanishes,

$$\int_0^1 \alpha_s f^{q,\text{DY}}(\hat{\tau}) d\hat{\tau} = 0 \quad (\text{E.65})$$

For the "Compton" subprocess $q + g \rightarrow \gamma^* + q$, we take the similar treatment and then have

$$\frac{1}{\sigma_0} \left(\frac{d\hat{\sigma}_{\text{DY}}^g}{d\hat{\tau}} \right)_+ = 2 \frac{\alpha_s}{2\pi} P_{g \rightarrow q\bar{q}}(\hat{\tau}) \log \left(\frac{M^2}{m_D^2} \right) + 2\alpha_s f^{g,\text{DY}}(\hat{\tau}), \quad (\text{E.66})$$

where the splitting function

$$P_{q \rightarrow qg}(\hat{\tau}) = \frac{1}{2} [\hat{\tau}^2 + (1 - \hat{\tau})^2], \quad (\text{E.67})$$

and

$$\begin{aligned} \alpha_s f^{g,\text{DY}}(\hat{\tau}) &= \frac{\alpha_s}{2\pi} \frac{1}{2} \left\{ [\hat{\tau}^2 + (1 - \hat{\tau})^2] \log \left(\frac{(1 - \hat{\tau})^2}{\hat{\tau}} \right) - \frac{3}{2} \hat{\tau}^2 + \hat{\tau} + \frac{3}{2} \right\} \\ &+ \frac{\alpha_s}{2\pi} P_{g \rightarrow q\bar{q}}(\hat{\tau}) \left(\frac{2}{\epsilon} + \gamma_E - \log(4\pi) \right). \end{aligned} \quad (\text{E.68})$$

Combining the "annihilation" term with the "Compton" term and including terms with the initial two partons interchanged, then the "Drell-Yan" cross section becomes (for one quark flavor)

$$\begin{aligned} s \frac{d\sigma_{\text{DY}}}{dM^2}(s, M^2) &= \frac{4\pi}{9} \frac{\alpha^2 e_q^2}{M^2} \int_{\tau}^1 \frac{dx_a}{x_a} \int_{\tau/x_a}^1 \frac{dx_b}{x_b} \left\{ \left(\bar{G}_{p \rightarrow q}^{(0)}(x_a) \bar{G}_{p \rightarrow \bar{q}}^{(0)}(x_b) \right. \right. \\ &+ \left. \bar{G}_{p \rightarrow \bar{q}}^{(0)}(x_a) \bar{G}_{p \rightarrow q}^{(0)}(x_b) \right) \left[\frac{\sigma_{\text{tot}}^{\text{DY}}}{\sigma_0} \delta(1 - \hat{\tau}) + \frac{\alpha_s}{2\pi} 2P_{q \rightarrow qg}(\hat{\tau}) \log \left(\frac{M^2}{\Lambda^2} \right) + 2\alpha_s f^{q,\text{DY}}(\hat{\tau}) \right] \\ &+ \left(\bar{G}_{p \rightarrow q}^{(0)}(x_a) \bar{G}_{p \rightarrow g}^{(0)}(x_b) + \bar{G}_{p \rightarrow g}^{(0)}(x_a) \bar{G}_{p \rightarrow q}^{(0)}(x_b) \right) \left[\frac{\alpha_s}{2\pi} P_{g \rightarrow q\bar{q}} \log \left(\frac{M^2}{\Lambda^2} \right) \right. \\ &+ \left. 2\alpha_s f^{g,\text{DY}}(\hat{\tau}) \right] + \left(\bar{G}_{p \rightarrow q}^{(0)}(x_a) \bar{G}_{p \rightarrow g}^{(0)}(x_b) + \bar{G}_{p \rightarrow g}^{(0)}(x_a) \bar{G}_{p \rightarrow q}^{(0)}(x_b) \right) \\ &\times \left. \left[\frac{\alpha_s}{2\pi} P_{g \rightarrow q\bar{q}}(\hat{\tau}) \log \left(\frac{M^2}{\Lambda^2} \right) + 2\alpha_s f^{g,\text{DY}}(\hat{\tau}) \right] \right\}, \end{aligned} \quad (\text{E.69})$$

where $\hat{\tau} = \tau/(x_a x_b)$ and

$$\frac{\sigma_{\text{tot}}^{\text{DY}}}{\sigma_0} = 1 + \alpha_s I_q^{\text{DY}} + \dots, \quad (\text{E.70})$$

with I_q^{DY} is given by eq. (E.61). The "little f " functions in the dimensional regularization scheme is given by eq. (E.64). The $\log(m_D^2)$ divergence has been absorbed into the $G_{p \rightarrow q}^{(0)}$ and $G_{p \rightarrow g}^{(0)}$ structure functions.

BIBLIOGRAPHY

- [1] S. Glashow, “Partial-symmetries of weak interactions”, Nucl. Phys. 22 (1961) 579.
- [2] S. Weinberg, “A Model of Leptons”, Phys. Rev. Lett. 19 (1967) 1264.
- [3] A. Salam, “Elementary Particle Theory”, edited by N. Svartholm (Almqvist and Wiksell, Stockholm, 1968), p.367.
- [4] S. Glashow, J. Iliopoulos, and L. Maiani, “Weak Interactions with Lepton-Hadron Symmetry”, Phys. Rev. D 2 (1970) 1285.
- [5] H. Georgi and S. Glashow, “Unified Weak and Electromagnetic Interactions without Neutral Currents”, Phys. Rev. Lett. 28 (1972) 1494.
- [6] H. D. Politzer, “Reliable Perturbative Results for Strong Interactions?”, Phys. Rev. Lett. 30 (1973) 1346.
- [7] H. D. Politzer, “Asymptotic freedom: An approach to strong interactions”, Phys. Rep. 14 (1974) 129
- [8] D. J. Gross and F. Wilczek, “Ultraviolet Behavior of Non-Abelian Gauge Theories”, Phys. Rev. Lett. 30 (1973) 1343.
- [9] D. J. Gross and F. Wilczek, “Asymptotically Free Gauge Theories. I”, Phys. Rev. D 8 (1973) 3633.
- [10] S. Weinberg, “Non-Abelian Gauge Theories of the Strong Interactions”, Phys. Rev. Lett. 31 (1973) 494.
- [11] S. Weinberg, “Current Algebra and Gauge Theories. II. Non-Abelian Gluons”, Phys. Rev. D 8 (1973) 4482.
- [12] S. Weinberg, “The making of the Standard Model”, Eur. Phys. J. C34 (2004) 5.
- [13] G. 't Hooft, “Renormalizable Lagrangians for massive Yang-Mills fields”, Nucl. Phys. B 33 (1971) 167.
- [14] G. 't Hooft, “Renormalization of massless Yang-Mills fields”, Nucl. Phys. B 33 (1971) 173.
- [15] G. 't Hooft and M. Veltman, “Regularization and renormalization of gauge fields”, Nucl. Phys. B 44 (1972) 189.
- [16] G. 't Hooft and M. Veltman, “Combinatorics of gauge fields”, Nucl. Phys. B 50 (1972) 318.

- [17] H. Fritzsch and P. Minowski, “Flavordynamics of quarks and leptons”, Phys. Rep. 73 (1981) 67.
- [18] B. F. L. Ward, Lecture for Quantum Field Theory 2.
- [19] Chris Quigg, “Gauge Theories of the Strong, Weak, and Electromagnetic Interactions, Frontiers in Physics”, The Benjamin-Cummings Publishing Company, Inc., 1983.
- [20] C. N. Yang and R. Mills, “Conservation of Isotopic Spin and Isotopic Gauge Invariance”, Phys. Rev. 96 (1954) 191.
- [21] Y. Nambu, “Axial Vector Current Conservation in Weak Interactions”, Phys. Rev. Lett. 4 (1960) 380.
- [22] Y. Nambu and G. Jona-Lasinio, “Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I”, Phys. Rev. 122 (1961) 345.
- [23] Y. Nambu and G. Jona-Lasinio, “Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. II”, Phys. Rev. 124 (1961) 246.
- [24] J. Goldstone, A. Salam and S. Weinberg, “Broken Symmetries”, Phys. Rev. 127 (1962) 965.
- [25] S. Bludman and A. Klein, “Broken Symmetries and Massless Particles”, Phys. Rev. 131 (1962) 2363.
- [26] E. S. Abers and B. W. Lee, “Gauge Theories”, Phys. Rep. 9 C (1973) 1.
- [27] F. Englert and R. Brout, “Broken Symmetry and the Mass of Gauge Vector Mesons”, Phys. Rev. Lett. 13 (1964) 9.
- [28] P. W. Higgs, “Broken Symmetries and the Masses of Gauge Bosons”, Phys. Rev. Lett. 13 (1964) 16.
- [29] G. S. Guralnik, C. R. Hagen and T. B. Kibble, “Global Conservation Laws and Massless Particles”, Phys. Rev. Lett. 13 (1964) 20.
- [30] A. Denner, “Techniques for the calculation of electroweak radiative corrections at the one-loop level and results for W-physics at LEP200”, Fortschr. Phys. 41 (1993) 4.
- [31] G. Passarino and M. Veltman, “One-loop corrections for e^+e^- annihilation into $\mu^+\mu^-$ in the Weinberg model”, Nucl. Phys. B 160 (1979) 151.
- [32] D. Bardin and G. Passarino, “The Standard Model in the Making”, Oxford University Press, 1999.
- [33] M. Consoli, “One-loop corrections to $e^+e^- \rightarrow e^+e^-$ in the Weinberg model”, Nucl. Phys. B 160 (1979) 208.

- [34] M. Veltman, “Radiative corrections to vector boson masses”, Phys. Lett. B 91 (1980) 95.
- [35] M. Green and M. Veltman, “Weak and E.M. radiative corrections to low-energy processes”, Nucl. Phys. B 169 (1980) 137; E: Nucl. Phys. B 175 (1980) 547.
- [36] F. A. Berends, R. Kleiss and S. Jadach, “Radiative corrections to muon pair and quark pair production in electron-positron collisions in the Z_0 region”, Nucl. Phys. B 202 (1982) 63.
- [37] I. B. Khriplovich, “Charge asymmetry of muon angular distribution in $e^+e^- \rightarrow \mu^+\mu^-$ process”, Sov. J. Nucl. Phys. 17 (1973) 576.
- [38] V. N. Popov and L.D. Faddeev, “Perturbation theory for gauge invariant fields”, Kiev Inst. Theor. Phys. Acad. Sci. preprint ITP (1967) 67-36.
- [39] L. D. Faddeev and V. N. Popov, “Feynman diagrams for the Yang-Mills field”, Phys. Lett. B 25 (1967) 29.
- [40] A. Sirlin, “Radiative corrections in the $SU(2)_L \times U(1)$ theory: A simple renormalization framework”, Phys. Rev. D 22 (1980) 971.
- [41] W. J. Marciano and A. Sirlin, “Radiative corrections to neutrino-induced neutral-current phenomena in the $SU(2)_L \times U(1)$ theory”, Phys. Rev. D 22 (1980) 2695.
- [42] A. Sirlin and W. J. Marciano, “Radiative corrections to $\nu_\mu + N \rightarrow \mu^- + X$ and their effect on the determination of ρ^2 and $\sin^2 \theta_W$ ”, Nucl. Phys. B 189 (1981) 442.
- [43] G. 't Hooft and M. Veltman, “Scalar one-loop integrals”, Nucl. Phys. B 153 (1979) 365.
- [44] A. Denner, U. Nierste and R. Scharf, “A compact expression for the scalar one-loop four-point function”, Nucl. Phys. B 367 (1991) 637.
- [45] F. A. Berends, R. Kleiss, P. De Causmaecker and R. Gastmans, “Single bremsstrahlung processes in gauge theories”, Phys. Lett. B 103 (1981) 124;
- [46] P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, “Helicity amplitudes for massless QED”, Phys. Lett. B 105 (1981) 215.
- [47] D. Dankaert, P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, “Four-jet production in e^+e^- annihilation”, Phys. Lett. B 114 (1982) 203.
- [48] P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, “Multiple bremsstrahlung in gauge theories at high energies (I). General formalism for quantum electrodynamics”, Nucl. Phys. B 206 (1982) 53.
- [49] F. A. Berends, R. Kleiss, P. De Causmaecker, R. Gastmans, W. Troost and T. T. Wu, “Multiple bremsstrahlung in gauge theories at high energies (II). Single bremsstrahlung”, Nucl. Phys. B 206 (1982) 61.

- [50] F. A. Berends, P. De Causmaecker, R. Gastmans, R. Kleiss, W. Troost and T. T. Wu, (CALKUL Collaboration), “Multiple bremsstrahlung in gauge theories at high energies: (III). Finite mass effects in collinear photon bremsstrahlung”, Nucl. Phys. B 239 (1984) 382.
- [51] Z. Xu, D.-H. Zhang, and L. Change, “Helicity amplitudes for multiple bremsstrahlung in massless non-abelian gauge theories”, Nucl. Phys. B 291 (1987) 392.
- [52] R. Kleiss, and W. J. Stirling, “Spinor techniques for calculating $pp \rightarrow W^\pm/Z^0 + \text{jets}$ ”, Nucl. Phys. B 262 (1985) 235.
- [53] Z. Xu, D.-H. Zhang and L. Change, Tsinghua University preprints TUTP-84/3, 84/4, and 84/5, Unpublished.
- [54] S. Jadach, B. F. L. Ward and Z. Was, “Global positioning of spin GPS scheme for half-spin massive spinors”, Eur. Phys. J. C 22 (2001) 423.
- [55] B. F. L. Ward, “Magic spinor product methods in loop integrals”, Phys. Rev. D 83 (2011) 113014.
- [56] Thomas Hahn, “LoopTools 2.15 User’s Guide”.
- [57] G. J. van Oldenborgh and J. A. M. Vermaseren, “New algorithms for one-loop integrals”, Z. Phys. C 46 (1990) 425.
- [58] S. Jadach, B. F. L. Ward and Z. Was, “Coherent exclusive exponentiation for precision Monte Carlo calculations”, Phys. Rev. D 63 (2001) 113009.
- [59] W. L. van Neerven and J. A. M. Vermaseren, “Large loop integrals”, Phys. Lett. B 137 (1984) 241.
- [60] A. Denner and S. Dittmaier, “Reduction of one-loop tensor 5-point integrals”, Nucl. Phys. B 658 (2003) 175.
- [61] A. Denner and S. Dittmaier, “Reduction schemes for one-loop tensor integrals” Nucl. Phys. B 734 (2006) 62.
- [62] J. C. Taylor, “Ward identities and charge renormalization of the Yang-Mills field”, Nucl. Phys. B 33 (1971) 436.
- [63] A. A. Slanov, “Ward identities in gauge theories”, Theor. Math. Phys. 10 (1973) 99.
- [64] T. Muta, “Foundations of Quantum Chromodynamics, An Introduction to Perturbative Methods in Gauge Theories”, World Scientific Publishing Co. Pte. Ltd., 1998.
- [65] W. Celmaster and R. J. Gonsalves, “Renormalization-prescription dependence of the quantum-chromodynamic coupling constant”, Phys. Rev. D 20 (1979) 1420.

- [66] P. Pascual and R. Tarrach, “Slavnov-Taylor identities in Weinberg’s renormalization scheme”, Nucl. Phys. B 174 (1980) 123.
- [67] G. ’t Hooft, “Dimensional regularization and the renormalization group”, Nucl. Phys. B 61 (1973) 455.
- [68] A. Peterman, “Renormalization group and the deep structure of the proton”, Phys. Rep. 53 (1979) 157.
- [69] G. Altarelli, “Partons in quantum chromodynamics”, Phys. Rep. 81 (1982) 1.
- [70] K. G. Wilson, “Renormalization Group and Strong Interactions”, Phys. Rev. D 3 (1971) 1818.
- [71] M. Gell-mann and F. E. Low, “Quantum Electrodynamics at Small Distances”, Phys. Rev. 95 (1954) 1300.
- [72] C. G. Callan, Jr., “Broken Scale Invariance in Scalar Field Theory”, Phys. Rev. D 2 (1970) 1541.
- [73] K. Symanzik, “Small distance behaviour in field theory and power counting”, Comm. Math. Phys. 18 (1970) 227.
- [74] S. Weinberg, “New Approach to the Renormalization Group”, Phys. Rev. D 8 (1973) 3497.
- [75] W. Caswell, “Asymptotic Behavior of Non-Abelian Gauge Theories to Two-Loop Order”, Phys. Rev. Lett. 33 (1974) 244.
- [76] D. R. T. Jones, “Two-loop diagrams in Yang-Mills theory”, Nucl. Phys. B 75 (1974) 531.
- [77] F. Block and A. Nordsieck, “Note on the Radiation Field of the Electron”, Phys. Rev, 52, 54 (1937).
- [78] D. R. Yennie, S. C. Frautschi and H. Suura, “The infrared divergence phenomena and high-energy processes. Annals of Physics. 13, 379-452 (1961).
- [79] B. F. L. Ward, “Renormalization-group-improved Yennie-Frautschi-Suura theory”, Phys. Rev. D 36, (1987) 939.
- [80] B. F. L. Ward, The lecture note for Quantum Field Theory 3.
- [81] J. M. Jauch and F. Rohrlich, Helv. Phys. Acta 27, 613 (1954).
- [82] D. R. Yennie and H. Suura, “Higher Order Radiative Corrections to Electron Scattering”, Phys. Rev. 105, (1957) 1378.
- [83] B. F. L. Ward, “Resummed Quantum Gravity”, arXiv:0610232.
- [84] S. Jadach and Z. Was, “Monte Carlo Simulation of the Process $e^+e^- \rightarrow \tau^+\tau^-$ Including Radiative $O(\alpha^3)$ QED Corrections, Mass and Spin”, Comput. Phys. Commun. 36 (1985) 191.

- [85] S. Jadach, B.F.L. Ward and Z. Was, “The Monte Carlo program KORALZ version 4.0 for lepton or quark pair production at LEP/SLC energies”, *Comput. Phys. Commun.* 79(1994) 503.
- [86] S. Jadach, E. Richter-Was, B.F.L. Ward, Z. Was, “Monte Carlo program BHLUMI 2.01 for Bhabha scattering at low angles with Yennie-Frautschi-Suura exponentiation”, *Comput. Phys. Commun.* 70(1992) 305.
- [87] S. Jadach, W. Placzek, E. Richter-Was, B. F. L. Ward, and Z. Was, “Upgrade of the Monte Carlo program BHLUMI for Bhabha scattering at low angles to version 4.04”, *Comput. Phys. Commun.* 102(1997) 229.
- [88] S. Jadach and B.F.L. Ward, “YFS2— The second-order Monte Carlo program for fermion pair production at LEP/SLC, with the initial state radiation of two hard and multiple soft photons”, *Comput. Phys. Commun.* 56(1990) 351.
- [89] S. Jadach, B.F.L. Ward, Z. Was, “The precision Monte Carlo event generator KK for two-fermion final states in e^+e^- collisions”, *Comput. Phys. Commun.* 130(2000) 260.
- [90] S. Jadach, B.F.L. Ward, Z. Was and S. Yost, *Phys. Rev.* D94, 074006.
- [91] S. Jadach, B.F.L. Ward, Z. Was and S. Yost “KKMC-hh: A Precision Event Generator for EW Radiative Corrections in Hadron Scattering”, [hep-ph/1811.03560](https://arxiv.org/abs/hep-ph/1811.03560).
- [92] S. Jadach, B.F.L. Ward, Z. Was and S. Yost, [hep-ph/1801.09509](https://arxiv.org/abs/hep-ph/1801.09509).
- [93] S.D. Drell and T.M. Yan, “Massive Lepton-Pair Production in Hadron-Hadron Collisions at High Energies”, *Phys. Rev. Lett.* 25, 316(1970).
- [94] S. Jadach, B.F.L. Ward, Z. Was, “The Monte Carlo program KORALZ version 4.0 for lepton or quark pair production at LEP/SLC energies”, *Comput. Phys. Commun.* 79(1994) 503.
- [95] S. Jadach, B.F.L. Ward and Z. Was, “Coherent exclusive exponentiation CEEEX: the case of the resonant e^+e^- collision”, *Phys.Lett.* B449(1999) 97
- [96] S. Jadach, B.F.L. Ward and Z. Was, “KK MC 4.22: Coherent exclusive exponentiation of electroweak corrections for $f\bar{f} \rightarrow f'\bar{f}'$ at the LHC and muon colliders”, *Phys.Rev. D* 88 (2013) 114022.
- [97] D. Bardin et al., “ZFITTER v.6.21: A semi-analytical program for fermion pair production in e^+e^- annihilation”, *Comput. Phys. Commun.* 133(2001) 229.
- [98] S. Jadach, J.H. Kühn and Z. Was, “TAUOLA - a library of Monte Carlo programs to simulate decays of polarized τ leptons”, *Comput. Phys. Commun.* 64(1990) 275.
- [99] S. Jadach, Z. Was, R. Decker and J.H. Kühn, “The tau decay library TAUOLA: Version 2.4”, *Comput. Phys. Commun.* 76 (1993) 361.

- [100] P. Golonka, B. Kersevan, T. Pierzchala, E. Richter-Was, Z. Was and M. Worek, “The tauola-photos-F environment for the TAUOLA and PHOTOS packages, release II”, *Comput. Phys. Commun.* 174 (2006) 818.
- [101] A. Buckley, James Ferrando, Stephen Lloyd, Karl Nordström, Ben Page, Martin Rufenacht, Marek Schönherr and Graeme Watt, “LHAPDF6: parton density access in the LHC precision era”, *Eur. phys. J. C* 75(2015) 3, 132.
- [102] S. Jadach, “Foam: Multi-dimensional general purpose Monte Carlo generator with self-adapting simplicial grid”, *Comput. Phys. Commun.* 130 (2000) 244.
- [103] F.A. Berends and R. Kleiss, “Distributions in the process $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$ ”, *Nucl. Phys. B* 177 (1981) 237 .
- [104] F.A. Berends, R. Kleiss and S. Jadach, “Monte Carlo stimulation of radiative corrections to the processes $e^+e^- \rightarrow \mu^+\mu^-$ and $e^+e^- \rightarrow \bar{q}q$ in the Z_0 region”, *Comput. Phys. Commun.* 29 (1983) 185.
- [105] F.A. Berends, R. Kleiss and S. Jadach, “Radiative corrections to muon pair and quark pair production in electron-positron collisions in the Z_0 region”, *Nucl. Phys. B* 202 (1982) 63.
- [106] F.A. Berends, W. Van Neerven, and G. Burgers, “Higher order radiative corrections at LEP energies”, *Nucl. Phys. B* 297 (1988) 429.
- [107] S. Jadach, B.F.L. Ward and Z. Was, “Coherent exclusive exponentiation CEEEX: the case of the resonant e^+e^- collision”, *Phys. Lett. B* 449 (1999) 97.
- [108] M. Greco, G. Pancheri-Srivastava and Y. Srivastava, “Radiative corrections for colliding beam resonances”, *Nucl. Phys. B* 101 (1975) 234.
- [109] M. Greco, G. Pancheri-Srivastava and Y. Srivastava, “Radiative corrections to $e^+e^- \rightarrow \mu^+\mu^-$ around the Z_0 ”, *Nucl. Phys. B* 171 (1980) 118;B 197 (1982) 543E.
- [110] Z. Was, “Radiative corrections to τ pair production around Z_0 ”, *Acta Phys. Pol. B* 18 (1987) 1099.
- [111] R. W. Brown, R. Decker and E. A. Paschos, “Weak Corrections to the $e^+e^- \rightarrow \mu^+\mu^-$ Asymmetry”, *Phys. Rev. Lett.* 52 (1984) 1192.
- [112] S. Jadach and Z. Was, “Monte Carlo simulation of the process $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^\pm \rightarrow X^\pm$ including radiative $O(\alpha^3)$ QED corrections, mass and spin effects”, *Comput. Phys. Commun.* 36 (1985) 191.
- [113] G. Burgers, “On the two-loop QED vertex correction in the high energy limit”, *Phys. Lett.* 164 B (1985) 167.
- [114] F. Berends, G. Burgers and W. Van Neerven, “QED radiative corrections to the reaction $e^+e^- \rightarrow Z\gamma$ ”, *Phys. Lett. B* 177 (1986) 191.
- [115] S. Jadach, “Practical guide to Monte Carlo”, 1999, hep-ph/9906056.

- [116] S. Jadach, “Comoving reference frames for multi-bremsstrahlung”, To be submitted to *Comput. Phys. Commun.*
- [117] S. Jadach, “Yennie–Frautschi–Suura soft photons in Monte Carlo event generators”, 1987, MPI-PAE/PTh 6/87, preprint of MPI Munich, unpublished.
- [118] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP*, 7 (2014) 79.
- [119] T. Stelzer and W. F. Long, “Automatic generation of tree level helicity amplitudes”, *Comput. Phys. Commun.* 81 (1994) 357.
- [120] H. Murayama, I. Watanabe, and K. Hagiwara, “HELAS: HELicity amplitude subroutines for Feynman diagram evaluations”.
- [121] C. Duhr and B. Fuks, “A superspace module for the FeynRules package”, arXiv:1102.4191.
- [122] C. Degrande et al., “UFO — The Universal FeynRules Output”.
- [123] T. Gleisberg and F. Krauss, “Automating dipole subtraction for QCD NLO calculations”, *Eur. Phys. J. C* 53 (2008) 501.
- [124] M.H. Seymour and C. Tevlin, “TeVJet: a general framework for the calculation of jet observables in NLO QCD”, arXiv:0803.2231.
- [125] Hasegawa, S. Moch and P. Uwer, “Automating dipole subtraction”, *Nucl. Phys. Proc. Suppl.* 183 (2008) 268.
- [126] R. Frederix, T. Gehrmann and N. Greiner, “Automation of the dipole subtraction method in MadGraph/MadEvent”, *JHEP* 09 (2008) 122.
- [127] C.F. Berger et al., “Precise predictions for $W + 4$ jet production at the Large Hadron Collider”, *Phys. Rev. Lett.* 106 (2011) 092001.
- [128] G. Ossola, C.G. Papadopoulos and R. Pittau, “CutTools: a program implementing the OPP reduction method to compute one-loop amplitudes”, *JHEP* 03 (2008) 042.
- [129] E. Boos et al, “Generic user process interface for event generations”, hep-ph/0109068
- [130] M. Aaboud et al. (ATLAS Collaboration), “Measurement of the W-boson mass in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, *Eur. Phys. J. C* 78 (2018) 110.
- [131] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, “Parton distributions incorporating QED contribution”, *Eur. Phys. J. C* 39 (2005) 155.

- [132] M. Gell-Mann, “Symmetries of Baryons and Mesons” , Phys. Rev. 125 (1962) 1067.
- [133] R. D. Field, “Applications of Perturbative QCD”, 1989